

The Reformulated Asymptotic Freedom

Sylwester Kornowski

Abstract: Within the Everlasting Theory I calculated the running coupling for the strong interactions applying three different methods. At very high energy there appears asymptote for 0.1139. The derivations are free from the methods characteristic for the Quantum Theory of Fields i.e. from the infinities, singularities, arbitrary assumptions, minimal subtraction, sliding scale and absolute parameter taken from experiment. We can make use of the prediction of existence of the asymptote to test the Everlasting Theory.

1. Introduction

Many experimental results lead to conclusion that inside baryons is a core. The Everlasting Theory [1] shows that due to the phase transitions of the modified Higgs field there appear the torus and ball in its centre both composed of the Einstein spacetime components. The torus is the black hole in respect of the strong interactions whereas the ball is the black hole in respect of the weak interactions.

Define energy of collision per nucleon as $E_N[\text{GeV}] = nm_N = m_N/\beta$ i.e. $\beta = m_N/E_N$, where $m_N = 0.939$ GeV. Then, the two formulae derived in the Everlasting Theory [1] for the upper and lower limits for the running coupling for the strong-weak interactions we can rewrite as follows:

$$\alpha_{sw} = \alpha_{sw,\text{central-value}} \pm \Delta\alpha_{sw} , \quad (1)$$

$$\alpha_{sw} = \{ \alpha_{w(\text{proton})}\beta^2 + b\beta + c \} \pm (b - b_1)\beta, \quad (2)$$

$$\alpha_{w(\text{proton})} = 0.0187229,$$

$$b = 0.36255,$$

$$c = 0.1139,$$

$$b - b_1 = 0.04415.$$

We can see that at very high energy there is asymptote $\alpha_{sw} = c = 0.1139$. The formula (2) follows from the internal structure of the core of baryons (the torus + ball), the law of conservation of spin and the uncertainty principle.

For $E_N \rightarrow \infty$ is $\alpha_{sw} = 0.1139$, for $E_N = 2.76$ GeV is 0.1140, for 91.19 GeV is 0.1176 ± 0.0005 , for 20 GeV is 0.1309 ± 0.0021 , for 10 GeV is 0.1481 ± 0.0041 , for 5 GeV is 0.1827 ± 0.0083 , for 2 GeV is 0.2882 ± 0.0207 and for 1 GeV is 0.4708 ± 0.0415 .

The running coupling in 1-, 2-, 3- and 4-loop approximation for QCD scale $\Lambda = 0.22$ GeV and $\alpha_s(M_Z) = 0.119$ is for $E_N = 2.76$ GeV in approximation 0.08 [2]. This value differs very much from the result obtained from formula (2) i.e. 0.114. We can use this significant difference to test the Everlasting Theory.

2. Calculations

On surface of the torus inside the core of baryons appear the gluon balls. Their energy $M[\text{GeV}]$ we can calculate from formulae (214)-(216) presented here [1]. We can rewrite these formulae as follows

$$M[\text{GeV}] = (C/E_N[\text{GeV}] + D)^{10}, \quad (3)$$

$$C = 0.52294,$$

$$D = 0.96868.$$

Calculate following derivative $\partial M/\partial E_N$

$$\partial M/\partial E_N = -10C(C/E_N + D)^9/E_N^2, \quad (4)$$

For $C/E_N \ll D$ i.e. for $E_N \gg C/D = 0.54$, we obtain

$$\partial M/\partial E_N = -F/E_N^2, \quad (5)$$

$$F = 3.927.$$

The value F is for the strongly interacting torus which mass is $X = 318.3 \text{ MeV}$ [1]. This torus produces gluons which energy m_g is the one fourth of the mass of the bound neutral pion (134.9661 MeV) [1]. The change of the X onto m_g changes the value of the F : $F' = Fm_g/X \approx 0.42$. So, we can rewrite the formula (5) as follows

$$\partial M/\partial E_N = -F'/E_N^2, \quad (6)$$

$$F' = 0.42.$$

Integrate the equation (6). There appears the integration constant H which we can interpret as the ratio of the mass of the torus X to the mass of nucleon m_N : $H = X/m_N$. We obtain

$$M = F'/E_N + H, \quad (7)$$

$$H = 0.339.$$

We can define running coupling α_{sw} as follows $\alpha_{sw} = M^2$. It leads to following formula

$$\alpha_{sw} = (F'/E_N + H)^2, \quad (8)$$

For $E_N \rightarrow \infty$ is $\alpha_{sw} = 0.115$, for $E_N = 2.76 \text{ GeV}$ is 0.115 , for 91.19 GeV is 0.118 , for 20 GeV is 0.13 , for 10 GeV is 0.15 , for 5 GeV is 0.18 , for 2 GeV is 0.30 and for 1 GeV is 0.58 . We can see that only for 1 GeV the obtained result from formula (8) is not close to the central value obtained from formula (2).

The formula (8) we can derive in different way. The gluon balls produced from the energy of collision of a nucleon look similarly to the ball in the centre of torus. Due to the confinement [1], the all gluon balls have the same radius as the ball in centre of torus. When energy of a gluon ball increases then increases the rotational energy of the carriers of the gluons the ball consists of. This energy does not increase mass density of the ball so it still is the black hole in respect of the weak interactions. For a rotating gluon ball is $E_b v r = \text{const}$. i.e. $E_b \sim 1/v$, where v is the spin speed of the rotating components of the Einstein spacetime i.e. the carriers of gluons. The relative mass R_{sw} responsible for asymptotic freedom, we can separate into two parts. The first part concerns the strong interactions of the torus that is the black hole in respect of the strong interactions. The R_{strong} should be $R_{strong} = X/m_N$, where $X = 0.3183 \text{ GeV}$ is the rest mass of the torus. The second part concerns the weak interactions of the created gluon balls. They are the black holes in respect of the weak interactions. Due to the relation $E_b \sim 1/v$, the intensity of weak interactions of the gluon balls decreases when energy increases. It leads to conclusion that the second part R_{weak} should be $R_{weak} = Y/E_N$, where $Y = 0.4241 \text{ GeV}$ is the rest mass of the ball in centre of the torus.

The above description leads to following formula

$$R_{sw} = R_{weak} + R_{strong} = Y/E_N + X/m_N. \quad (9)$$

Similar as previously we can define the running coupling as R^2 so we obtain

$$\alpha_{sw} = (Y/E_N + X/m_N)^2. \quad (10)$$

This formula is correct for $E_N \gg E_o = Ym_N/X = 1.25 \text{ GeV}$. Due to the quadrupole symmetry for the weak interactions, the lower limit for the E_N should be $4E_o = 5 \text{ GeV}$ and such value is in the old asymptotic freedom.

3. Summary

I described the asymptotic freedom applying three different methods. They lead to the same or very close theoretical results. The theoretical results are consistent or very close to experimental data. This means that experimental results lead to the core of baryons composed of the torus and ball in its centre. Within the Everlasting Theory I derived the internal structure of the core of baryons on base of the phase transitions of the modified Higgs field which I refer to as the Newtonian spacetime. This spacetime consists of internally structureless tachyons that carry the inertial mass only. The modified Higgs field is gravitationally the massless field. Moreover, the mean spin of the tachyons is in approximation 10^{67} times lower than the reduced Planck constant so the modified Higgs field is practically the scalar field.

The old QCD [3], [4] should be inconsistent with experimental data at very high energies. It follows from the fact that the Quantum Theory of Fields neglects the internal structure of the bare fermions (i.e. the torus + ball) and neglects the atom-like structure of baryons [1].

I described the origin of the mass scale 5 GeV which appears in the old asymptotic freedom.

The new asymptotic freedom is free from the methods characteristic for the Quantum Theory of Fields i.e. from the infinities, singularities, arbitrary assumptions, minimal subtraction, sliding scale and absolute parameter taken from experiment.

We can make use of the prediction of existence of the asymptote 0.1139 to test the Everlasting Theory.

References

- [1] S. Kornowski (2012). "The Everlasting Theory and Special Number Theory".
<http://www.rxiv.org/abs/1203.0021> [v2].
- [2] S. Bethke (2008). "Experimental Tests of Asymptotic Freedom".
<http://arxiv.org/pdf/hep-ex/0606035v2.pdf> , p. 12.
- [3] D.J. Gross, F. Wilczek (1973). "Ultraviolet behaviour of non-abelian gauge theories".
Physical Review Letters **30** (26).
- [4] H.D. Politzer (1973). "Reliable perturbative results for strong interactions". *Physical Review Letters* **30** (26).