

Author Michael John Sarnowski; Email thiele@charter.net

I. Abstract

The calculation of the Planck constant has been elusive. Here it is shown, thanks to the Rydberg constant and the equation for force of charge and gravity. The equation shows that the fundamental constants are likely from a very complex yet consistent action of the aether.

II. Calculations

It is known that the Rydberg infinity constant is the most precise physical measurement in physics and that it can also be calculated from fundamental constants and the mass of the electron. It's value is shown below

Equation 1.0
$$R = \frac{Meq^4}{8\varepsilon^2 h^3 c} = 1.0973731568539 * 10^7 m^{-1}$$

We found from "Discrete Calculations of Charge and Gravity with Planck Spinning Spheres and Kaluza Spinning Spheres", Michael John Sarnowski (2) that the following equation for q, $q^2 = T\pi^3 hc\varepsilon(Me) / 2Mn$ modeled Discrete unit charge. We can rearrange the equation for charge and the equation for the Rydberg constant as follows.

Equation 2.0
$$\frac{q^4}{\varepsilon^2} = \frac{8Rh^3c}{Me}$$

Equation 3.0
$$\frac{q^2}{\varepsilon} = \frac{T\pi^3 hcMe}{2Mn}$$

Equation 3.0 can then be squared and equalized and then solved for h which yields

Equation 4.0
$$h = \frac{T^2 \pi^6 c Me^3}{32Mn^2 R} = 6.62606935 * 10^{-34} \text{ joulesec}$$

This compares to the codata value of $6.62606957(30) * 10^{-34} \text{ joulesec}$

The value from equation 4.0 is within one standard deviation of the Codata value.

Note the following

Author Michael John Sarnowski; Email thiele@charter.net

$$1) \quad T^2 = \frac{(M_p - M_e)^2 + M_n^2 + M_n^2}{M_n^2}$$

2) Where h =planck's constant, c =speed of light in a vacuum, G =gravitational constant, and M_n =rest mass of the neutron, M_p =rest mass of the proton, and M_e =rest mass of the electron and R =Rydberg constant.

Appendix A

Fundamental Physical Constants (18)

1. $c=2.99792458 \times 10^8$ m/s
2. $h=6.626\ 06957(33) \times 10^{-34}$ J s
3. Mass of Neutron = $M_n= 1.674\ 927\ 351(74) \times 10^{-27}$ kg
4. Mass of Proton = $M_p= 1.672\ 621\ 777(74) \times 10^{-27}$ kg
5. Mass of Electron = $M_e = 9.109\ 382\ 91(40) \times 10^{-31}$ kg.
6. q = unit charge = $1.602\ 176\ 565(35) \times 10^{-19}$ C
7. ϵ = Dielectric Permittivity = $8.854187817 \times 10^{-12}$
8. $G= 6.67384(80) \times 10^{-11}$ m³ kg⁻¹ s⁻²

9. $T= 1.730942781$ where

$$T^2 = \frac{((M_p - M_e)^2 + M_n^2 + M_n^2)}{M_n^2}$$

$$10.0 \quad R = \frac{M_e q^4}{8 \epsilon^2 h^3 c} = 1.0973731568539 \times 10^7 \text{ m}^{-1}$$

Definitions

References

- 1) http://en.wikipedia.org/wiki/Rydberg_constant
- 2) <http://www.vixra.org/pdf/1403.0502v5.pdf>
- 3) <http://www.vixra.org/pdf/1404.0055v2.pdf>
- 4) <http://vixra.org/pdf/1404.0035v1.pdf>
- 5) http://en.wikipedia.org/wiki/Gravitational_coupling_constant

Calculation of the Planck Constant and Gravitational constant

Author Michael John Sarnowski; Email thiel@charter.net