

Certain Single-Valued Neutrosophic Graphs

Muhammad Akram and Saba Siddique

Department of Mathematics, University of the Punjab,

New Campus, Lahore, Pakistan.

E-mail: m.akram@pucit.edu.pk, sabasabiha6@gmail.com

Abstract

Neutrosophic sets are the generalization of the concept of fuzzy sets and intuitionistic fuzzy sets. Neutrosophic models give more flexibility, precisions and compatibility to the system as compared to the classical, fuzzy and intuitionistic fuzzy models. In this research paper, we present certain types of single-valued neutrosophic graphs, including regular single-valued neutrosophic graphs, totally regular single-valued neutrosophic graphs, edge regular single-valued neutrosophic graphs and totally edge regular single-valued neutrosophic graphs. We also investigate some of their related properties.

Keywords: Edge regular single-valued neutrosophic graphs, Totally edge regular single-valued neutrosophic graphs.

Mathematics Subject Classification 2010: 03E72, 68R10, 68R05

1 Introduction

Fuzzy set theory is the generalized concept of classical set theory. In classical set theory, there are only two possibilities, that is, either the statement is true or false. However, many statements have variable values which can be handled more accurately using fuzzy set theory. In 1965, Zadeh [20] introduced the notion of fuzzy sets to handle the problems with uncertainties. Fuzzy set theory [20] plays a vital role in complex phenomena which is not easily characterized by classical set theory. In 1983, Atanassov [6] proposed the notion of intuitionistic fuzzy sets as a generalization of fuzzy sets. He added a new component which determines the falsity membership degree in the definition of fuzzy sets. The idea of intuitionistic fuzzy sets is more meaningful as well as intensive due to the presence of truth membership degree, indeterminacy membership degree and falsity membership degree, where the indeterminacy membership degree of intuitionistic fuzzy sets is its hesitation part by default. The truth membership degree and the falsity membership degree are more or less independent from each other, the only requirement is that the sum of these two degrees is not greater than one. Smarandache [9-10] introduced the idea of neutrosophic sets by combining the non-standard analysis. Neutrosophic set is a mathematical tool for dealing real life problems having imprecise, indeterminacy and inconsistent data. Neutrosophic set theory, as a generalization of classical set theory, fuzzy set theory and intuitionistic fuzzy set theory, is applied in a variety of fields, including control theory, decision making problems, topology, medicines and in many more real life problems. Wang et al. [19] presented the notion of single-valued neutrosophic sets to apply neutrosophic sets in real life problems more conveniently. A single-valued neutrosophic set has three components: truth membership degree, indeterminacy membership degree and falsity membership degree. These three components of a single-valued neutrosophic set are not dependent and their values are contained in the standard unit interval $[0, 1]$. Single-valued neutrosophic sets are the generalization of intuitionistic fuzzy sets. Single-valued neutrosophic sets have been a new hot research topic and many researchers have addressed this issue. Majumdar and Samanta [11] studied similarity and entropy of single-valued neutrosophic sets.

Graph theory has become a powerful conceptual framework for modeling and solution of combinatorial problems that arise in various areas, including computer sciences, engineering and mathematics. Single-valued neutrosophic graphs, as the generalization of graphs, have many properties which are the basis of different techniques that are used in modern mathematics. Dhavaseelan et al. [10] defined strong neutrosophic graphs. Broumi et al. [8, 9] portrayed single-valued neutrosophic graphs. Akram and Shahzadi [1] introduced the notion of neutrosophic soft graphs with applications. Akram and Shahzadi [4] highlighted some flaws in the definitions of Broumi et al. [8] and Shah-Hussain

[16]. Akram [3] introduced the notion of single-valued neutrosophic planar graphs. Akram et al. [2] also introduced the single-valued neutrosophic hypergraphs. Representation of graphs using intuitionistic neutrosophic soft sets was discussed in [5]. In this research paper, we present certain types of single-valued neutrosophic graphs, including regular single-valued neutrosophic graphs, totally regular single-valued neutrosophic graphs, edge regular single-valued neutrosophic graphs and totally edge regular single-valued neutrosophic graphs.

2 Single-valued neutrosophic graphs

Definition 2.1. [1] A single-valued neutrosophic graph $G = (X, Y)$ is a pair, where $X : N \rightarrow [0, 1]$ is a single-valued neutrosophic set on N and $Y : N \times N \rightarrow [0, 1]$ is a single-valued neutrosophic relation on N such that

$$\begin{aligned} t_Y(st) &\leq \min\{t_X(s), t_X(t)\}, \\ i_Y(st) &\leq \min\{i_X(s), i_X(t)\}, \\ f_Y(st) &\leq \max\{f_X(s), f_X(t)\}, \end{aligned}$$

for all $s, t \in N$, where $t_Y(st) = 0$, $i_Y(st) = 0$ and $f_Y(st) = 0$, for all $s, t \in N \times N - L$. X and Y are called the single-valued neutrosophic vertex set of G and the single-valued neutrosophic edge set of G , respectively. A single-valued neutrosophic relation Y is said to be symmetric if $t_Y(st) = t_Y(ts)$, $i_Y(st) = i_Y(ts)$ and $f_Y(st) = f_Y(ts)$, for all $s, t \in N$.

Example 2.1. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4, s_5\}$ and $L = \{s_1s_2, s_1s_4, s_1s_5, s_2s_3, s_2s_4, s_3s_4, s_4s_5\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 1.




Figure 1: Single-valued neutrosophic graph G

Definition 2.2. A single-valued neutrosophic path \mathcal{P} is a sequence of distinct vertices $s = s_1, s_2, s_3, \dots, s_n = t$ such that, for all k , $t_Y(s_k s_{k+1}) > 0$, $i_Y(s_k s_{k+1}) > 0$ and $f_Y(s_k s_{k+1}) > 0$. A single-valued neutrosophic path is said to be a single-valued neutrosophic cycle if $s = t$.

Example 2.2. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4, s_5\}$ and $L = \{s_1s_5, s_2s_3, s_2s_4, s_3s_4, s_4s_5\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 2.




Figure 2: Single-valued neutrosophic graph G

The path \mathcal{P} from s_2 to s_1 is shown with thick lines and the cycle \mathcal{C} from s_2 to s_2 is shown with dashed lines in Fig. 2.

Definition 2.3. [9] The order and the size of a single-valued neutrosophic graph G are denoted by $\mathcal{O}(G)$ and $\mathcal{S}(G)$, respectively, and are defined as

$$\begin{aligned}\mathcal{O}(G) &= \left(\sum_{s \in N} t_X(s), \sum_{s \in N} i_X(s), \sum_{s \in N} f_X(s) \right), \\ \mathcal{S}(G) &= \left(\sum_{st \in L} t_Y(st), \sum_{st \in L} i_Y(st), \sum_{st \in L} f_Y(st) \right).\end{aligned}$$

Definition 2.4. [9] The degree and the total degree of a vertex s of a single-valued neutrosophic graph G are denoted by $\mathcal{D}_G(s) = (\mathcal{D}_t(s), \mathcal{D}_i(s), \mathcal{D}_f(s))$ and $\mathcal{TD}_G(s) = (\mathcal{TD}_t(s), \mathcal{TD}_i(s), \mathcal{TD}_f(s))$, respectively, and are defined as

$$\begin{aligned}\mathcal{D}_G(s) &= \left(\sum_{s \neq t} t_Y(st), \sum_{s \neq t} i_Y(st), \sum_{s \neq t} f_Y(st) \right), \\ \mathcal{TD}_G(s) &= \left(\sum_{s \neq t} t_Y(st) + t_X(s), \sum_{s \neq t} i_Y(st) + i_X(s), \sum_{s \neq t} f_Y(st) + f_X(s) \right),\end{aligned}$$

for $st \in L$, where $s \in N$.

Example 2.3. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3\}$ and $L = \{s_1s_2, s_2s_3, s_3s_1\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 3.




Figure 3: Single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{O}(G) = (1.5, 1.7, 1.6)$, $\mathcal{S}(G) = (0.9, 0.9, 1.7)$, $\mathcal{D}_G(s_1) = (0.5, 0.6, 1.1)$, $\mathcal{D}_G(s_2) = (0.7, 0.5, 1.2)$, $\mathcal{D}_G(s_3) = (0.6, 0.7, 1.1)$, $\mathcal{TD}_G(s_1) = (0.9, 1.2, 1.6)$, $\mathcal{TD}_G(s_2) = (1.2, 0.9, 1.9)$ and $\mathcal{TD}_G(s_3) = (1.2, 1.4, 1.5)$.

Definition 2.5. A single-valued neutrosophic graph $G = (X, Y)$ is called a regular single-valued neutrosophic graph of degree (m_1, m_2, m_3) if $\mathcal{D}_G(s) = (m_1, m_2, m_3)$, for all $s \in N$.

Example 2.4. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4\}$ and $L = \{s_1s_2, s_2s_3, s_3s_4, s_4s_1\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 4.




Figure 4: Regular single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.2, 1.2, 0.8) = \mathcal{D}_G(s_2) = \mathcal{D}_G(s_3) = \mathcal{D}_G(s_4)$. Hence G is a regular single-valued neutrosophic graph.

Definition 2.6. A single-valued neutrosophic graph $G = (X, Y)$ is called a totally regular single-valued neutrosophic graph of degree (n_1, n_2, n_3) if $\mathcal{TD}_G(s) = (n_1, n_2, n_3)$, for all $s \in N$.

Example 2.5. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4, s_5, s_6\}$ and $L = \{s_1s_2, s_2s_3, s_3s_4, s_4s_5, s_5s_6, s_6s_1\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 5.




Figure 5: Totally regular single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.3, 0.5, 1.1) = \mathcal{D}_G(s_2) = \mathcal{D}_G(s_3) = \mathcal{D}_G(s_4) = \mathcal{D}_G(s_5) = \mathcal{D}_G(s_6)$ and $\mathcal{TD}_G(s_1) = (0.6, 0.9, 1.7) = \mathcal{TD}_G(s_2) = \mathcal{TD}_G(s_3) = \mathcal{TD}_G(s_4) = \mathcal{TD}_G(s_5) = \mathcal{TD}_G(s_6)$. Hence G is a totally regular single-valued neutrosophic graph.

Remark 2.1. The above two concepts are independent, that is, it is not necessary that totally regular single-valued neutrosophic graph is regular single-valued neutrosophic graph and vice versa.

Example 2.6. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4\}$ and $L = \{s_1s_2, s_2s_3, s_3s_4, s_4s_1\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 6.




Figure 6: Totally regular single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.5, 0.7, 0.7)$, $\mathcal{D}_G(s_2) = (0.3, 0.5, 0.5)$, $\mathcal{D}_G(s_3) = (0.2, 0.4, 0.5)$, $\mathcal{D}_G(s_4) = (0.4, 0.6, 0.7)$ and $\mathcal{TD}_G(s_1) = (1.2, 1.1, 1.3) = \mathcal{TD}_G(s_2) = \mathcal{TD}_G(s_3) = \mathcal{TD}_G(s_4)$. Therefore, G is a totally regular single-valued neutrosophic graph but not a regular single-valued neutrosophic graph.

Definition 2.7. The degree and the total degree of an edge st of a single-valued neutrosophic graph G are denoted by $\mathcal{D}_G(st) = (\mathcal{D}_t(st), \mathcal{D}_i(st), \mathcal{D}_f(st))$ and $\mathcal{TD}_G(st) = (\mathcal{TD}_t(st), \mathcal{TD}_i(st), \mathcal{TD}_f(st))$, respectively, and are defined as

$$\mathcal{D}_G(st) = \mathcal{D}_G(s) + \mathcal{D}_G(t) - 2(t_Y(st), i_Y(st), f_Y(st)),$$

$$\mathcal{TD}_G(st) = \mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st)).$$

Example 2.7. Consider a crisp graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3\}$ and $L = \{s_1s_2, s_1s_3\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 7.




Figure 7: Single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.5, 0.2, 1.3)$, $\mathcal{D}_G(s_2) = (0.3, 0.1, 0.6)$, and $\mathcal{D}_G(s_3) = (0.2, 0.1, 0.7)$.

- The degree of each edge is given as:

$$\begin{aligned} \mathcal{D}_G(s_1s_2) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_2) - 2(t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (0.7, 0.5, 0.8) + (0.5, 0.4, 0.7) - 2(0.3, 0.1, 0.6), \\ &= (0.2, 0.1, 0.7). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_1s_3) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_3) - 2(t_Y(s_1s_3), i_Y(s_1s_3), f_Y(s_1s_3)), \\ &= (0.7, 0.5, 0.8) + (0.4, 0.2, 0.6) - 2(0.2, 0.1, 0.7), \\ &= (0.3, 0.1, 0.6). \end{aligned}$$

- The total degree of each edge is given as:

$$\begin{aligned} \mathcal{TD}_G(s_1s_2) &= \mathcal{D}_G(s_1s_2) + (t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (0.2, 0.1, 0.7) + (0.3, 0.1, 0.6), \\ &= (0.5, 0.2, 1.3). \end{aligned}$$

$$\begin{aligned} \mathcal{TD}_G(s_1s_3) &= \mathcal{D}_G(s_1s_3) + (t_Y(s_1s_3), i_Y(s_1s_3), f_Y(s_1s_3)), \\ &= (0.3, 0.1, 0.6) + (0.2, 0.1, 0.7), \\ &= (0.5, 0.2, 1.3). \end{aligned}$$

Definition 2.8. The maximum degree of a single-valued neutrosophic graph G is defined as $\Delta(G) = (\Delta_t(G), \Delta_i(G), \Delta_f(G))$, where

$$\Delta_t(G) = \max\{\mathcal{D}_t(s) : s \in N\},$$

$$\Delta_i(G) = \max\{\mathcal{D}_i(s) : s \in N\},$$

$$\Delta_f(G) = \max\{\mathcal{D}_f(s) : s \in N\}.$$

Definition 2.9. The minimum degree of a single-valued neutrosophic graph G is defined as $\delta(G) = (\delta_t(G), \delta_i(G), \delta_f(G))$, where

$$\delta_t(G) = \min\{\mathcal{D}_t(s) : s \in N\},$$

$$\delta_i(G) = \min\{\mathcal{D}_i(s) : s \in N\},$$

$$\delta_f(G) = \min\{\mathcal{D}_f(s) : s \in N\}.$$

Example 2.8. Consider the single-valued neutrosophic graph $G = (X, Y)$ as shown in Fig. 7. By direct calculations, we have $\Delta(G) = (0.5, 0.2, 1.3)$ and $\delta(G) = (0.2, 0.1, 0.6)$.

Definition 2.10. Let N be a nonempty set. A single-valued neutrosophic graph $G = (X, Y)$ on N is said to be an edge regular single-valued neutrosophic graph if every edge in G has the same degree (q_1, q_2, q_3) .

Remark 2.2. Let $G = (X, Y)$ be an edge regular single-valued neutrosophic graph. Then G is said to be an equally edge regular single-valued neutrosophic graph if $q_1 = q_2 = q_3$.

Example 2.9. Consider a graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3\}$ and $L = \{s_1s_2, s_1s_3, s_2s_3\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 8.




Figure 8: Edge regular single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.2, 0.4, 1.2)$, $\mathcal{D}_G(s_2) = (0.2, 0.4, 1.2)$ and $\mathcal{D}_G(s_3) = (0.2, 0.4, 1.2)$. The degree of each edge is given below:

$$\begin{aligned} \mathcal{D}_G(s_1s_2) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_2) - 2(t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (0.2, 0.4, 1.2) + (0.2, 0.4, 1.2) - 2(0.1, 0.2, 0.6), \\ &= (0.2, 0.4, 1.2). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_1s_3) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_3) - 2(t_Y(s_1s_3), i_Y(s_1s_3), f_Y(s_1s_3)), \\ &= (0.2, 0.4, 1.2) + (0.2, 0.4, 1.2) - 2(0.1, 0.2, 0.6), \\ &= (0.2, 0.4, 1.2). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_2s_3) &= \mathcal{D}_G(s_2) + \mathcal{D}_G(s_3) - 2(t_Y(s_2s_3), i_Y(s_2s_3), f_Y(s_2s_3)), \\ &= (0.2, 0.4, 1.2) + (0.2, 0.4, 1.2) - 2(0.1, 0.2, 0.6), \\ &= (0.2, 0.4, 1.2). \end{aligned}$$

It is easy to see that each edge of single-valued neutrosophic graph G has the same degree. Hence G is an edge regular single-valued neutrosophic graph.

Definition 2.11. Let N be a nonempty set. A single-valued neutrosophic graph $G = (X, Y)$ on N is said to be a totally edge regular single-valued neutrosophic graph if every edge in G has the same total degree (p_1, p_2, p_3) .

Example 2.10. Consider a graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3\}$ and $L = \{s_1s_2, s_1s_3, s_2s_3\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 9.




Figure 9: Totally edge regular single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.8, 0.8, 0.7)$, $\mathcal{D}_G(s_2) = (0.9, 0.8, 0.7)$ and $\mathcal{D}_G(s_3) = (0.9, 0.8, 0.4)$.

- The degree of each edge is given below:

$$\begin{aligned} \mathcal{D}_G(s_1s_2) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_2) - 2(t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (0.8, 0.8, 0.7) + (0.9, 0.8, 0.7) - 2(0.4, 0.4, 0.5), \\ &= (0.9, 0.8, 0.4). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_1s_3) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_3) - 2(t_Y(s_1s_3), i_Y(s_1s_3), f_Y(s_1s_3)), \\ &= (0.8, 0.8, 0.7) + (0.9, 0.8, 0.4) - 2(0.4, 0.4, 0.2), \\ &= (0.9, 0.8, 0.7). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_2s_3) &= \mathcal{D}_G(s_2) + \mathcal{D}_G(s_3) - 2(t_Y(s_2s_3), i_Y(s_2s_3), f_Y(s_2s_3)), \\ &= (0.9, 0.8, 0.7) + (0.9, 0.8, 0.4) - 2(0.5, 0.4, 0.2), \\ &= (0.8, 0.8, 0.7). \end{aligned}$$

It is easy to see that $\mathcal{D}_G(s_1s_2) \neq \mathcal{D}_G(s_1s_3) \neq \mathcal{D}_G(s_2s_3)$. So G is not an edge regular single-valued neutrosophic graph.

- The total degree of each edge is calculated as:

$$\begin{aligned} \mathcal{TD}_G(s_1s_2) &= \mathcal{D}_G(s_1s_2) + (t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (1.3, 1.2, 0.9). \end{aligned}$$

$$\begin{aligned} \mathcal{TD}_G(s_1s_3) &= \mathcal{D}_G(s_1s_3) + (t_Y(s_1s_3), i_Y(s_1s_3), f_Y(s_1s_3)), \\ &= (1.3, 1.2, 0.9). \end{aligned}$$

$$\begin{aligned} \mathcal{TD}_G(s_2s_3) &= \mathcal{D}_G(s_2s_3) + (t_Y(s_2s_3), i_Y(s_2s_3), f_Y(s_2s_3)), \\ &= (1.3, 1.2, 0.9). \end{aligned}$$

It is easy to see that each edge of single-valued neutrosophic graph G has the same total degree. So G is a totally edge regular single-valued neutrosophic graph.

Remark 2.3. A single-valued neutrosophic graph G is an edge regular single-valued neutrosophic graph if and only if $\Delta_{\mathcal{D}}(G) = \delta_{\mathcal{D}}(G) = (q_1, q_2, q_3)$.

Example 2.11. Consider a graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4\}$ and $L = \{s_1s_2, s_2s_3, s_3s_4\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 10.




Figure 10: Single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.1, 0.4, 0.2)$, $\mathcal{D}_G(s_2) = (0.3, 0.9, 0.3)$, $\mathcal{D}_G(s_3) = (0.3, 0.9, 0.3)$ and $\mathcal{D}_G(s_4) = (0.1, 0.4, 0.2)$.

- The degree of each edge is given below:

$$\begin{aligned} \mathcal{D}_G(s_1s_2) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_2) - 2(t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (0.1, 0.4, 0.2) + (0.3, 0.9, 0.3) - 2(0.1, 0.4, 0.2), \\ &= (0.2, 0.5, 0.1). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_2s_3) &= \mathcal{D}_G(s_2) + \mathcal{D}_G(s_3) - 2(t_Y(s_2s_3), i_Y(s_2s_3), f_Y(s_2s_3)), \\ &= (0.3, 0.9, 0.3) + (0.3, 0.9, 0.3) - 2(0.2, 0.5, 0.1), \\ &= (0.2, 0.8, 0.4). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_3s_4) &= \mathcal{D}_G(s_3) + \mathcal{D}_G(s_4) - 2(t_Y(s_3s_4), i_Y(s_3s_4), f_Y(s_3s_4)), \\ &= (0.3, 0.9, 0.3) + (0.1, 0.4, 0.2) - 2(0.1, 0.4, 0.2), \\ &= (0.2, 0.5, 0.1). \end{aligned}$$

It is easy to see that $\mathcal{D}_G(s_1s_2) \neq \mathcal{D}_G(s_2s_3)$. So G is not an edge regular single-valued neutrosophic graph.

- The total degree of each edge is calculated as:

$$\begin{aligned} \mathcal{TD}_G(s_1s_2) &= \mathcal{D}_G(s_1s_2) + (t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (0.3, 0.9, 0.3). \end{aligned}$$

$$\begin{aligned} \mathcal{TD}_G(s_2s_3) &= \mathcal{D}_G(s_2s_3) + (t_Y(s_2s_3), i_Y(s_2s_3), f_Y(s_2s_3)), \\ &= (0.4, 1.3, 0.5). \end{aligned}$$

$$\begin{aligned} \mathcal{TD}_G(s_3s_4) &= \mathcal{D}_G(s_3s_4) + (t_Y(s_3s_4), i_Y(s_3s_4), f_Y(s_3s_4)), \\ &= (0.3, 0.9, 0.3). \end{aligned}$$

It is easy to see that $\mathcal{TD}_G(s_1s_2) \neq \mathcal{TD}_G(s_2s_3)$. So G is not a totally edge regular single-valued neutrosophic graph.

Remark 2.4. A complete single-valued neutrosophic graph G may not be an edge regular single-valued neutrosophic graph.

Example 2.12. Consider a graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4\}$ and $L = \{s_1s_2, s_1s_3, s_1s_4, s_2s_3, s_2s_4, s_3s_4\}$. The corresponding complete single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 11.




Figure 11: Complete single-valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (1.1, 1.2, 1.9)$, $\mathcal{D}_G(s_2) = (0.9, 1.2, 1.9)$, $\mathcal{D}_G(s_3) = (1.4, 1.2, 2.4)$ and $\mathcal{D}_G(s_4) = (1.4, 1.4, 2.0)$. The degree of each edge is given below:

$$\begin{aligned} \mathcal{D}_G(s_1s_2) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_2) - 2(t_Y(s_1s_2), i_Y(s_1s_2), f_Y(s_1s_2)), \\ &= (1.1, 1.2, 1.9) + (0.9, 1.2, 1.9) - 2(0.3, 0.3, 0.5), \\ &= (1.4, 2.0, 2.8). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_1s_3) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_3) - 2(t_Y(s_1s_3), i_Y(s_1s_3), f_Y(s_1s_3)), \\ &= (1.1, 1.2, 1.9) + (1.4, 1.2, 2.4) - 2(0.4, 0.4, 0.8), \\ &= (1.7, 1.6, 2.7). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_1s_4) &= \mathcal{D}_G(s_1) + \mathcal{D}_G(s_4) - 2(t_Y(s_1s_4), i_Y(s_1s_4), f_Y(s_1s_4)), \\ &= (1.1, 1.2, 1.9) + (1.4, 1.4, 2.0) - 2(0.4, 0.5, 0.6), \\ &= (1.7, 1.6, 2.7). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_2s_3) &= \mathcal{D}_G(s_2) + \mathcal{D}_G(s_3) - 2(t_Y(s_2s_3), i_Y(s_2s_3), f_Y(s_2s_3)), \\ &= (0.9, 1.2, 1.9) + (1.4, 1.2, 2.4) - 2(0.3, 0.4, 0.8), \\ &= (1.7, 1.6, 2.7). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_2s_4) &= \mathcal{D}_G(s_2) + \mathcal{D}_G(s_4) - 2(t_Y(s_2s_4), i_Y(s_2s_4), f_Y(s_2s_4)), \\ &= (0.9, 1.2, 1.9) + (1.4, 1.4, 2.0) - 2(0.3, 0.5, 0.6), \\ &= (1.7, 1.6, 2.7). \end{aligned}$$

$$\begin{aligned} \mathcal{D}_G(s_3s_4) &= \mathcal{D}_G(s_3) + \mathcal{D}_G(s_4) - 2(t_Y(s_3s_4), i_Y(s_3s_4), f_Y(s_3s_4)), \\ &= (1.4, 1.2, 2.4) + (1.4, 1.4, 2.0) - 2(0.7, 0.4, 0.8), \\ &= (1.4, 1.8, 2.8). \end{aligned}$$

It is easy to see that each edge of single-valued neutrosophic graph G has not the same degree. Therefore, G is a complete single-valued neutrosophic graph but not an edge regular single-valued neutrosophic graph.

Theorem 2.1. Let $G = (X, Y)$ be a single-valued neutrosophic graph. Then

$$\sum_{st \in L} \mathcal{D}_G(st) = \sum_{st \in L} \mathcal{D}_{G^*}(st)(t_Y(st), i_Y(st), f_Y(st)),$$

where $\mathcal{D}_{G^*}(st) = \mathcal{D}_{G^*}(s) + \mathcal{D}_{G^*}(t) - 2$ for all $s, t \in N$.

Theorem 2.2. Let $G = (X, Y)$ be a single-valued neutrosophic graph. Then

$$\sum_{st \in L} \mathcal{T}\mathcal{D}_G(st) = \sum_{st \in L} \mathcal{D}_{G^*}(st)(t_Y(st), i_Y(st), f_Y(st)) + \mathcal{S}(G),$$

where $\mathcal{D}_{G^*}(st) = \mathcal{D}_{G^*}(s) + \mathcal{D}_{G^*}(t) - 2$ for all $s, t \in N$.

Proof. Since the total degree of each edge in a single-valued neutrosophic graph G is $\mathcal{T}\mathcal{D}_G(st) = \mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st))$. Therefore,

$$\begin{aligned} \sum_{st \in L} \mathcal{T}\mathcal{D}_G(st) &= \sum_{st \in L} (\mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st))), \\ \sum_{st \in L} \mathcal{T}\mathcal{D}_G(st) &= \sum_{st \in L} \mathcal{D}_G(st) + \sum_{st \in L} (t_Y(st), i_Y(st), f_Y(st)), \\ \sum_{st \in L} \mathcal{T}\mathcal{D}_G(st) &= \sum_{st \in L} \mathcal{D}_{G^*}(st)(t_Y(st), i_Y(st), f_Y(st)) + \mathcal{S}(G). \end{aligned}$$

This completes the proof. □

Theorem 2.3. Let $G^* = (N, L)$ be an edge regular crisp graph of degree q and $G = (X, Y)$ be an edge regular single-valued neutrosophic graph of degree (q_1, q_2, q_3) of G^* . Then the size of G is $(\frac{mq_1}{q}, \frac{mq_2}{q}, \frac{mq_3}{q})$, where $|L| = m$.

Proof. Let $G = (X, Y)$ be an edge regular single-valued neutrosophic graph of an edge regular crisp graph $G^* = (N, L)$. Therefore, $\mathcal{D}_G(st) = (q_1, q_2, q_3)$ and $\mathcal{D}_{G^*}(st) = q$ for each edge $st \in L$. Since,

$$\begin{aligned} \sum_{st \in L} \mathcal{D}_G(st) &= \sum_{st \in L} \mathcal{D}_{G^*}(st)(t_Y(st), i_Y(st), f_Y(st)), \\ \sum_{st \in L} (q_1, q_2, q_3) &= q \sum_{st \in L} (t_Y(st), i_Y(st), f_Y(st)), \\ m(q_1, q_2, q_3) &= q\mathcal{S}(G), \\ (mq_1, mq_2, mq_3) &= q\mathcal{S}(G), \\ \mathcal{S}(G) &= (\frac{mq_1}{q}, \frac{mq_2}{q}, \frac{mq_3}{q}). \end{aligned}$$

This completes the proof. □

Theorem 2.4. Let $G^* = (N, L)$ be an edge regular crisp graph of degree q and $G = (X, Y)$ be a totally edge regular single-valued neutrosophic graph of degree (p_1, p_2, p_3) of G^* . Then the size of G is $(\frac{mp_1}{q+1}, \frac{mp_2}{q+1}, \frac{mp_3}{q+1})$, where $|L| = m$.

Proof. Let $G = (X, Y)$ be a totally edge regular single-valued neutrosophic graph of an edge regular crisp graph $G^* = (N, L)$. Therefore, $\mathcal{D}_G(st) = (p_1, p_2, p_3)$ and $\mathcal{D}_{G^*}(st) = q$ for each edge $st \in L$. Since,

$$\begin{aligned} \sum_{st \in L} \mathcal{T}\mathcal{D}_G(st) &= \sum_{st \in L} \mathcal{D}_{G^*}(st)(t_Y(st), i_Y(st), f_Y(st)) + \mathcal{S}(G), \\ \sum_{st \in L} (p_1, p_2, p_3) &= q \sum_{st \in L} (t_Y(st), i_Y(st), f_Y(st)) + \mathcal{S}(G), \\ m(p_1, p_2, p_3) &= q\mathcal{S}(G) + \mathcal{S}(G), \\ (mp_1, mp_2, mp_3) &= (q+1)\mathcal{S}(G), \\ \mathcal{S}(G) &= (\frac{mp_1}{q+1}, \frac{mp_2}{q+1}, \frac{mp_3}{q+1}). \end{aligned}$$

This completes the proof. □

Theorem 2.5. Let $G^* = (N, L)$ be a crisp graph. Suppose that $G = (X, Y)$ be an edge regular single-valued neutrosophic graph of degree (q_1, q_2, q_3) and a totally edge regular single-valued neutrosophic graph of degree (p_1, p_2, p_3) of G^* . Then the size of G is $m(p_1 - q_1, p_2 - q_2, p_3 - q_3)$, where $|L| = m$.

Proof. Let $G = (X, Y)$ be an edge regular single-valued neutrosophic graph and a totally edge regular single-valued neutrosophic graph of a crisp graph $G^* = (N, L)$. Therefore, $\mathcal{D}_G(st) = (q_1, q_2, q_3)$ and $\mathcal{TD}_G(st) = (p_1, p_2, p_3)$ for each edge $st \in L$. Since,

$$\begin{aligned}\mathcal{TD}_G(st) &= \mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st)), \\ \sum_{st \in L} \mathcal{TD}_G(st) &= \sum_{st \in L} \mathcal{D}_G(st) + \sum_{st \in L} (t_Y(st), i_Y(st), f_Y(st)), \\ m(p_1, p_2, p_3) &= m(q_1, q_2, q_3) + \mathcal{S}(G), \\ \mathcal{S}(G) &= m(p_1 - q_1, p_2 - q_2, p_3 - q_3).\end{aligned}$$

This completes the proof. \square

Theorem 2.6. Let $G^* = (N, L)$ be a crisp graph, which is a cycle on m vertices. Suppose that $G = (X, Y)$ be a single-valued neutrosophic graph of G^* . Then $\sum_{s_k \in N} \mathcal{D}_G(s_k) = \sum_{s_k s_l \in L} \mathcal{D}_G(s_k s_l)$.

Proof. Let $G = (X, Y)$ be a single-valued neutrosophic graph of G^* . Suppose that G^* be a cycle $s_1, s_2, s_3, \dots, s_m, s_1$ on m vertices. Then

$$\begin{aligned}\sum_{s_k s_l \in L} \mathcal{D}_G(s_k s_l) &= \mathcal{D}_G(s_1 s_2) + \mathcal{D}_G(s_2 s_3) + \dots + \mathcal{D}_G(s_m s_1), \\ &= [\mathcal{D}_G(s_1) + \mathcal{D}_G(s_2) - 2(t_Y(s_1 s_2), i_Y(s_1 s_2), f_Y(s_1 s_2))] [\mathcal{D}_G(s_2) \\ &+ \mathcal{D}_G(s_3) - 2(t_Y(s_2 s_3), i_Y(s_2 s_3), f_Y(s_2 s_3))] + \dots + [\mathcal{D}_G(s_m) \\ &+ \mathcal{D}_G(s_1) - 2(t_Y(s_m s_1), i_Y(s_m s_1), f_Y(s_m s_1))], \\ &= 2\mathcal{D}_G(s_1) + 2\mathcal{D}_G(s_2) + \dots + 2\mathcal{D}_G(s_m) - 2(t_Y(s_1 s_2), i_Y(s_1 s_2), f_Y(s_1 s_2)), \\ &- 2(t_Y(s_2 s_3), i_Y(s_2 s_3), f_Y(s_2 s_3)) - \dots - 2(t_Y(s_m s_1), i_Y(s_m s_1), f_Y(s_m s_1)), \\ &= 2 \sum_{s_k \in N} \mathcal{D}_G(s_k) - 2 \sum_{s_k s_l \in L} (t_Y(s_k s_l), i_Y(s_k s_l), f_Y(s_k s_l)), \\ &= \sum_{s_k \in N} \mathcal{D}_G(s_k) + \sum_{s_k \in N} \mathcal{D}_G(s_k) - 2 \sum_{s_k s_l \in L} (t_Y(s_k s_l), i_Y(s_k s_l), f_Y(s_k s_l)), \\ &= \sum_{s_k \in N} \mathcal{D}_G(s_k) + 2 \sum_{s_k s_l \in L} (t_Y(s_k s_l), i_Y(s_k s_l), f_Y(s_k s_l)) \\ &- 2 \sum_{s_k s_l \in L} (t_Y(s_k s_l), i_Y(s_k s_l), f_Y(s_k s_l)), \\ &= \sum_{s_k \in N} \mathcal{D}_G(s_k).\end{aligned}$$

This completes the proof. \square

Theorem 2.7. Let $G = (X, Y)$ be a single-valued neutrosophic graph. Then Y is a constant function if and only if the following statements are equivalent:

- (a) G is an edge regular single-valued neutrosophic graph,
- (b) G is a totally edge regular single-valued neutrosophic graph.

Proof. Let $G = (X, Y)$ be a single-valued neutrosophic graph. Suppose that Y is a constant function, then $t_Y(st) = l_1, i_Y(st) = l_2, f_Y(st) = l_3$ for all $st \in L$.

(a) \Rightarrow (b): Assume that G is an edge regular single-valued neutrosophic graph, i.e., $\mathcal{D}_G(st) = (q_1, q_2, q_3)$, for each edge $st \in L$. This implies that $\mathcal{TD}_G(st) = (l_1 + q_1, l_2 + q_2, l_3 + q_3)$ for each edge $st \in L$. This shows that G is an edge regular single-valued neutrosophic graph of degree $(l_1 + q_1, l_2 + q_2, l_3 + q_3)$.

(b) \Rightarrow (a): Suppose that G is a totally edge regular single-valued neutrosophic graph, i.e., $\mathcal{TD}_G(st) = (p_1, p_2, p_3)$ for all $st \in L$. This implies that $\mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st)) = (p_1, p_2, p_3)$. This implies that $\mathcal{D}_G(st) = (p_1, p_2, p_3) - (t_Y(st), i_Y(st), f_Y(st))$. This implies that $\mathcal{D}_G(st) = (p_1 - l_1, p_2 - l_2, p_3 - l_3)$ for each edge $st \in L$. Thus G is an edge regular single-valued neutrosophic graph of degree $(p_1 - l_1, p_2 - l_2, p_3 - l_3)$. Hence the statements (a) and (b) are equivalent.

Conversely, suppose that (a) and (b) are equivalent. Assume that Y is not a constant function. This implies that $(t_Y(st), i_Y(st), f_Y(st)) \neq (t_Y(uv), i_Y(uv), f_Y(uv))$ for at least one pair of edges $st, uv \in L$. Assume that G is an edge regular single-valued neutrosophic graph. This implies that $\mathcal{D}_G(st) = \mathcal{D}_G(uv) = (q_1, q_2, q_3)$. This implies that $\mathcal{TD}_G(st) = \mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st)) = (q_1, q_2, q_3) + (t_Y(st), i_Y(st), f_Y(st))$ and $\mathcal{TD}_G(uv) = \mathcal{D}_G(uv) + (t_Y(uv), i_Y(uv), f_Y(uv)) = (q_1, q_2, q_3) + (t_Y(uv), i_Y(uv), f_Y(uv))$. Since $(t_Y(st), i_Y(st), f_Y(st)) \neq (t_Y(uv), i_Y(uv), f_Y(uv))$. This implies that $\mathcal{TD}_G(st) \neq \mathcal{TD}_G(uv)$. This shows that G is not a totally edge regular single-valued neutrosophic graph, which contradicts our supposition. Now, suppose that G is a totally edge regular single-valued neutrosophic graph, i.e., $\mathcal{TD}_G(st) = \mathcal{TD}_G(uv) = (p_1, p_2, p_3)$. This implies that $\mathcal{TD}_G(st) = \mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st)) = \mathcal{D}_G(uv) + (t_Y(uv), i_Y(uv), f_Y(uv))$. This implies that $\mathcal{D}_G(st) - \mathcal{D}_G(uv) = (t_Y(st), i_Y(st), f_Y(st)) - (t_Y(uv), i_Y(uv), f_Y(uv))$. Since $(t_Y(st), i_Y(st), f_Y(st)) \neq (t_Y(uv), i_Y(uv), f_Y(uv))$. This implies that $\mathcal{D}_G(st) - \mathcal{D}_G(uv) \neq 0$. This implies that $\mathcal{D}_G(st) \neq \mathcal{D}_G(uv)$. This shows that G is not an edge regular single-valued neutrosophic graph, which contradicts our supposition. Hence Y is a constant function. \square

Theorem 2.8. *Let $G = (X, Y)$ be a single-valued neutrosophic graph. Assume that G is both edge regular single-valued neutrosophic of degree (q_1, q_2, q_3) and totally edge regular single-valued neutrosophic graph of degree (p_1, p_2, p_3) . Then Y is a constant function.*

Proof. The proof is obvious. \square

Remark 2.5. The converse of theorem 2.8 may not be true in general, that is, a single-valued neutrosophic graph $G = (X, Y)$, where Y is a constant function, may or may not be edge regular and totally edge regular single-valued neutrosophic graph.

Example 2.13. Consider a graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4\}$ and $L = \{s_1s_2, s_2s_3, s_3s_4\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 12.




Figure 12: Single valued neutrosophic graph G

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.1, 0.2, 0.4)$, $\mathcal{D}_G(s_2) = (0.2, 0.4, 0.8)$, $\mathcal{D}_G(s_3) = (0.2, 0.4, 0.8)$ and $\mathcal{D}_G(s_4) = (0.1, 0.2, 0.4)$. The degree of each edge is $\mathcal{D}_G(s_1s_2) = (0.1, 0.2, 0.4)$, $\mathcal{D}_G(s_2s_3) = (0.2, 0.4, 0.8)$ and $\mathcal{D}_G(s_3s_4) = (0.1, 0.2, 0.4)$. The total degree of each edge is $\mathcal{TD}_G(s_1s_2) = (0.2, 0.4, 0.8)$, $\mathcal{TD}_G(s_2s_3) = (0.3, 0.6, 1.2)$. It is clear from above calculations that G is neither an edge regular nor a totally edge regular single-valued neutrosophic graph.

Theorem 2.9. *Let $G = (X, Y)$ be a single-valued neutrosophic graph of $G^* = (N, L)$, where Y is a constant function. If G is a regular single-valued neutrosophic graph. Then G is an edge regular single-valued neutrosophic graph.*

Proof. Assume that Y is a constant function, that is, $t_Y(st) = l_1$, $i_Y(st) = l_2$ and $f_Y(st) = l_3$ for all $st \in L$. Suppose that G is a regular single-valued neutrosophic graph, that is, $\mathcal{D}_G(s) = (m_1, m_2, m_3)$ for all $s \in N$. Now

$$\begin{aligned}\mathcal{D}_G(st) &= \mathcal{D}_G(s) + \mathcal{D}_G(t) - 2(t_Y(st), i_Y(st), f_Y(st)), \\ &= (m_1, m_2, m_3) + (m_1, m_2, m_3) - 2(l_1, l_2, l_3), \\ &= 2(m_1 - l_1, m_2 - l_2, m_3 - l_3),\end{aligned}$$

for all $st \in L$. Hence G is an edge regular single-valued neutrosophic graph. \square

Theorem 2.10. *Let $G = (X, Y)$ be a single-valued neutrosophic graph of $G^* = (N, L)$, where Y is a constant function. If G is a regular single-valued neutrosophic graph. Then G is a totally edge regular single-valued neutrosophic graph.*

Proof. Let Y be a constant function, that is, $t_Y(st) = l_1$, $i_Y(st) = l_2$ and $f_Y(st) = l_3$ for all $st \in L$. Assume that G is a regular single-valued neutrosophic graph, that is, $\mathcal{D}_G(s) = (m_1, m_2, m_3)$ for all $s \in N$. Then G is an edge regular single-valued neutrosophic graph, that is, $\mathcal{D}_G(st) = (q_1, q_2, q_3)$. Now

$$\begin{aligned}\mathcal{T}\mathcal{D}_G(st) &= \mathcal{D}_G(st) + (t_Y(st), i_Y(st), f_Y(st)), \\ &= (q_1, q_2, q_3) + (l_1, l_2, l_3), \\ &= 2(q_1 + l_1, q_2 + l_2, q_3 + l_3),\end{aligned}$$

for all $st \in L$. Hence G is a totally edge regular single-valued neutrosophic graph. \square

Theorem 2.11. *Let $G^* = (N, L)$ be a regular crisp graph. Suppose that $G = (X, Y)$ is a single-valued neutrosophic graph of G^* . Then G is both regular and totally edge regular single-valued neutrosophic graph if and only if Y is a constant function.*

Proof. Let $G^* = (N, L)$ be a regular crisp graph. Suppose that $G = (X, Y)$ is a single-valued neutrosophic graph of G^* . Suppose that G is both regular and totally edge regular single-valued neutrosophic graph, that is, $\mathcal{D}_G(s) = (m_1, m_2, m_3)$ for all $s \in N$ and $\mathcal{T}\mathcal{D}_G(st) = (p_1, p_2, p_3)$ for all $st \in L$. Now

$$\begin{aligned}\mathcal{T}\mathcal{D}_G(st) &= \mathcal{D}_G(s) + \mathcal{D}_G(t) - (t_Y(st), i_Y(st), f_Y(st)), \quad \forall st \in L, \\ (p_1, p_2, p_3) &= (m_1, m_2, m_3) + (m_1, m_2, m_3) - (t_Y(st), i_Y(st), f_Y(st)), \\ (t_Y(st), i_Y(st), f_Y(st)) &= (2m_1 - p_1, 2m_2 - p_2, 2m_3 - p_3),\end{aligned}$$

for all $st \in L$. Hence Y is a constant function. Conversely, let Y be a constant function, that is, $t_Y(st) = l_1$, $i_Y(st) = l_2$ and $f_Y(st) = l_3$ for all $st \in L$. So

$$\begin{aligned}\mathcal{D}_G(s) &= \sum_{st \in L} (t_Y(st), i_Y(st), f_Y(st)), \quad \forall s \in N, \\ &= \sum_{st \in L} (m_1, m_2, m_3), \\ &= (m_1, m_2, m_3)\mathcal{D}_{G^*}(s), \\ &= (m_1, m_2, m_3)m.\end{aligned}$$

This implies that $\mathcal{D}_G(s) = (mm_1, mm_2, mm_3)$ for all $s \in L$. Thus G is a regular single-valued neutrosophic graph. Now

$$\begin{aligned}\mathcal{T}\mathcal{D}_G(st) &= \sum_{sa \in L, s \neq a} (t_Y(sa), i_Y(sa), f_Y(sa)) + \sum_{at \in L, a \neq t} (t_Y(at), i_Y(at), f_Y(at)), \\ &+ (t_Y(st), i_Y(st), f_Y(st)) \quad \forall st \in L, \\ &= \sum_{sa \in L, s \neq a} (l_1, l_2, l_3) + \sum_{at \in L, a \neq t} (l_1, l_2, l_3) + (l_1, l_2, l_3), \\ &= (l_1, l_2, l_3)(\mathcal{D}_{G^*}(s) - 1) + (l_1, l_2, l_3)(\mathcal{D}_{G^*}(t) - 1) + (l_1, l_2, l_3), \\ &= (l_1, l_2, l_3)(s - 1) + (l_1, l_2, l_3)(t - 1) + (l_1, l_2, l_3), \\ &= (2l_1, 2l_2, 2l_3)(s - 1) + (l_1, l_2, l_3),\end{aligned}$$

for all $st \in L$. Hence G is a totally edge regular single-valued neutrosophic graph. \square

Theorem 2.12. Let $G^* = (N, L)$ be a crisp graph. Suppose that $G = (X, Y)$ is a single-valued neutrosophic graph of G^* . Then Y is a constant function if and only if G is an edge regular single-valued neutrosophic graph.

Proof. Let G be a regular single-valued neutrosophic graph, that is, $\mathcal{D}_G(s) = (m_1, m_2, m_3)$, for all $s \in N$. Suppose that Y is a constant function, that is, $t_Y(st) = l_1$, $i_Y(st) = l_2$ and $f_Y(st) = l_3$, for all $st \in L$. Now

$$\begin{aligned} \mathcal{D}_G(st) &= \mathcal{D}_G(s) + \mathcal{D}_G(t) - 2(t_Y(st), i_Y(st), f_Y(st)), \quad \forall st \in L. \\ &= (m_1, m_2, m_3) + (m_1, m_2, m_3) - 2(l_1, l_2, l_3), \end{aligned}$$

this implies that $\mathcal{D}_G(st) = 2(m_1, m_2, m_3) - 2(l_1, l_2, l_3)$, for all $st \in L$. Hence G is an edge regular single-valued neutrosophic graph.

Conversely, assume that G is an edge regular single-valued neutrosophic graph, that is, $\mathcal{D}_G(st) = (q_1, q_2, q_3)$ for each edge $st \in L$. Now

$$\begin{aligned} \mathcal{D}_G(st) &= \mathcal{D}_G(s) + \mathcal{D}_G(t) - 2(t_Y(st), i_Y(st), f_Y(st)), \quad \forall st \in L, \\ (q_1, q_2, q_3) &= (m_1, m_2, m_3) + (m_1, m_2, m_3) - 2(t_Y(st), i_Y(st), f_Y(st)), \end{aligned}$$

this implies that $(t_Y(st), i_Y(st), f_Y(st)) = \frac{(q_1, q_2, q_3) - (2m_1, 2m_2, 2m_3)}{2}$, for all $st \in L$. Thus Y is a constant function. \square

Definition 2.12. Let G^* be an edge regular crisp graph. Then a single-valued neutrosophic graph G of G^* is said to be a partially edge regular single-valued neutrosophic graph.

Example 2.14. It can be seen in example 2.12 that G^* is an edge regular crisp graph. Therefore, G is a partially edge regular single-valued neutrosophic graph.

Definition 2.13. Let G^* be an edge regular crisp graph. A single-valued neutrosophic graph G of G^* is said to be a full edge regular single-valued neutrosophic graph if it is both edge regular and partially edge regular.

Example 2.15. Consider a graph $G^* = (N, L)$ such that $N = \{s_1, s_2, s_3, s_4\}$ and $L = \{s_1s_2, s_2s_3, s_3s_4, s_1s_4\}$. The corresponding single-valued neutrosophic graph $G = (X, Y)$ is shown in Fig. 13.

By direct calculations, we have $\mathcal{D}_G(s_1) = (0.4, 0.8, 0.8)$, $\mathcal{D}_G(s_2) = (0.4, 0.8, 0.8)$, $\mathcal{D}_G(s_3) = (0.4, 0.8, 0.8)$ and $\mathcal{D}_G(s_4) = (0.4, 0.8, 0.8)$. The degree of each edge is $\mathcal{D}_G(s_1s_2) = (0.4, 0.8, 0.8)$, $\mathcal{D}_G(s_2s_3) = (0.4, 0.8, 0.8)$, $\mathcal{D}_G(s_3s_4) = (0.4, 0.8, 0.8)$ and $\mathcal{D}_G(s_1s_4) = (0.4, 0.8, 0.8)$. It is clear from calculations that G is full edge regular single-valued neutrosophic graph.




Figure 13: Full edge regular single-valued neutrosophic graph

Theorem 2.13. Let $G = (X, Y)$ be a single-valued neutrosophic graph of a crisp graph $G^* = (N, L)$, where Y is a constant function. Then G is full edge regular single-valued neutrosophic graph if it is full regular single-valued neutrosophic graph.

Proof. Let $G = (X, Y)$ be a single-valued neutrosophic graph of a crisp graph $G^* = (N, L)$. Suppose that Y is a constant function, that is, $(t_Y(st), i_Y(st), f_Y(st)) = (l_1, l_2, l_3)$ for each edge $st \in L$. Assume that G is full regular single-valued neutrosophic graph. Then G is both regular and partially regular. Therefore, $\mathcal{D}_G(s) = (m_1, m_2, m_3)$ and

$\mathcal{D}_{G^*}(s) = m$ for all $s \in N$. Since $\mathcal{D}_{G^*}(st) = \mathcal{D}_{G^*}(s) + \mathcal{D}_{G^*}(t) - 2$ for all $st \in L$. This shows that $\mathcal{D}_{G^*}(st) = 2m - 2$. Therefore, G^* is an edge regular single-valued neutrosophic graph. Now

$$\begin{aligned} \mathcal{D}_G(st) &= \mathcal{D}_G(s) + \mathcal{D}_G(t) - 2(t_Y(st), i_Y(st), f_Y(st)), \quad \forall st \in L. \\ &= (m_1, m_2, m_3) + (m_1, m_2, m_3) - 2(l_1, l_2, l_3), \end{aligned}$$

this implies that $\mathcal{D}_G(st) = 2(m_1 - l_1, m_2 - l_2, m_3 - l_3)$. This shows that G is an edge regular single-valued neutrosophic graph. Hence G is a full edge regular single-valued neutrosophic graph. \square

3 Conclusion

Neutrosophic sets are the generalization of the concept of fuzzy sets and intuitionistic fuzzy sets. Neutrosophic models give more flexibility, precisions and compatibility to the system as compared to the classical, fuzzy and intuitionistic fuzzy models. In this research paper, we have discussed certain types of edge-regular single-valued neutrosophic graphs. Here, we have established some theorems on single valued neutrosophic graphs. It is known that the single-valued neutrosophic graphs are successfully used to analysis the uncertain situation in the real World. Thus we aim to widen our research of fuzzification to (1) single-valued neutrosophic soft graphs, (2) single-valued neutrosophic rough fuzzy graphs, (3) Roughness in neutrosophic graphs and (4) Application of intuitionistic neutrosophic graphs in decision support systems.

References

- [1] Akram, M., and Shahzadi, S. (2017), *Neutrosophic soft graphs with applicatioon*, Journal of Intelligent & Fuzzy Systems, **32**(1), 841-858.
- [2] Akram, M., Shahzadi, S. and A. Borumand saeid. (2016), *Single-valued neutrosophic hypergraphs*, TWMS Journal of Applied and Engineering Mathematics, (In press).
- [3] Akram, M., (2016), *Single-valued neutrosophic planar graphs*, International Journal of Algebra and Statistics, **5**(2), 157-167.
- [4] Akram, M., Shahzadi, G. (2017), *Operations on single-valued neutrosophic graphs*, Journal of Uncertain System, **11**:1-26.
- [5] Akram, M., Shahzadi, S. (2016), *Representation of graphs using intuitionistic neutrosophic soft sets*, Journal of Mathematical Analysis, **7**(6), 31-53.
- [6] Atanassov K., (1986), *Intuitionistic fuzzy sets*, Fuzzy sets and Systems, **20**, 87-96.
- [7] Bhattacharya P., (1987), *Some remarks on fuzzy graphs*, Pattern Recognition Letter, **6**, 297-302.
- [8] Broumi, S., Talea, M., Bakali A., and Smarandache F., (2016), *Single-valued neutrosophic graphs*, Journal of New Theory, **10**, 86-101.
- [9] Broumi, S., Talea, M., Bakali A., and Smarandache F., (2016), *Single Valued Neutrosophic Graphs: Degree, Order and Size*, IEEE International Conference on Fuzzy Systems, 2444-2451.
- [10] Dhavaseelan, R., Vikramaprasad, R., and Krishnaraj V., (2015), *Certain types of neutrosophic graphs*, International Journal of Mathematical Sciences and Applications, **5**(2), 333-339.
- [11] Majumdar P., and Samanta S. K., (2014), *On similarity and entropy of neutrosophic sets*, Journal of Intelligent & Fuzzy Systems, **26**(3), 1245-1252.
- [12] Mordeson, J. N. and Nair, P. S. (2001), *Fuzzy graphs and fuzzy hypergraphs*, Physica Verlag, Heidelberg.
- [13] Nagoorgani A., and Radha K., (2008), *Regular properties of fuzzy graphs*, Bulletin of Pure and Applied Sciences, **27E**(2), 411-419.

- [14] Radha K., and Kumaravel N., (2014), *Some properties of edge regular fuzzy graphs*, Jamal Academic Research Journal, Special issue, 121-127.
- [15] Rosenfeld A., (1975), *Fuzzy graphs, Fuzzy Sets and Their Applications*, Academic Press, New York, 77-95.
- [16] Shah, N., and Hussain, A. (2016). *Neutrosophic soft graphs*, Neutrosophic Set and Systems, **11**, 31 – 44.
- [17] Smarandache F., (1999), *A unifying field in logics. neutrosophy: neutrosophic probability, set and logic. Rehoboth.*, American Research Press.
- [18] Smarandache F., (2006), *Neutrosophic set - a generalization of the intuitionistic fuzzy set*, Granular Computing. 2006 IEEE International Conference, 38-42,DOI: 10.1109/GRC.2006.1635754.
- [19] Wang H., Smarandache F., Zhang Y. Q., and Sunderraman R., (2010), *Single-valued neutrosophic sets*, Multispace and Multistruct, **4**, 410-413.
- [20] Zadeh L. A., (1965), *Fuzzy sets*, Information and control, **8**, 338-353.