Logical Errors of Special Relativity

by

Florian Michael Schmitt

April 2018, Berlin

Abstract

This paper will reveal a fundamental logical error of Special Relativity. It will be shown, strictly deducting from Albert Einstein's own concept, that lengths being measured with clocks do not differ between two systems in relative motion to each other. Furthermore, it is shown that Special Relativity de facto puts two different events into relation and concludes from this the loss of simultaneity.

1. Introduction

Most laymen, but also more and more experts who are confronted with the ideas of Special Relativity, have a faint feeling of lacking of logical reason about the basic structure of theory. In particular, the claim of invariance of the speed of light and equivalence of the relative uniformity of motion are under suspect. Although mathematics and the limited ability of abstraction make it difficult to fully understand the theory of special relativity, we will point out the fundamental fallacy right at the beginning. We will strictly follow Albert Einstein's original paper and try to visualize the mathematical assumptions for clarity.

2. Clock Synchronisation

We first cite Albert Einstein from "On the Electrodynamics of Moving Bodies", Chapter A,1.:

If there is a clock at point A in space, then an observer located at A can evaluate the time of events in the immediate vicinity of A by finding the positions of the hands of the clock that are simultaneous with these events. If there is another clock at point B that in all respects resembles the one at A, then the time of events in the immediate vicinity of B can be evaluated by an observer at B. But it is not possible to compare the time of an event at A with one at B without a further stipulation. So far we have defined only an "A-time" and a "B-time," but not a common "time" for A and B. The latter can now be determined by establishing bydefinition that the "time" required for light to travel from A to B is equal to the "time" it requires to travel from B to A. For, suppose a ray of light leaves from A for B at "A-time" t_A , is reflected from B toward A at "B-time" t_B , and arrives back at A at "A-time" t'_A . The two clocks are synchronous by definition if

$$t_B - t_A = t'_A - t_B.$$

We assume that it is possible for this definition of synchronism to be free of contradictions, and to be so for arbitrarily

Excerpt 1: Synchronization of two clocks

Now we visualize the principle of synchronization of two clocks in A and B in a stationary reference system, strictly based on the description given by Albert Einstein. To be conform to Albert Einstein's text we chose to denominate the system at rest with S'. Light beam is orange, the rod is black, the involved clocks are highlighted in red:



v=0

Fig. 1: Synchronization of two clocks

Again we cite Albert Einstein regarding synchronization of additional clocks:

ELECTRODYNAMICS OF MOVING BODIES

many points, and therefore that the following relations are generally valid:

- 1. If the clock at B runs synchronously with the clock at A, the clock at A runs synchronously with the clock at B.
- 2. If the clock at A runs synchronously with the clock at B as well as with the clock at C, then the clocks at B and C also run synchronously relative to each other.

Excerpt 2: Synchronisation of two more clocks

We now imagine to have two more clocks at C and D and we again synchronize such clocks with B and therefore according to Albert Einstein also with A. Positions of C and D appear random and we will show its purpose later:



v=0

Fig. 2: Synchronisation of two more clocks

Strictly according to Albert Einstein we now have four synchronized clocks in A, B, C and D.

3. Length Measurement Methods

Albert Einstein is proposing two different ways of length measurement. The first method is to simply have a scale and measure the length l of a rod from within any moving reference system S:

Take a rigid rod at rest; let its length, measured by a measuring rod that is also at rest, be l. Now imagine the axis of the rod placed along the X-axis of the rest coordinate system, and the rod then set into uniform parallel translational motion (with velocity v) along the X-axis in the direction of increasing x. We now inquire about the length of the *moving* rod, which we imagine to be ascertained by the following two operations:

a. The observer moves together with the aforementioned measuring rod and the rigid rod to be measured, and measures the length of the rod by laying out the measuring rod in the same way as if the rod to be measured, the observer, and the measuring rod were all at rest.

Excerpt 3: Measurement using scale(operation a.)

Again the visualization for this, measurement of the rod length l from inside the moving system S (bluish). The rod is blue, the measuring scale green:



Fig. 3: Measurement using scale (operation a.)

The second method is measuring the length of the rod r_{AB} by means of the position of the two ends of the rod at a given time t, to be measured with synchronized clocks from a stationary system S':

b. Using clocks at rest and synchronous in the rest system as outlined in section 1, the observer determines at which points of the rest system the beginning and end of the rod to be measured are located at some given time t. The distance between these two points, measured with the rod used before—but now at rest—is also a length that we can call the "length of the rod."

Excerpt 4: Measurement using clocks (operation b.)

It has to be stressed that for now we still only deal with measuring of a rigid rod that is in relative motion, once being measured from within the moving system S (bluish), once from within the stationary system S' (grey). We do <u>not</u> deal with measurement of the length of any light beam. Also it must be emphasized that measurement of the moving rod from the stationary system S' takes place within one instant, as if the two clocks were stopwatches (measurement of the rod with a scale according to Albert Einstein "at some given time t", i.e. instant). The clocks being involved with the measurement are highlighted in red:



 $v = \frac{1}{4}V$

Fig. 4: Measurement using clocks in stationary system (operation b.)

Now Albert Einstein announces that he will show the two ways of measurement giving different lengths for the rod. Also he is claiming that classic physics assume both measurement operations must be equal. We will prove the opposite, that according to classic physics it is not equal, but according to Albert Einstein.

The length determined using operation (b), which we shall call "the length of the (moving) rod in the rest system," will be determined on the basis of our two principles, and we shall find that it differs from l.

Current kinematics tacitly assumes that the lengths determined by the above two operations are exactly equal to each other, or, in other words, that at the time t a moving rigid body is totally replaceable, in geometric respects, by the same body when it is at rest in a particular position.

Excerpt 5: Differing lengths in moving system

4. Non- Simultaneity within two reference systems moving relative to each other

Now Albert Einstein introduces a light beam that is traveling along the moving rod towards a mirror from which it is being reflected back.

Further, we imagine the two ends (A and B) of the rod equipped with clocks that are synchronous with the clocks of the rest system, i.e., whose readings always correspond to the "time of the system at rest" at the locations the clocks happen to occupy; hence, these clocks are "synchronous in the rest system."

We further imagine that each clock has an observer comoving with it, and that these observers apply to the two clocks the criterion for the synchronous rate of two clocks formulated in section 1. Let a ray of light start out from A at time² t_A ; it is reflected from B at time t_B , and arrives back at A at time t'_A . Taking into account the principle of the constancy of the velocity of light, we find that

$$t_B - t_A = \frac{r_{AB}}{V - v}$$

Excerpt 6: Measurement using clocks in moving system

The difference Δt_1 $(t_B - t_A)$ and Δt_2 $(t_A - t_B)$ is computed strictly according to classic speed addition (rather than on the assumption of "the principle of the constancy of the velocity of light"), as shown by the following visualization.

The light beam (orange) is within the moving system S (bluish), the traveling length of the beam is being measured by classic speed addition, i.e. light travels from A to the right with speed V while the rod (blue) is traveling to the right with speed v inside the moving system. According to classic physics (and to Albert Einstein) movement of the source is irrelevant.

The beam then hits the mirror on the moving rod in B. Albert Einstein has not mentioned at this how the measurement is done at the stationary frame at this same instant, and we show the stationary system with its measuring scale for clarification.



Fig. 5: Measurement using clocks in moving system

Now Albert Einstein claims that Δt_1 $(t_B - t_A)$ in one direction is different from Δt_2 $(t_A - t_B)$ in the opposite direction, hence the principle outlined in excerpt 1 and fig. 1 is violated and thus synchronicity of clocks is broken.

and

$$t_A' - t_B = \frac{r_{AB}}{V + v}.$$

where r_{AB} denotes the length of the moving rod, measured in the rest system. Observers co-moving with the rod would thus find that the two clocks do not run synchronously, while observers in the system at rest would declare them to be running synchronously.

Thus we see that we cannot ascribe absolute meaning to the concept of simultaneity; instead, two events that are simultaneous when observed from some particular coordinate system can no longer be considered simultaneous when observed from a system that is moving relative to that system.

Excerpt 7: Violation of simultaneity

5. Critic

At first we have to have a closer look to the mathematic scheme, in order to see what is being herewith presented to us. As for excerpt 1 we have:

$$t_B - t_A = t_A' - t_B \tag{1}$$

And as per excerpt 6 and 7 we have:

$$t_B - t_A = \frac{r_{AB}}{(V - v)}$$

 $t_A' - t_B = \frac{r_{AB}}{(V + v)}$

Now it is suggesting itself to do a simple operation, we insert the two formulae from (2) into (1):

$$t_B - t_A = t_A' - t_B$$
$$\Rightarrow \frac{r_{AB}}{(V - v)} = \frac{r_{AB}}{(V + v)}$$

$$\Rightarrow (V-v) = (V+v)$$

The above could be true only if light speed were infinite. In any other case we obtain:

(2)

$$v = -v$$

This already gives us a faint glimpse of the problem's root.

We clarify now, how such could happen. At first, we have added the two additional clocks to the stationary system (thus A', B', C', D') that we have already synchronized according to fig. 2, and we have added them equally on both systems.

Secondly we get rid of the basic mistake that is to relate <u>the traveling length of one light beam</u> within the moving system against <u>the measured length of a rigid rod</u> within the stationary system, i.e. we get rid of the fact that Albert Einstein did mix an instantaneous event (measuring a length at one given time t) with a period event (measuring the length of a light beam in one period $t_B - t_A$), i.e. by no means the same events.

Instead, let us examine the question of how one and the same light beam within the moving system S is actually measured in both systems (S and S') that are in relative motion to each other.

The first event is the emergence of the beam from the source in A. In both systems clocks in A and A' give equal and simultaneous readings. The second event is when the light beam hits the mirror on the rod in B while the rod has moved with system S, and again we get equal and simultaneous readings, but this time from clocks B and C' (keep in mind that all clocks are synchronized). At the third event the light beam hits back to the source in A and we obtain equal and simultaneous readings this time from clocks in A and D'. Obviously the distances that have been covered by the light beam are equal in both systems, represented by the orange beam in S and the green scale in S'. We have highlighted in red such clocks (being all synchronized according to Albert Einstein's rule) that are involved with measurement in each reference system at the respective simultaneous events.



Fig. 6: Scheme with correctly measured light beam (with clocks) in two systems in relative motion

5. Conclusion

As we can easily see, we have now the following results:

- 1) All clocks are synchronized.
- 2) Clocks in A and A' are showing the same reading for the same event and instant and are on the very same locality. Likewise for clocks in B and C' and clocks in A and D' for their respective events, instances and localities.
- 3) Clock readings in both reference systems result in the same traveling length of the light beam.
- 4) Equal events and instants take place simultaneously and at the same locality.
- 5) Light speed is equal and constant in both reference systems.
- 6) Both reference systems are equivalent.
- 7) Simultaneous events are observed as simultaneous events in both reference systems.

We have shown that Special Relativity de facto puts two different events into relation, which leads to the claim that simultaneity does not exist within two systems in uniform relative motion. All further conclusions of Special Relativity must be put on the touchstone.

References and Acknowledgments:

[1] Einstein, Albert, Zur Elektrodynamik bewegter Körper, Annalen der Physik und Chemie. 17, 1905,
S. 891–921, 1905.