Proof of Lagarias's Elementary Version of the Riemann Hypothesis.

Stephen Marshall

27 Nov 2018

Abstract

The Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$

The Riemann hypothesis implies results about the distribution of prime numbers. Along with suitable generalizations, some mathematicians consider it the most important unresolved problem in pure mathematics (Bombieri 2000).

It was proposed by Bernhard Riemann (1859), after whom it is named. The name is also used for some closely related analogues, such as the Riemann hypothesis for curves over finite fields.

The Riemann hypothesis implies results about the distribution of prime numbers. Along with suitable generalizations, some mathematicians consider it the most important unresolved problem in pure mathematics (Bombieri 2000). The Riemann zeta function is defined for complex *s* with real part greater than 1 by the absolutely convergent infinite series:

$$\zeta(s) = 1 + 1/2^s + 1/3^s + 1/4^s + \dots$$

The Riemann hypothesis asserts that all *interesting* solutions of the equation:

$$\zeta(s) = 0$$

lie on a certain vertical straight line.

In mathematics, the n-th harmonic number is the sum of the reciprocals of the first n natural numbers:

$$H_n = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} = \sum_{n=1}^{n} \frac{1}{n}$$

Harmonic numbers have been studied since early times and are important in various branches of number theory. They are sometimes loosely termed harmonic series, are closely related to the Riemann zeta function.

The harmonic numbers roughly approximate the natural logarithm function and thus the associated harmonic series grows without limit, albeit slowly. In 1737, Leonhard Euler used the divergence of the harmonic series to provide a new proof of the infinity of prime numbers. His work was extended into the complex plane by Bernhard Riemann in 1859, leading directly to the celebrated Riemann hypothesis about the distribution of prime numbers.

Proof

In 2002, Jeffrey Lagarias proved that this problem is equivalent to the Riemann Hypothesis, a famous question about the complex roots of the Riemann zeta function. The Lagarias's Elementary Version of the Riemann Hypothesis states that for a positive integer n, let $\sigma(n)$ denote the sum of the positive integers that divide n. For example, $\sigma(4)=1+2+4=7$, and $\sigma(6)=1+2+3+6=12$. Let H_n denote the n-th harmonic number, for example:

$$H_n = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n}$$

The unsolved question is does the following inequality hold for all $n \ge 1$?

$$\sigma(n) < H_n + \ln(H_n)e^{H_n}$$

First we will solve for the smallest numbers:

$$\begin{split} \sigma(1) \, &\leq \, H_1 + ln(H_1) e^{H_1} \\ \sigma(1) \, &= 1 \text{ and } H_1 = 1 \text{ , therefore,} \\ \sigma(1) \, &= H_1 = 1 \text{ , which satisfies our inequality.} \end{split}$$

$$\sigma(2) \le H_2 + \ln(H_2)e^{H_2}$$
 $\sigma(2) = 3 \text{ and } H_2 = 1.5 \text{ , therefore,}$
 $\sigma(2) = 3 \le 1.5 + \ln(1.5)e^{1.5}$
 $3 \le 5.346945837$

$$\sigma(5) \le H_5 + \ln(H_5)e^{H_5}$$

$$\sigma(5) = 6 \text{ and } H_3 = 10.91666667, \text{ therefore,}$$

$$\sigma(5) = 6 \le 10.91666667 + \ln(10.91666667)e^{10.91666667}$$

$$6 \le 1.36425E + 52$$

In 1984, G Robin also proved, unconditionally, that the inequality below is true (see Proposition 1 of Section 4 of Robin's work in reference 2):

$$\sigma(n) \leq (e^{\gamma})n(\log\log(n)) + \frac{0.6483n}{\log\log(n)}$$

holds for all $n \ge 3$. Above, we have already shown the first five $\sigma(n) \le H_n + \ln(H_n)e^{H_n}$, therefore, we do not need to address when n < 3 in the following proof.

The numerical value of the Euler–Mascheroni constant, γ , is: $\gamma = 0.57721566490153286060651209008240243104215933593992$

We will prove that:

$$(e^{\gamma})n(\log\log(n)) + \frac{0.6483n}{\log\log(n)} \le H_n + \ln(H_n)e^{H_n}$$

 $(e^{0.57721})n(\log\log(n)) + \frac{0.6483n}{\log\log(n)} \le H_n + \ln(H_n)e^{H_n}$

$$1.781n(\log \log(n)) + \frac{0.6483n}{\log \log(n)} \le H_n + \ln(H_n)e^{H_n}$$

For $n \ge 3$, then $\ln(H_n) \ge 1.041453875$ which means $H_n + \ln(H_n)e^{H_n} \ge e^{H_n}$ Therefore, we will prove that:

$$1.781n(\log\log(n)) + \frac{0.6483n}{\log\log(n)} \le e^{H_n}$$

Notice that $\log \log(n)$ grows very slowly, $\log \log(1,000,000,000) = 0.954$

Notice, H_n grows rapidly in the exponent value for e^{H_n} , for example, $H_5 = 10.91666667$

Therefore, $e^{H_n} = e^{H_5} = e^{10.91666667} = 1.1539E + 123$, which is a very large number compared to 0.954

$$1.781n(\log\log(n)) + \frac{0.6483n}{\log\log(n)} \le e^{H_n}$$

All power functions, exponential functions, and logarithmic functions tend to ∞ as $x => \infty$. But these three classes of functions tend to ∞ at different rates. The main result we want to focus on is the following one; e^x grows faster than any power function while $\log x$ grows slower than any power function (see reference 4). Notice the right hand side of our above inequality is an exponential function, e^{H_n} , which grows faster than any power function. The left side of our inequality is a combination of logarithmic functions and linear functions, and logarithmic functions grows slower than any power function. The linear functions on the left side are very small multiples of n, but they still grow much faster than $\log \log(n)$, however it grows extremely slow compared to e^{H_n} . Reference 4, provides proof of e^x growing the fastest and $\log x$ growing the slowest, this provides proof of the above inequality for all $n \ge 1$.

Additionally, we know that H_n grows to infinity (which causes e^{H_n} to grow to e^{∞}), while we do not know whether n, on the left hand side of the inequality grows to infinity.

Now we will return to our earlier inequalities.

$$\sigma(n) < e^{\gamma} \left(\log \log(n) \right) + \frac{0.6483n}{\log \log(n)}, \text{ for } n \ge 3$$

Since we have proven,
$$1.781n(\log\log(n)) + \frac{0.6483n}{\log\log(n)} \le e^{H_n}$$

Then it follows that, $\sigma(n) \leq e^{H_n}$

And,
$$\sigma(n) \leq e^{H_n} \leq H_n + \ln(H_n)e^{H_n}$$
, for $n \geq 3$

Therefore, we have proven that: $\sigma(n) \leq H_n + \ln(H_n)e^{H_n}$, for all $n \geq 1$, since we solved for $n \geq 5$

Thus, we have proven Lagarias's Elementary Version of the Riemann Hypothesis. And since it's proof is equivalent to Riemann Hypothesis, we have also proven Riemann Hypothesis.

The author must express many thanks to Bernhard Riemann who proposed the Riemann hypothesis in 1859. Again, the author thanks Bernhard Riemann for all of his work. Also the author wishes to express his eternal gratitude to Jeffrey Lagarias who proved that his Lagarias's Elementary Version is equivalent to the Riemann Hypothesis, a famous question about the

complex roots of the Riemann zeta function. Without Lagarias work, the author could not have proved the Riemann Hypothesis.

References:

- 1. An Elementary Problem Equivalent to the Riemann Hypothesis, Jeffery C. Lagarias, July 29, 2001.
- 2. G. Robin, Grandes valeurs de la fonction somme des diviseurs et hypothese de Riemann, J Math. Pures Appl. **63** (1984) 187-213.
- 3. Guy, Richard K., Unsolved problems in number theory, (3rd edition). Springer-Verlag (2004).
- 4. From Euclid to Present: A Collection of Proofs regarding the Infinitude of Primes, Lindsey Harrison, December 14, 2013.
- 5. (Dun) W. Dunham,"Journey Through Genius: The Great Theorems of Mathematics," Penguin Books, John Wiley and Sons, 1990.
- 6. Orders of Growth, Keith Conrad
- 7. Is two to the power of infinity more than infinity?, Igor Ostrovsky, April 2011