

Fundamental Solution of the Turbulence Problem avoiding Hypotheses

Rolf Warnemünde

© 2019 CC BY-NC-ND

Rolf Warnemünde

E-Mail: rolf.warnemuende@t-online.de

Abstract

Locality, natural causality and deterministics are the fundamental principles of this treatise. A dense fluctuating point set with unique linking to physical movements facilitates the setting up of partial differential equations. So a clear definition of a turbulent fluid is established. Stochasticity in the sense of an ensemble theory is considered by distributions of motion quantities of an unlimited number of parallelly existent deterministic systems. First, particle transport equations of

1. *Brownian motion as molecular self diffusion,*
2. *Stochastic transport by longitudinal fluctuations of a continuum,*
3. *Stochastic transport by turbulent continuum-fluctuations*

are developed. So Transition probabilities of moving quantities are evolved.

The connection of deterministics and stochastics in the sense of an ensemble theory enables a complete deterministic turbulence equation set. The result is pure geometrodynamics of turbulence on one side and pure geometrodynamics of deformation on the other side. At the end, the incorrectness of the known equation set of laminar fluid dynamics for turbulence problems is discussed.

Contents

Contents	4
1. Preface	8
2. Introduction	11
I. Prerequisites	15
3. Definition of a moved fluid-continuum	16
3.1. Introduction	16
3.2. Definition of moved fluid-elements	16
3.3. Laminar moved fluids	17
3.4. Turbulently moved fluids	18
4. Distribution functions	21
4.1. Introduction	21
4.2. Ensemble consideration of molecular self-diffusion	22
4.3. Ensemble consideration of stochastic particle transport in a continuum of longitudinal fluctuations	24
4.4. Ensemble consideration of stochastic particle transport in turbulently moved continua	25
4.5. Definition of Markov Processes with natural causality	27

II. Stochastically continuous transport of passive scalar particles within the meaning of an ensemble-theory	30
5. Introduction	31
6. Brownian motion as molecular self-diffusion	33
6.1. Introduction	33
6.2. Transport equation of molecular self-diffusion	33
6.3. Brownian motion as Markov Process with natural causality	38
6.4. Approximation formula	41
6.5. Appendix: equations for the spherical hamonics components	45
7. Stochastic transport by longitudinal fluctuations of a continuum	47
7.1. Introduction	47
7.2. Transport by Markov Processes with natural causality	48
7.3. Calculation of the exchange-term	51
7.4. Calculation of the exchange-coefficients Υ_l	53
7.5. Reconstruction of the transition probabilities $\overline{W}_{t_\epsilon}$	56
7.6. Approximation formula	58
8. Stochastic transport by turbulent continuum-fluctuations	60
8.1. Introduction	60
8.2. The transport as Markov Process with natural causality	61
8.3. Calculation of the exchange-term	64
8.4. Calculation of the exchange-coefficients Υ_l	66
8.5. Reconstruction of the transition probabilities $\overline{W}_{t_\epsilon}$	69
9. Appendix	73
9.1. Legendre-Polynomials	73
9.2. Spherical harmonics	75
9.3. Turbulence-functions	76

9.4. Euler-angles as fluctuation properties of the turbulent particle transport	77
III. Deterministic continuum-fluctuations and their stochastic view within the meaning of an ensemble-theory	79
10. Introduction	80
11. Deterministic turbulent mass-transport and its stochastic formulation	82
11.1. Introduction	82
11.2. The transition: stochastic theory \longleftrightarrow deterministic theory	82
11.3. The deterministic equations of turbulence	85
12. Stochastic and deterministic general vector fields	90
12.1. Introduction	90
12.2. The Transition: stochastic theory \longleftrightarrow deterministic theory	91
12.3. The deterministic fluctuation-equations	93
12.3.1. The vacuum Maxwell Equations	97
12.4. Surfacelike deformation-fluctuations in 3-dimensional space	98
12.5. 1-dimensional deformation-fluctuations in 3-dimensional space	99
13. Geometrodynamics of turbulence	101
13.1. Introduction	101
13.2. The complete set of turbulence-equations	103
13.3. Comments on the application of the complete equation system	105
13.4. The impossibility of calculating turbulence fields only knowing $\frac{\partial \vec{v}}{\partial t}(\vec{x}, t)$	107
13.5. Summary	109
13.6. Appendix A: The basic equations of laminar fluid-dynamics	109
13.6.1. The equation of continuity	109
13.6.2. The Navier-Stokes-equations	110
13.6.3. The equation of energy	112

CONTENTS

13.7. Appendix B	116
13.7.1. The Helmholtz-decomposition-theorem	116
14. Summary	118
Bibliography	120

1. Preface

After his graduation in physics the author has got the impression that all the essential problems in physics had been solved. Accordingly William K. George (Professor of Turbulence Emeritus) states: “*I remember despairing as a student that it seemed that all the problems were solved, and there was really nothing fundamental left to do.*” [10] At an advanced age, after decades of research and development in industry, this impression has fundamentally changed. Last but not least the frequently occurring confrontations with problems of turbulence for example in connection with pollutant propagation in turbulently moved atmosphere or heat exchange processes in turbulently moved boundary layers¹ has shown, that much remains to be done. Furthermore the usual interpretation of quantum mechanics (the probability problem) and the question of its completeness got to thinking, too.

The variety of different “turbulence theories“² would never have had a chance, to find a useful application in Industry. As for turbulence not even a trace of a useful definition exists. Although treatises (for instance theses) have been published by the most famous physicists of the 20th century like Heisenberg, it was nearly impossible for the industry concerning research and development to invest money and time. On the other hand hydrodynamics or rather turbulence didn’t play a significant role studying theoretical physics.

The question, how an optimal theory of physics is characterized is qualitatively easy answered but difficult to realize in a practical and specific way: It has to give a 1 to 1 mapping of reality. Anyway there has to be a suitable definition of that part of reality, whose processes are to be analyzed. Sometimes the necessary mathematics is not known. In this case one has to create formulations, which lead to calculable solutions. The identifying of their existence and definiteness may be sufficient for mathematics but for physics and especially industrial research they are not.

Theories in physics (especially in classical physics) are described by equations of motion with geometrodynamical connection (location, time, velocity and their space- und timelike partial derivations) and additional physical assumptions. These

¹Reynold’s law of similarity has proved to be entirely unsuitable in the case of turbulence.

²According to the opinion of the author: they all do not deserve this title.

assumptions include in the simplest case constants for instance the velocity of light in the Maxwell Equations of vacuum ³, or pure physical terms like in the Navier-Stokes-Equations of hydrodynamics, which describe the acceleration field of the fluid continuum by state variables of thermodynamics. (At this an appropriate definition of the continuum is somewhat difficult.) These equations are in the laminar flow range well confirmed by experiments, the applications beyond these limiting cases remain at least dubious. The degree of hypothesizing in such theories is least, if the mathematical formulation can be reduced to geometrodynamics without assumptions and without losing the **unique** reference to physical movements.

If it works calculating the velocity field of a heterogeneous mass as smeared mass distribution in a pure geometrodynamical way, the future smeared mass distribution can be determined exactly from the initially realized distribution.⁴ This is the result of the well known continuity equation:

$$\frac{\partial}{\partial t}\boldsymbol{\rho} + \vec{\nabla} \cdot (\boldsymbol{\rho} \cdot \vec{v}) = 0.$$

Calculations applying today's theory of hydrodynamics do not result in velocity fields of pure geometrodynamics. If the known hydrodynamic equations were correct ⁵, one could speak of a coupling of geometrodynamics and thermodynamics. Then there were an analogy to the theory of General Relativity, which means a coupling of space-time geometry with energy-momentum-density of matter. The development of suitable evolution equations in General Relativity is of great challenge for today's research.

Initially there was the intention to look for a mathematical formulation of deterministic transport processes of aerosols in turbulently moved fluids by a statistical transport theory within the meaning of an ensemble consideration. The developed transport equations would have been difficult to access experimentally, so that the second part of the research project arose, in which the relation of stochastic theory of natural causality with a suitable deterministic theory was established. Further theoretical researches of stochastic continuum fluctuations of 3-dimensional vector fields lead to pure geometrodynamics of turbulently fluctuating continua as well as fluctuating deformations. In doing so old line of thoughts of Einstein and Wheeler are not resumed. The resulting geometrodynamics is a consequent finding by avoiding

³these equations in this treatise are shown to be the consequence of geometrodynamics except the mentioned experimental velocity of light

⁴Generally, more cannot be achieved calculating a fluid. It requires defining a mathematical continuum corresponding to a discontinuously smeared matter

⁵Generally, they are not correct. Turbulence manifestations are the common case, laminar hydrodynamics is the limiting case.

hypotheses such that the unique relation of mathematical formulation and physical reference is not lost.

It is the opinion of the author, that the language of physics consists of two interacting parts-so to speak form and content-, whereby the form is linked to mathematical formulation and the content is linked to verbalized description.⁶ In this way the present theory is developed. It is plausible that in the act of unifying certain fields of physics a joint mathematical formulation should take shape.

Generally, classical physics is not the limiting case of quantum field theory ($\lim \hbar \rightarrow 0$) as maybe the limiting cases (of classical physics) Newton Mechanics and electrodynamics. In this treatise the electrodynamic field is shown to be a limiting case of formally classical physics but at the end of central importance. Einstein's General Relativity and the geometrodynamics of continuum physics of this treatise are no limiting cases of today's quantum field theory.

Overall, the author sees the way, which is pursued, as a path to reality in analogy to the book "The Road to Reality" of Penrose[24]. However Penrose doubts in the last chapter the correctness of the road outlined by himself (as it is nowadays generally tried in physics) not at least because the unification of Maxwell Field and Gravitational Field and quantization of the gravitational field failed. The author contends, that the prerequisite for understanding new developments is the understanding of their starting point, classical physics. Nowadays classical physics has serious shortcomings on one side and on the other side contains methods to ensure a natural causality that is the simultaneous consideration of position and momentum, which is lost with the so called modern physics (quantum mechanics, quantum field theory, etc.). This treatise wants to cure the main deficits of classical physics via the way of solving the turbulence problem, the main problem of continuum physics, avoiding usual hypotheses.

⁶the physicist is the moderator considering the interacting influences

2. Introduction

Feynman[9]: *“Nobody in physics has really been able to analyze it mathematically satisfactorily in spite of its importance to the sister sciences. It is the analysis of circulating or turbulent fluids.”*

The description of turbulent movements within the framework of continuum mechanics turned out to be difficult since more than 160 years. However, laminar fluid movements can be calculated by the known basic equations successfully confirmed in experiments: equation of continuity, Navier-Stokes-Equations and energy equation. The efforts, treating movements of turbulence in a similar way, must be considered as failures. There are substantial reasons for believing, that the above equations describing turbulent collective movements of **non-homogenously distributed molecular matter** are inadequate. This was the situation that inspired the idea, to explain the phenomenon of turbulence by stochastic methods. In that context, particularly approaches of Kolmogorov are to be mentioned, which lead to spectral energy distributions, assuming highly hypothetically, that turbulence is statistically isotropical und homogeneous. Between them there is a wide range of models with physically not well founded hypotheses. Overall, this leads to the statement of Feynman cited at the beginning, whereupon not much has changed since then.

This situation is characterized in recent treatises as for example by Trinh, Khanh Tuoc [16] in the following way:

“ the study of turbulence is immediately hampered by the surprising lack of a clear and concise definition of the physical process. Tsinober (2001) has published a long list of attempts at a definition by some of the most noted researchers in turbulence. The most common descriptions are vague: “a motion in which an irregular fluctuation (mixing, or eddying motion) is superimposed on the main stream” (Schlichting 1960), “a fluid motion of complex and irregular character” (Bayly, Orszag, Herbert, 1988)

or negative as in the breakdown of laminar flow (Reynolds' experiment 1883). Some of the definitions are quite controversial like Saffman's (1981) "One of the best definition of turbulence is that it is a field of random chaotic vorticity" because the words random and chaotic would imply that a formal mathematical solution, which is necessarily deterministic, does not exist. Perhaps the most accurate definition can be attributed to Bradshaw (1971) "The only short but satisfactory answer to the question "what is turbulence" is that it is the general-solution of the Navier-Stokes equation". This definition cannot be argued with but it is singularly unhelpful since no general solution of the NS yet exists 160 years after they were formulated."

or by McDonough [22] " In particular, a turbulent flow can be expected to exhibit all of the following features:

- 1. disorganized, chaotic, seemingly random behavior;*
- 2. nonrepeatability (i.e., sensitivity to initial conditions);*
- 3. extremely large range of length and time scales (but such that the smallest scales are still sufficiently large to satisfy the continuum hypothesis);*
- 4. enhanced diffusion (mixing) and dissipation (both of which are mediated by viscosity at molecular scales);*
- 5. three dimensionality, time dependence and rotationality (hence, potential flow cannot be turbulent because it is by definition irrotational);*
- 6. intermittency in both space and time."*

Fluctuation elements of the presented theory always form a dense point set, i.e. a definition of a continuum of such fluctuation elements is important deducing equations of motion in form of partial differential equations. On the other hand a concept of a stochastic theory of a fluctuating continuum within the meaning of an ensemble theory is deduced. Fundamental principles of this treatise as well as in the whole classical physics are **locality, causality and deterministics**. In this treatise particular emphasis is placed on specially defined **natural causality**, which in contrary to **Newtonian causality** of point mechanics only knows finite velocities. Discussed stochastics arises from statistics with an in thought experiment supposed unlimited ensemble of locally equivalent deterministic processes.¹ Finally turbulence is derived as a deterministic process and traced back to geometrodynamics of turbulence. The fluid element movements are described by interacting vortex- and vector-curvature

¹If one wants, it can be seen as a many-world-theory of classical physics. However this is done creating beforehand unknown equations of the deterministic processes (in contrary to Everett's many-world-theory of quantum mechanics).

fields, whereas geometrodynamics of deformation \vec{d} is characterised by a fluctuating curvature-tensor-field of 2nd degree, alternatively described by interacting

$$\frac{\partial \vec{d}}{\partial t}, \quad \text{rot}(\vec{d})$$

fields.

The interrelations of the deterministic and an associated stochastic theory enable a complete equation system of turbulently moved continua. The formulation of stochastically fluctuating processes of continua within the meaning of an ensemble theory is innovative for physics and mathematics. The known Navier-Stokes-Equations are not integrated in the complete equation system of turbulent moved fluids. The inclosed acceleration field $\frac{d\vec{v}}{dt}$ of the associated momentum equation can not be described by thermodynamic variables and viscose frictions.

$$\frac{d\vec{v}}{dt} \neq -\frac{1}{\rho} \vec{\nabla} p + \nu \Delta \vec{v} + (\zeta + \frac{\nu}{3}) \vec{\nabla} (\vec{\nabla} \cdot \vec{v})$$

The complete system of equations consists of 12 equations with 12 unknowns and contains only variables of motion in form of the vector fields: velocity, vortex, curvature and an acceleration field. So the developed theory of turbulence proves to be a geometrodynamics in a 3+1 dimensional Euclidian Space. Thermodynamics and matter distribution do not occur explicitly. This theory of variables of motion is principally exact. A smeared distribution of matter over Space-Time results by evaluation of the calculated velocity vector field and the equation of continuity. Currently, in a parallel treatise, this system of equations will be solved numerically for different physical situations: pipes, concentric pipes and rectangular channels. The numerical solutions are two times continuously differentiable in space coordinates.[28]

Some potential solutions for General Relativity and Cosmology result unconstrainedly. They are preceded by definitions of dense fluid continua and distribution functions for an unlimited ensemble of locally equivalent continua.

Generally, the background theory describes two geometrodynamics of different structures, each faced by a stochastic theory of continuum fluctuations with **natural causality**. No additional physical model considerations enter into the theory.²

Overall, the following problems are treated or solved respectively:

- Transport Theory of turbulently moved Particles

The stochastic transport equations considering consequently the continuum behaviour include coefficients, which stem from deterministic theory. The fluctuation variables are Euler Angles. These thoughts are important for example

²for example thermodynamics in the usual Navier-Stokes-Equations

considering pollutant dispersion in the lower atmosphere. The Theory offers several options of numerical evaluation depending on using the related partial differential equation, its presentation as integral equation or using particle trajectory-simulations in the related deterministic turbulence field (which can be calculated too).

- Continuum Theory of turbulent Fluctuations

The physics of turbulence is described by certain geometrodynamics for related turbulent movements. The associated stochastic description represents a Markov-Process with **natural causality**. A Markov Process as the stochastic description of turbulence is generally denied, because the understanding of such a stochastic process is too restricted. Formulating a transition probability for space-time-points

$$(\vec{x}, t) \longrightarrow (\vec{x}', t'),$$

is not enough. It depends on the velocity in (\vec{x}, t) reaching point (\vec{x}', t') . Arriving at the point (\vec{x}', t') has to be the consequence of a velocity in (\vec{x}, t) , which will be regulated by means of a transition probability of velocities factored out in vortex and related curvature vectors. The existence of a velocity as causal factor for transport in space-time we call **natural causality**. Explicitly, this is forbidden in quantum mechanics. That is why there are principal differences between turbulence theory on one side and quantum mechanics and quantum field theory on the other side. The phase space appearing usually in Boltzmannlike formulations is replaced by an in every point (\vec{x}, t) assumed velocity distribution of an ensemble of parallelly existing deterministic systems. Transition probabilities are explicitly calculated and are depending on the measurement accuracy of the measured motion quantities and degenerate exactly measured to δ -functions.

- Continuum Fluctuations of general 3-dimensional Vector Fields

Their physical interpretation enables for example the calculation of

- deformation fields (geometrodynamics of deformation)
- turbulence fields (geometrodynamics of turbulence)
- Maxwell Fields (they prove to be limiting cases of general deformation fields)

Part I.

Prerequisites

3. Definition of a moved fluid-continuum

3.1. Introduction

A proper definition of turbulence, which is based on a fluctuating, dense point set, does not exist. But this is necessary establishing equations of movement in form of partial differential equations. The known Navier-Stokes equations are only providing sufficient solutions for laminar problems. Below a fluctuating fluid is defined, which is associated uniquely to a dense set of space points of the considered time. This definition is the prerequisite developing stochastic theories of turbulent transport of continuously moved particles within the meaning of an ensemble theory, a deterministic theory and the connection of stochastic and a deterministic turbulence.

3.2. Definition of moved fluid-elements

At every time, space points (\vec{x}) are assigned to fluid elements in a unique correspondence. As this applies to every space point (\vec{x}) of the fluid field, the set of fluid elements is seen as a continuum. A Continuum of fluid element points (simply called fluid elements) is considered, where a fluid environment of non infinitesimal size is uniquely allocated to every fluid element point. Two infinitesimally neighboring fluid elements differ apart from their distance by their velocities and not quite identical material distributions of their neighborhoods. The neighborhoods of two nearby fluid elements overlap. A fluid element is shifted moving the material of its neighborhood. Though the material of such a fluid element may have changed marginally after an infinitesimal time interval t_ϵ , it can be identified principally by its prior material status. As every molecule possesses its own identity, there has to be at least an infinitesimally greater difference of material distribution to the neighborhoods of other fluid elements.

The neighborhoods exchange material with neighborhoods of adjacent fluid elements and vary their thermodynamic state (a local thermodynamic state does not necessarily exist). Their size is not infinitesimal, because a local thermodynamic state (if

physically existent) has to be detectable at least in thought experiment. The open neighborhoods have equally sized spherical shapes, generally. Near a solid border they are described by parts of spheres. Infinitesimally adjacent fluid elements possess overlapping neighborhoods. In an ε -surrounding they move in parallel. So one obtains a fluid, which is assumed to be a dense fluctuating point set, though there is no continuous matter distribution in Space-Time. That means it is possible to follow theoretically the history of every fluid element, though it has exchanged a lot of its initial material altering its local thermodynamic state.

Recapitulated:

Every space point (\vec{x}) of the open point set of a considered fluid area is at every time in unique correspondence to a fluid element. The fluid is an abstract, dense set of fluctuating fluid elements, which do not generally correspond to material points.

3.3. Laminar moved fluids

A continuum of moved fluid elements is considered each uniquely assigned to a neighborhood and a velocity.

$$\vec{v}_{t_\epsilon} = \frac{\vec{x}_2 - \vec{x}_1}{t_\epsilon} \quad (3.1)$$

The fluid elements move along sufficiently often continuously differentiable trajectories. The accuracies of the considered motion quantities are determined by t_ϵ -measurement processes t_ϵ characterising the accuracy. Deriving the transport equation of turbulent particle transport a limes consideration ($\lim t_\epsilon \rightarrow 0$) is subjected. The whole of the velocities create a velocity vector field having $\mathbf{rot}(\vec{v}) \neq \mathbf{0}$ generally.¹ Though $\mathbf{rot}(\vec{v})$ has dimension [1/sec], it does not refer to a rotation of laminar flow. In an infinitesimally surrounding area of a space-time-point (\vec{x}_0, t_0) a fluid flow can be defined locally² by parallelly moved fluid elements. Considering without loss of generality a fluid movement of velocity $\vec{v}(\vec{x}_0) = (v_x, 0, 0)$ in a space point \vec{x}_0 in cartesian coordinates, the velocity is described in an ϵ -neighborhood and parallel to the x-coordinate as follows:

$$\vec{v}(\vec{x}) = \begin{pmatrix} v_x(\vec{x}) \\ v_y(\vec{x}) \\ v_z(\vec{x}) \end{pmatrix} = \begin{pmatrix} v_x(\vec{x}_0) + \left. \frac{\partial v_x}{\partial x} \right|_{\vec{x}_0} \cdot \Delta x + \left. \frac{\partial v_x}{\partial y} \right|_{\vec{x}_0} \cdot \Delta y + \left. \frac{\partial v_x}{\partial z} \right|_{\vec{x}_0} \cdot \Delta z + \dots \\ \left. \frac{\partial v_y}{\partial x} \right|_{\vec{x}_0} \cdot \Delta x + \left. \frac{\partial v_y}{\partial y} \right|_{\vec{x}_0} \cdot \Delta y + \left. \frac{\partial v_y}{\partial z} \right|_{\vec{x}_0} \cdot \Delta z + \dots \\ \left. \frac{\partial v_z}{\partial x} \right|_{\vec{x}_0} \cdot \Delta x + \left. \frac{\partial v_z}{\partial y} \right|_{\vec{x}_0} \cdot \Delta y + \left. \frac{\partial v_z}{\partial z} \right|_{\vec{x}_0} \cdot \Delta z + \dots \end{pmatrix}$$

¹in english literature $\mathbf{curl}(\vec{v}) \neq \mathbf{0}$ is used but in turbulence the name \mathbf{rot} is more adapted as will be seen

²except in stagnation points

The velocity components $\mathbf{v}_y(\vec{\mathbf{x}})$ und $\mathbf{v}_z(\vec{\mathbf{x}})$ **osculate** at the velocity $\vec{\mathbf{v}}(\vec{\mathbf{x}}_0) = (v_x, 0, 0)$ spatially approaching (constant time t_0),

$$\begin{aligned}\mathbf{v}_y(x_0, y, z_0) &\longrightarrow \mathbf{v}_y(x_0, y_0, z_0) = \mathbf{0} \\ \mathbf{v}_z(x_0, y_0, z) &\longrightarrow \mathbf{v}_z(x_0, y_0, z_0) = \mathbf{0}\end{aligned}$$

That means especially, that all the partial derivations by y- or z-coordinate of 1. order of $\mathbf{v}_y(\vec{\mathbf{x}})$ and $\mathbf{v}_z(\vec{\mathbf{x}})$ disappear in the point (x_0, y_0, z_0) .

$$\lim_{z \rightarrow z_0} \frac{\Delta \mathbf{v}_y}{\Delta z} \Big|_{\vec{\mathbf{x}}_0} = \lim_{y \rightarrow y_0} \frac{\Delta \mathbf{v}_z}{\Delta y} \Big|_{\vec{\mathbf{x}}_0} = \mathbf{0} \quad (3.2)$$

$$\vec{\mathbf{x}}_0 = (x_0, y_0, z_0)$$

Applying the differential quotients in the $\vec{\nabla} \times$ -operator expressed in cartesian coordinates gives for the fluid velocity

$$(\vec{\nabla} \times \vec{\mathbf{v}})|_{\vec{\mathbf{x}}_0} = \begin{pmatrix} 0 \\ \frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \\ \frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \end{pmatrix} \Big|_{\vec{\mathbf{x}}_0}, \quad \vec{\mathbf{v}}(\vec{\mathbf{x}}_0) = (v_x, 0, 0) \quad (3.3)$$

The orthogonality of $\vec{\nabla} \times \vec{\mathbf{v}} \perp \vec{\mathbf{v}}$ is a fundamental quality ³⁴ and a necessary condition for continuous fluid flow.

In this orthogonality velocity vector fields differ from deformation vector fields.

3.4. Turbulently moved fluids

Trying to identify the state of movement of a fluid element in turbulent fluids by a velocity $\vec{\mathbf{v}}_{t_\epsilon}$ it should be recognized, that the state of movement is not yet determined, as the path in every space point (except in turning points) is uniquely adapted by an infinitesimal circle segment. In the infinitesimal neighborhood of a path point the velocity is identified by an instantaneous axis of rotation $\vec{\omega}_{t_\epsilon}$ and a radius vector $\vec{\mathbf{r}}_{t_\epsilon}$.⁵

$$\boxed{\vec{\mathbf{v}}_{t_\epsilon} = \vec{\omega}_{t_\epsilon} \times \vec{\mathbf{r}}_{t_\epsilon}} \quad (3.4)$$

³this relation can not be found in literature.

⁴This is one reason why the known millenium prize question does not lead to a solution of the turbulence problem. However the validity problem of the Navier-Stokes-equations is more fatal.

⁵That is why turbulence can not be uniquely identified by experiments of local velocity statistics.

The considered vectorial motion quantities $\vec{\omega}_{t_\epsilon}$ und \vec{r}_{t_ϵ} are determined by t_ϵ -measurement processes, which are calculated later on by a limes process $\lim t_\epsilon \rightarrow 0$. A fluid element originating from the point \vec{x}_0 crossing \vec{x}_1 after the time t_ϵ reaches \vec{x}_2 after a further time t_ϵ .

$$\vec{x}_0 \xrightarrow{t_\epsilon} \vec{x}_1 \xrightarrow{t_\epsilon} \vec{x}_2$$

By these 3 points a circle segment is uniquely drawn crossing point \vec{x}_1 with radius vector \vec{r}_{t_ϵ} and speed of rotation $\vec{\omega}_{t_\epsilon}$. The local state of motion can not be described by velocity only, neither statistically nor deterministically.⁶

Thus the fluid element in the space-time-point (\vec{x}, t) is identified principally by the contents of the matter of its neighborhood and state of movement expressed by $\vec{\omega}_{t_\epsilon}$ and \vec{r}_{t_ϵ} . In that way defined fluid elements move on sufficiently often continuously differentiable trajectories. They lead considering a continuum of fluctuating fluid elements to multiply continuously differentiable vector fields of motion. The continuum of moved fluid elements represent the turbulently collectiv movement of a discontinuously spaced Matter.

The field of turbulence is described by the two vector fields $\vec{\omega}_{t_\epsilon}$ and \vec{b}_{t_ϵ} ,

$$\vec{b}_{t_\epsilon} = \vec{r}_{t_\epsilon} / r_{t_\epsilon}^2 \quad \text{-curvature vector field.} \quad (3.5)$$

In addition, the results show that

$$\vec{\omega}_{t_\epsilon} = \frac{1}{2} \mathbf{rot}(\vec{v}_{t_\epsilon}). \quad (3.6)$$

$\mathbf{rot}(\vec{v})$ has the meaning of a local rotation in the frame of turbulence. An infinitesimal disturbance of stationary pipe flow leads to an change of the significance of $\mathbf{rot}(\vec{v})$, where $\mathbf{rot}(\vec{v})$ does not correspond to a rotation initially. Whether starting motions of turbulence are suppressed, depends on an existent viscosity. These decelerations are generally weak. The beginning of turbulent movements avoid Newtonian friction as well as pressure gradients by means of hereto orthogonal motions.

Vortex fields in turbulence (local rotation fields will be identified with vortex fields) and radius fields may have turning points (\vec{x}, t) along the paths of the fluid elements, which means $\vec{\omega} = \mathbf{0}$ und $\vec{r} = \infty$.⁷ In this case the velocities are to be calculated by interpolation or extrapolation of the neighborhood, for example. In the theory a further method will be shown. The fluid elements are accompanied by a moving frame of $\vec{\omega}, \vec{b}$ and \vec{v} along their paths.

In the following it is outlined, how locally Lagrangian and Eulerian formulations of fluid dynamics are reassembled in the turbulence theory. So deterministic considerations are found via stochastic descriptions, which could be designated as Lagrangian. Nevertheless, Lagrangian paths are calculated only after the determin-

⁶This statement contadicts that of [32]

⁷The temporal and spatial neighborhood of a turning point does not have such singular properties.



Figure 3.1.: Turbulences understood by Leonardo da Vinci

istic turbulence field is determined. These relations will become clear in later chapters.

4. Distribution functions

4.1. Introduction

Stochastically physical processes generally refer to random transports of physical quantities from $\vec{A} = (\vec{x}_1, t_1)$ to $\vec{B} = (\vec{x}_2, t_2)$, where a diffusion equation results at the end of all the discussions as can be seen in the well known treatise of Chandrasekhar [3]. This takes place in accordance with the Langevin equation, all known attempts characterizing Brownian motion and applying Fokker-Planck-equation, too. The diffusion equation is subjected to a Newtonian causality, that means the related propagation speed is unlimited. This is not the case in nonrelativistic physics beyond Newtonian mechanics, generally, as shown in the further course of this treatise.

In this context the Boltzmann Equation, which is only applicable for extensively diluted gases, constitutes a particularity. Despite surprising successes the importance of this equation is obviously not appropriately appreciated. In first approximation the Navier-Stokes equations are derived from this equation. A linear version can be classified as key-equation of nuclear reactor physics and is used for radioactive shielding problems in its stationary formulation.¹ These equations are based on a 6-dimensional phase-space with the apparent disadvantage, that using distribution functions $f(\vec{x}, t, \vec{v})$ a small but not infinitesimal phase space volume element $\Delta x \cdot \Delta y \cdot \Delta z \cdot \Delta v_x \cdot \Delta v_y \cdot \Delta v_z = \Delta V$ is to be believed surrounding the phase space point (\vec{x}, \vec{v}) . This situation is mathematically dissatisfying, as only a finite number of molecules can be existent inside this Volume, and executing $\lim \Delta V \rightarrow 0$ there remain no molecules representing $f(\vec{x}, t, \vec{v})$. Despite this contrariness the Boltzmann Equation, in general or linear form, is successful considering the results.²

Such a situation exists in other fields of physics, too. For example, no mathematically satisfying definition of a continuum is existent justifying partial differential equations like the Navier-Stokes-equations. Nevertheless they have performed satisfactorily in the case of laminar fluid dynamics, but extending to the general case of turbulence the known equations of laminar fluid dynamics fail. Deficient mathematical justification is sometimes balanced by experiments, not always. In the special case of

¹1968, associated with this the author has developed in his diploma thesis a numerical method (Monte-Carlo) for solving the linearly stationary Boltzmann Equation, without knowing this equation, simply by simulating the stochastic elementary processes.

²The Boltzmann Equation is the single equation describing the transition from a nonexisting local thermodynamic balance to a local thermodynamic balance.

kinetics, which is used for describing molecular self-diffusion, an ensemble theory is applied, which could be used for developing the Boltzmann Equation, too, avoiding the stated contrariness. On the other side the detailed mathematical formulations and their results do not alter by such a modified interpretation. But as phase space considerations are not possible for stochastically interpreted deterministic continuum fluctuations an equivalent treatment of ensemble theory will be used for all discussed problems.

The used distribution functions $f(\vec{x}, t, \vec{v})$ are not functions of the 6-dimensional phase space as usually applied in statistical mechanics but regular functions of space time with a probability density distribution of motion quantities in every point (\vec{x}, t) obtained by an unlimited ensemble of parallelly equivalent systems. In the case of the stationary linear radiation transport equation³ very different elementary particles like neutrons, electrons, α – particles, γ – particles etc. are simultaneously calculated by this equation.

A suitable ensemble-consideration is helpful to avoid Newtonian causality (see section 4.5), and to get rid of the mathematical inconsistency of ΔV of limited size with limited number of included particles. So mathematically not justified applications of related partial differential equations are avoided. This interpretation not altering the mathematical formulations in connection with gas kinetics the turbulence is lead to new relations.

4.2. Ensemble consideration of molecular self-diffusion

The specifically used construction may appear somehow artificially, but it is supposed to illustrate the classification of the usual diffusion equation as an approximate equation of a primary, with natural causality endowed transport equation. The particle density distributions are gained in thought experiment by an unlimited number of ensemble systems, which exist simultaneously. Their functions are sufficiently often, continuously differentiable in space and time. Regarding the quantities of motion the continuity condition is sufficient. This situation may be generated as follows:

An ensemble of parallel, extensively diluted monomolecular systems is considered to be in local thermodynamic ballance. They are all seen as statistically equivalent. They generally differ locally in an ϵ - neighborhood. Permitting for some

³This is a slight modification of the linear Boltzmann Equation with a streaming function $\frac{1}{v}f(\vec{x}, t, \vec{\Omega})$ instead of distribution function $f(\vec{x}, t, \vec{v})$, and the different velocities before and after molecular collisions are considered in the differential cross sections of the collision integral only [31].

time equivalent molecules to enter in all systems by equally distributed sources such that the additional quantity of gas is insignificant in relation to the original quantity of the gas, the additional part of the gas will be in the same statistical balance in all the systems of the ensemble after a short time, though it has not reached a homogenous distribution. While the added part of gas consists in every single system of a limited number of molecules only, it is possible to formulate a sufficiently often continuously differentiable particle density distribution $f(\vec{x}, \vec{v}, t)$ for the added gas part, as the statistics of the velocities relates to the whole ensemble.

The expectation value of a suitable particle density may be constructed as follows. Around the point (\vec{x}, t) an equal-sized volume $\Delta V_{(\vec{x}, t)}$ is chosen out of all representatives μ of the ensemble in which a subset ΔN_μ of molecules is located. μ identifies a single representative of the ensemble. μ passing all values from 1 to ∞ an unlimited number of ensemble-representatives is taken into account. The expectation value of a particle density of this ensemble-consideration in point (\vec{x}, t) in the small volume $\Delta V_{(\vec{x}, t)}$ of the neighborhood of (\vec{x}, t) results in

$$\langle \rho(\vec{x}, t) \rangle_{\Delta V} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mu=1}^n \frac{\Delta N_\mu}{\Delta V_{(\vec{x}, t)}}. \quad (4.1)$$

Contracting the volume $\Delta V_{(\vec{x}, t)}$ to the point (\vec{x}, t) one has

$$\langle \rho(\vec{x}, t) \rangle = \lim_{\Delta V_{(\vec{x}, t)} \rightarrow 0} \langle \rho(\vec{x}, t) \rangle_{\Delta V}. \quad (4.2)$$

This function of expectation values is sufficiently often continuously differentiable in its depending variables, especially in space and time. In accordance with the velocities, consisting of amount and direction of motion, their distribution density is separated in these quantities as follows ⁴

$$\boxed{\begin{aligned} \langle \rho(\vec{x}, t) \rangle &= \int_0^\infty \int_{4\pi} f(\vec{x}, t, \vec{v}) d\vec{\Omega} dv = \int_{4\pi} h(\vec{x}, t, \vec{\Omega}) d\vec{\Omega} \\ f(\vec{x}, t, \vec{v}) &= f(\vec{x}, t, v\vec{\Omega}) = h(\vec{x}, t, \vec{\Omega}) g(v) \\ \int_0^\infty g(v) dv &= 1, \quad \bar{v} = \int_0^\infty g(v) v dv \\ \vec{v} &= v \cdot \vec{\Omega}. \end{aligned}} \quad (4.3)$$

So the necessary connection is given by

⁴A separation ansatz $f(\vec{x}, t, v\vec{\Omega}) = h(\vec{x}, t, \vec{\Omega})g(\vec{x}, t, v\vec{\Omega})$ is generally to be chosen.

$$h(\vec{x}, t, \vec{\Omega}) = \int_0^\infty f(\vec{x}, t, \mathbf{v}\vec{\Omega}) dv. \quad (4.4)$$

This enables the derivation of a transport equation of $h(\vec{x}, t, \vec{\Omega})$ the amounts of the velocities occurring as constant coefficients only.

At this the distribution of the velocity amounts $\mathbf{g}(\mathbf{v})$ is separated from the direction distribution $h(\vec{x}, t, \vec{\Omega})$. The diffusing particles possess a gaussian distribution and the equipartition law is applied. This is the prerequisite for deriving a suitable transport equation below and afterwards in less than first approximation a diffusion equation with constant diffusion coefficients.

This special distribution of the velocity amounts (equipartition law) corresponds to the assumed situation of molecular self-diffusion of chapter 6. The gained expectation value of the density does not exactly equal the value measured in an ensemble representative performed in a small volume. Thus one has

$$\langle \rho(\vec{x}, t) \rangle \approx \rho(\vec{x}, t). \quad (4.5)$$

An exact measurement (this has nothing to do with a measurement of the macroscopic state quantity density) would result in

$$\langle \rho(\vec{x}, t) \rangle \neq \rho(\vec{x}, t) = \begin{cases} 1 & \text{one particle existent in point } (\vec{x}, t) \\ 0 & \text{else} \end{cases}. \quad (4.6)$$

4.3. Ensemble consideration of stochastic particle transport in a continuum of longitudinal fluctuations

Similarly, further considerations occur to section 4.2. Around the point (\vec{x}, t) an equal-sized volume $\Delta \mathbf{V}_{(\vec{x}, t)}$ is chosen out of all representatives μ of the ensemble in which a subset $\Delta \mathbf{N}_\mu$ of particles ⁵ is located. μ identifies a single representative of the ensemble. The expectation value of a particle density results in

$$\langle \rho(\vec{x}, t) \rangle = \lim_{\Delta \mathbf{V}_{(\vec{x}, t)} \rightarrow 0} \left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mu=1}^n \frac{\Delta \mathbf{N}_\mu}{\Delta \mathbf{V}_{(\vec{x}, t)}} \right). \quad (4.7)$$

⁵the particles have to be of a size adapting the identical movement of present fluid element of the fluctuating continuum

This function of expectations is sufficiently often continuously differentiable in its variables of space and time. By the separation of the velocity in amount and direction the distribution density is described as follows

$$\begin{aligned}
 < \rho(\vec{x}, t) > = \int_0^\infty \int_{4\pi} f(\vec{x}, t, \vec{v}) d\vec{\Omega} dv = \int_{4\pi} \bar{f}(\vec{x}, t, \vec{\Omega}) d\vec{\Omega} \\
 f(\vec{x}, t, \vec{v}) &= G(\vec{x}, t, v \vec{\Omega}) \bar{f}(\vec{x}, t, \vec{\Omega}) \\
 \int_0^\infty G(\vec{x}, t, v \vec{\Omega}) dv &= 1, \quad \bar{v}(\vec{x}, t, \vec{\Omega}) = \int_0^\infty G(\vec{x}, t, v \vec{\Omega}) v dv \\
 \vec{v} &= v \cdot \vec{\Omega}.
 \end{aligned} \tag{4.8}$$

So the necessary combination

$$\bar{f}(\vec{x}, t, \vec{\Omega}) = \int_0^\infty f(\vec{x}, t, v \vec{\Omega}) dv. \tag{4.9}$$

is achieved deriving a transport equation for $\bar{f}(\vec{x}, t, \vec{\Omega})$. In this transport equation the velocities occur as coefficients of the averaged velocity amounts in dependence of space, time and direction.

The gained expectation value of the density fails to comply with the measured ensemble representative, that is

$$< \rho(\vec{x}, t) > \neq \rho(\vec{x}, t) \tag{4.10}$$

For a single ensemble representative the distribution function f degenerates to a delta-function

$$f \rightarrow \delta(\vec{v}_{(\vec{x}, t)}, \vec{v}) \tag{4.11}$$

with

$$\begin{aligned}
 \int_{\vec{v}} \delta(\vec{v}_{(\vec{x}, t)}, \vec{v}) d\vec{v} &= 1 \\
 \int_{\vec{v}} \delta(\vec{v}_{(\vec{x}, t)}, \vec{v}) \vec{v} d\vec{v} &= \vec{v}_{(\vec{x}, t)}.
 \end{aligned} \tag{4.12}$$

4.4. Ensemble consideration of stochastic particle transport in turbulently moved continua

The fluid fluctuations of different ensemble-representatives may arise at the same time by equivalent macroscopic-physical processes with different infinitesimal perturbations. That is why different fluent motions are created in (\vec{x}, t) in every of the

parallel-systems. The simultaneous release of passive particles retracing uniquely the motions of fluid movements may have taken place by a distribution of similar point sources in all parallel systems. So an own particle distribution is developed in every individual system in space-time. The statistical recording running over the whole ensemble leads to continuously differentiable distribution functions of a limited number of particles ⁶ in a single system.

Further considerations follow analogously to section 4.2. Around the point (\vec{x}, t) an equal-sized volume $\Delta \mathbf{V}_{(\vec{x}, t)}$ is chosen out of all representatives μ of the ensemble in which a subset $\Delta \mathbf{N}_\mu$ of particles is located. μ identifies a single representative of the ensemble. The expectation value of a particle density results in

$$\langle \rho(\vec{x}, t) \rangle = \lim_{\Delta \mathbf{V}_{(\vec{x}, t)} \rightarrow 0} \left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mu=1}^n \frac{\Delta \mathbf{N}_\mu}{\Delta \mathbf{V}_{(\vec{x}, t)}} \right). \quad (4.13)$$

This function of expectation values arises out of a distribution function f of motion quantities

$$\begin{aligned} \vec{\omega} &= \vec{\omega}(\vec{x}, t) && \text{rotation speed} \\ \vec{r} &= \vec{r}(\vec{x}, t) && \text{radius vector} \\ \vec{v} &= \vec{\omega}(\vec{x}, t) \times \vec{r}(\vec{x}, t) && \text{velocity vector} \end{aligned} \quad (4.14)$$

that means

$$f = f(\vec{x}, t, \vec{\omega}, \vec{r}). \quad (4.15)$$

A separation results in

$$\begin{aligned} \langle \rho(\vec{x}, t) \rangle &= \int_{\vec{r}} \int_{\vec{\omega}} f(\vec{x}, t, \vec{\omega}, \vec{r}) d\vec{\omega} d\vec{r} = \int_{2\pi} \int_{4\pi} \bar{f}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) d\vec{\Omega} d\vec{\Theta} \\ f(\vec{x}, t, \vec{\omega}, \vec{r}) &= G(\vec{x}, t, \vec{\omega}, \vec{r}) \bar{f}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \\ \int_0^\infty \int_0^\infty G(\vec{x}, t, \omega \vec{\Omega}, r \vec{\Theta}) d\omega dr &= 1, \quad \int_0^\infty \int_0^\infty G(\vec{x}, t, \omega \vec{\Omega}, r \vec{\Theta}) \omega r d\omega dr = \bar{v}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \\ \int_0^\infty \int_0^\infty G(\vec{x}, t, \omega \vec{\Omega}, r \vec{\Theta}) r d\omega dr &= \bar{r}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}), \quad \int_0^\infty \int_0^\infty G(\vec{x}, t, \omega \vec{\Omega}, r \vec{\Theta}) \omega d\omega dr = \bar{\omega}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \\ \bar{v}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \bar{\omega}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \cdot \bar{r}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \\ \bar{v}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \bar{\omega}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \times \bar{r}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}). \end{aligned}$$

(4.16)

Such the necessary combination is given by

⁶the particles have to be of a size adapting the identical movement of present fluid element of the fluctuating continuum

$$\bar{f}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) = \int_0^\infty \int_0^\infty f(\vec{x}, t, \vec{\omega} \cdot \vec{\Omega}, \vec{r} \cdot \vec{\Theta}) d\omega dr. \quad (4.17)$$

This enables a transport equation of $\bar{f}(\vec{x}, t, \vec{\Omega}, \vec{\Theta})$ with the averaged amounts of rotation velocities and radius-vectors as coefficients.

The resulting expectation value of the density does not equal the measured value of a single ensemble representative. That is

$$\langle \rho(\vec{x}, t) \rangle \neq \rho(\vec{x}, t) \quad (4.18)$$

Limiting to one system of the ensemble the distribution function degenerates to a delta-function

$$f \rightarrow \delta(\vec{\omega}_{(\vec{x}, t)}, \vec{r}_{(\vec{x}, t)}; \vec{\omega}, \vec{r}) \quad (4.19)$$

with

$$\begin{aligned} \int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x}, t)}, \vec{r}_{(\vec{x}, t)}; \vec{\omega}, \vec{r}) d\vec{\omega} d\vec{r} &= 1 \\ \int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x}, t)}, \vec{r}_{(\vec{x}, t)}; \vec{\omega}, \vec{r}) \vec{\omega} d\vec{\omega} d\vec{r}' &= \vec{\omega}_{(\vec{x}, t)} \\ \int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x}, t)}, \vec{r}_{(\vec{x}, t)}; \vec{\omega}, \vec{r}) \vec{r} d\vec{\omega} d\vec{r}' &= \vec{r}_{(\vec{x}, t)}. \end{aligned} \quad (4.20)$$

4.5. Definition of Markov Processes with natural causality

The probabilistic theory is related to random distributions of velocities $\vec{\pi}$ moving from (\vec{x}, t) to $(\vec{x} + \vec{\pi}t_\epsilon, t + t_\epsilon)$. These velocity distributions may get together of vortex and curvature vector fields

$$\vec{\pi} = \vec{\omega} \times \frac{\vec{b}}{b^2}.$$

The transport from $(\vec{x} - t_\epsilon \vec{\pi}', t - t_\epsilon)$ to (\vec{x}, t) is additionally controlled by transition probabilities

$$W_{t_\epsilon} = W_{t_\epsilon}(\vec{x}, t, \vec{\pi}, \vec{\pi}'),$$

resulting in

$$f_{t_\epsilon}(\vec{x}, t, \vec{\pi}) = \int_{\vec{\pi}'} W_{t_\epsilon}(\vec{x}, t, \vec{\pi}, \vec{\pi}') f_{t_\epsilon}(\vec{x} - t_\epsilon \vec{\pi}', t - t_\epsilon, \vec{\pi}') d\vec{\pi}'.$$

Such a relation we call a Markov Process of natural causality. According to Sen [26] there is a so called Newtonian causality in nonrelativistic physics implying the possibility of unlimited velocities. However Newtonian causality is restricted to Newtonian mechanics and stochastic processes of physics ending with diffusion equations when applied practically.⁷ This applies not for formulations of the general or linear Boltzmann Equation. In electrodynamics the velocity of light is the limiting velocity. In this treatise one essential statement is: classical physics is generally not Newtonian. Further on

1. is shown, that diffusion equations can only be approximations of an exact description. The diffusion equation is related to an unlimited propagation speed. The diffusion coefficient is correlated with the velocity of sound. Exact descriptions lead via Boltzmannlike formulations.
2. is shown, that the second Newtonian law applies to fluid dynamics in limiting cases only. In field theories as fluid dynamics not force- but acceleration fields are expressed. These are generally not free of rot (equivalently curl) in contrary to a Newtonian force field. That is why it is reasonable to distinguish conservative from non conservative acceleration fields. In classical physics one has normally non conservative fields. (Though for students a contrary impression may occur.)

The Newtonian causality proves to be a limiting case of non relativistic classical physics. Subsequently a causal Markov Process is continuously used or derived. Overarching master equations can not exist, physically. The transition probabilities W_{t_ϵ} depend on a time quantity t_ϵ related to continuum fluctuations of measurement accuracy according to vectorial motion quantities. For $t_\epsilon \rightarrow 0$ (exact motion quantities) the transition probability W_{t_ϵ} degenerates to a δ -function.

Simultaneous details of space and momentum are not possible in the context of quantum mechanics. The Schrödinger Equation for free particles

$$i\hbar \frac{\partial \psi(\vec{x}, t)}{\partial t} = -\frac{\hbar^2}{2\mu} \vec{\nabla}^2 \psi(\vec{x}, t) \quad (4.21)$$

can be transformed into a linear homogenous integral equation [12] [14]

⁷This statement applies to the Fokker-Planck and Langevin equation. See, for example, Chandrasekhar[3]

$$\psi(\vec{x}, t) = i \int G(\vec{x}, t; \vec{x}', t') \psi(\vec{x}', t') d\vec{x}'. \quad (4.22)$$

The Green function

$$G(\vec{x}, t; \vec{x}', t') = \left\langle \vec{x} \left| \exp\left(-\frac{i}{\hbar}(t - t')\mathbf{H}\right) \right| \vec{x}' \right\rangle \quad (4.23)$$

is called Feynman kernel, too.

In the case of the diffusion equation

$$\frac{\partial \rho(\vec{x}, t)}{\partial t} = D \vec{\nabla}^2 \rho(\vec{x}, t) \quad (4.24)$$

an equivalent integral equation the Green function understood as transition probability from (\vec{x}', t') to (\vec{x}, t) exists with

$$\rho(\vec{x}, t) = \int_{V'} G(\vec{x}, t; \vec{x}', t') \rho(\vec{x}', t') d\vec{x}' \quad (4.25)$$

and the Green function

$$G(\vec{x}, t; \vec{x}', t') = \left(\frac{1}{4\pi D(t - t')} \right)^{\frac{3}{2}} e^{-\frac{(\vec{x} - \vec{x}')^2}{4\pi D(t - t')}}. \quad (4.26)$$

Equations based on a "heat-kernel"-structure are not exact in classical physics (as well as the Newtonian mechanics).

In quantum mechanics and quantum field theory natural causality is not possible because of the uncertainty principle. In Relativity there is the maximal possible velocity, the velocity of light. A geometrodynamic equation system of turbulence found further down does not contain such limiting velocities, explicitly. Velocity fields are calculated uniquely by an initial field giving to GR compatible results after mapping from Einstein Space into a suitable observer-space. Using other initial conditions higher velocities are possible.

Part II.

Stochastically continuous transport
of passive scalar particles within
the meaning of an ensemble-theory

5. Introduction

$$\begin{aligned}
\frac{\partial \bar{f}}{\partial t} + \bar{\omega} \cdot \bar{r} \vec{\Omega} \times \vec{\Theta} \cdot \nabla \bar{f} &= \frac{-1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \frac{l(l+1)}{2} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}) \\
&\Downarrow \\
\bar{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, \vec{\Theta}, t) &= \int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}, \vec{\Omega}', \vec{\Theta}') \bar{f}_{t_\epsilon}(\vec{x} - t_\epsilon \cdot \vec{v}' \vec{\Omega}' \times \vec{\Theta}', \vec{\Omega}', \vec{\Theta}'), t - t_\epsilon) d\vec{\Omega}' d\vec{\Theta}'
\end{aligned}$$

The aim is the derivation of a stochastic transport equation of turbulent, passively moved scalar particles being based on particle balancing. Stochastics is always understood as the randomness of motions. From point \vec{A} originated a point \vec{B} is approached in consequence of a random velocity. The particle motions are totally adjusted to the fluid motions of the fluctuating continuum and reproduce single fluid motions in detail.¹ The used stochastics is based for one thing on an ensemble-consideration and on the other hand on a locally formulated motion process. So an equation is achieved owning local coefficients depending on space and time as well as the states of movement. The field of coefficients can be determined in principal in every desired level of detail by the deterministic turbulence theory of chapter 13. The transport equation is a partial differential equation shown to be equivalent to a derived integral equation. The respective stochastic process is immediately recognized as Markovian of natural causality. We call it causal Markov Process.

Chapter 6: Within the framework of kinetic theory a physical situation is selected handling the linear Boltzmann Equation. This equation is extensively studied in nuclear reactor physics called neutron Boltzmann Equation[31].² Using the above described ensemble consideration a statistical particle balance is formulated by local velocities and their unsteady changes by local cross sections. The resulting mathematical ties help the developments in further chapters as guideline and answer the question, which analogies exist between kinetic theory and turbulent stochastic continuum transport.

¹The particles are assumed to have a suitable weight

²The insights of this theory had little impact on similar fields of physics.

Chapter 7: The motion of passive scalar particles by longitudinal continuum fluctuations is examined. In the centre of the consideration is the development of transition probability densities of velocities. They depend as well as the velocities and their particle density distributions on the accuracy of a measuring process indexed by t_ϵ . $\lim t_\epsilon \rightarrow 0$ means exact measurements and the transition probabilities result into δ -functions. They have the property of test functions of the distribution theory with immediate physical meaning. Calculating them a transport equation in form of a partial differential equation as well as an equivalent integral equation is derived.

Chapter 8: In analogy to chapter 7 the motion by turbulent continuum fluctuations of passive particles is examined. The fluctuation directions are expressed by Eulerian angles and the distribution functions are developed by generalized spherical harmonics (we call them turbulence functions). A pair of equations is created consisting of a partial differential equation and an equivalent integral equation as in the cases of molecular self diffusion and the longitudinal (1+3)-dimensional continuum fluctuations. The three physical situations can be compared all the more as in the three cases the transition probabilities are explicitly formulated.

6. Brownian motion as molecular self-diffusion

$$\begin{aligned} \frac{\partial}{\partial t} h + \vec{v} \cdot \vec{\Omega} \nabla h &= \frac{1}{\tau} \sum_{l=1}^{+\infty} \gamma_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \\ h_{t_\epsilon}(\vec{x}, \vec{v}_{t_\epsilon} \vec{\Omega}, t) &= \int_{4\pi} \int_0^\infty W_{t_\epsilon}(\vec{\Omega}, v' \vec{\Omega}') h_{t_\epsilon}(\vec{x} - v' \vec{\Omega}' t_\epsilon, v' \vec{\Omega}', t - t_\epsilon) dv' d\vec{\Omega}' \end{aligned}$$

6.1. Introduction

Brownian motion is understood as a disordered thermic motion of molecules in gases or fluids creating a disordered motion of suspended, sufficiently small particles. This Brownian motion is all the more livelier the smaller the particle quantity is. With increasing particle sizes the detailed molecular influence on the particle movement disappears and having suitable sizes the particles reproduce the turbulent fluid motions. The phenomenon of small particles was first examined by Einstein and Smoluchowski.

Subsequently, the case of very small particles that is the statistical development of the molecular distribution of a gas is evaluated. In the treatises of Einstein[8] and Smoluchowski[27] the considerations lead in each case to a diffusion equation, which contains two fundamental deficiencies, though being sufficient for the purpose at that time. The propagation speed concerning a diffusion equation is unlimited. Immediately after switching a point particle source on there is at least an infinitesimal influence in arbitrary distance. In close proximity to a point source the solution of a diffusion equation shows a $\sim \frac{1}{r}$ -behaviour. But it should be $\sim \frac{1}{r^2}$.

6.2. Transport equation of molecular self-diffusion

Examining the molecular self-diffusion in a highly diluted gas in thermodynamic equilibrium the linear Boltzmann equation will be derived. It is a linear integro-differential equation statistically describing the transport of diffusing particles by

cross sections of the interacting particles. The whole gas medium is regarded as divided into two parts, a main part and an additional very small part. The diffusing of the small part in the main part without changing the statistical properties of the main part is considered. (See section 4.2) Due to the low density of the diffusing molecules a relevant self interaction within the small part can be excluded. Regarding the spatiotemporal development the velocity distribution density $\mathbf{g}(\mathbf{v})$ is normalised to 1.

$$\int_0^\infty \mathbf{g}(\mathbf{v}) d\mathbf{v} = 1 \quad (6.1)$$

The diffusing part is depicted by

$$f(\vec{\mathbf{x}}, \vec{\mathbf{v}}, t) = f(\vec{\mathbf{x}}, t, v\vec{\Omega}) = h(\vec{\mathbf{x}}, t, \vec{\Omega})\mathbf{g}(\mathbf{v}) \quad (6.2)$$

I.e. the velocity distribution is independent of space-time $(\vec{\mathbf{x}}, t)$ and direction $(\vec{\Omega})$ (equipartition theorem).

This allows to talk about an expectation value for every space-time point of the particle density

$$\langle \Phi(\vec{\mathbf{x}}, t) \rangle = \int_0^\infty \int_{4\pi} f(\vec{\mathbf{x}}, t, \vec{\mathbf{v}}) d\vec{\Omega} dv = \int_{4\pi} h(\vec{\mathbf{x}}, t, \vec{\Omega}) d\vec{\Omega}. \quad (6.3)$$

However the measured value of the density $\Phi(\vec{\mathbf{x}}, t)$ is only a good approximation of the expectation value

$$\langle \Phi(\vec{\mathbf{x}}, t) \rangle \approx \Phi(\vec{\mathbf{x}}, t). \quad (6.4)$$

This ceases to apply for the particle transport by fluctuating continua.

The total derivative of the distribution function f in direction of the velocity $v\vec{\Omega}$ results in

$$\frac{d}{dt} f(\vec{\mathbf{x}}, t, v\vec{\Omega}) = \frac{\partial}{\partial t} f + v\vec{\Omega} \nabla f. \quad (6.5)$$

The change of the particle density distribution for the velocity $\vec{\mathbf{v}} = v\vec{\Omega}$ is balanced by collisions of molecules modifying the velocities with a certain probability expressed by differential cross sections. Defining $\frac{1}{v}f(\vec{\mathbf{x}}, \vec{\mathbf{v}}, t)$ as particle stream¹ the following

¹Hereby no stream in the meaning of deterministic fluid dynamics is defined!

balance equation can be noted

$$\frac{1}{v} \frac{d}{dt} f(\vec{x}, t, v\vec{\Omega}) = \frac{1}{v} \frac{\partial}{\partial t} f + \vec{\Omega} \nabla f = I_+ - I_- \quad (6.6)$$

I_+ corresponds to molecules coming from other directions $\vec{\Omega}'$.
 I_- corresponds to molecules leaving direction $\vec{\Omega}$.

The particle distribution density varying in space this expression has to be different from zero. Otherwise the particle distribution density remains constant. So the assumed initial distribution is variously dispersed in space.

The momentum exchange is determined on one side by the cross sections of the impact partners and on the other side by the number of particles arriving at location \vec{x} and time t per unit area with the velocity \vec{v}' pivoting into the velocity \vec{v} . This growth of the number of particles per time and unit-area is

$$I_+ = \rho \int_0^\infty \int_{4\pi} \sigma(\vec{v} \cdot \vec{v}') \cdot f(\vec{x}, t, v'\vec{\Omega}') dv' d\vec{\Omega}' \quad (6.7)$$

with

$$\begin{aligned} \rho &= \text{constant density of the main part of the gas} \\ \sigma(\vec{v} \cdot \vec{v}') &= \text{differential cross section, symmetrical in } \vec{v} \text{ and } \vec{v}'. \end{aligned}$$

The particles simultaneously changing their velocity \vec{v} into another \vec{v}' the appropriate decrease of particle number per time- and area-unit is expressed by

$$I_- = \rho \int_0^\infty \int_{4\pi} \sigma(\vec{v} \cdot \vec{v}') \cdot f(\vec{x}, t, v\vec{\Omega}) dv d\vec{\Omega} = \Sigma(v) f(\vec{x}, t, v\vec{\Omega}) \quad (6.8)$$

due to

$$\Sigma(v) = \rho \int_0^\infty \int_{4\pi} \sigma(\vec{v} \cdot \vec{v}') dv' d\vec{\Omega}' \quad [m^{-1}]. \quad (6.9)$$

$\Sigma(v)$ represents the total macroscopic cross section and the transport equation results in the molecular self diffusion equation

$$\boxed{\frac{1}{v} \frac{\partial}{\partial t} f + \vec{\Omega} \cdot \nabla f = \rho \int_0^\infty \int_{4\pi} \sigma(\vec{v} \cdot \vec{v}') \cdot f(\vec{x}, t, v' \vec{\Omega}') dv' d\vec{\Omega}' - \Sigma(v) f}. \quad (6.10)$$

Further on, considering the molecules being in statistical balance throughout the whole gas, i.e. the same Boltzmann Distribution $\mathbf{g}(\mathbf{v})$ is existing everywhere, an integration of $\int_0^\infty (6.10) dv$ results in a manageable equation as follows.

Defining

$$\bar{\Sigma} = \int_0^\infty \Sigma(v) g(v) dv \quad (6.11)$$

$$\bar{\sigma}(\vec{\Omega} \cdot \vec{\Omega}') = \int_0^\infty \int_0^\infty \sigma(\vec{v} \cdot \vec{v}') g(v') dv dv' \quad (6.12)$$

$$\bar{v} = \int_0^\infty v g(v) dv \quad (6.13)$$

gives²

$$\frac{\partial}{\partial t} h + \bar{v} \vec{\Omega} \cdot \nabla h = \bar{v} \rho \int_{4\pi} \bar{\sigma}(\vec{\Omega} \cdot \vec{\Omega}') \cdot h(\vec{x}, t, \vec{\Omega}') d\vec{\Omega}' - \bar{v} \bar{\Sigma} \cdot h(\vec{x}, t, \vec{\Omega}). \quad (6.14)$$

Developing by spherical harmonics (see appendix 9.2) yield in

$$\bar{\sigma}(\vec{\Omega} \cdot \vec{\Omega}') = \sum_{l=0}^{+\infty} \sigma_l P_l(\cos(\alpha)) = \sum_{l=0}^{+\infty} \sigma_l \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}') \quad (6.15)$$

and

$$\begin{aligned} h(\vec{x}, t, \vec{\Omega}) &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} h_{lm}(\mathbf{x}, t) P_{lm}(\vec{\Omega}) \\ &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} h_{lm}(\mathbf{x}, t) P_{lm}^*(\vec{\Omega}). \end{aligned} \quad (6.16)$$

²Such an equation corresponds in nuclear reactor physics to the one group neutron-transport-equation regardless of absorptions-and fission effects.

These developments inserted into (6.14) and executing the respective integrations lead to

$$\begin{aligned} \frac{\partial}{\partial t} h + \bar{v} \vec{\Omega} \cdot \nabla h &= \rho \bar{v} \sum_{l=1}^{+\infty} \sigma_l \frac{4\pi}{2l+1} \sum_{m=-l}^{m=+l} h_{lm} P_{lm}^*(\vec{\Omega}) - \bar{v} \bar{\Sigma} \cdot \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \\ &= \sum_{l=1}^{+\infty} \bar{v} \left\{ \rho \sigma_l \frac{4\pi}{2l+1} - \bar{\Sigma} \right\} \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \end{aligned} \quad (6.17)$$

It holds

$$\bar{\Sigma} = 4\pi \rho \sigma_0 = \rho \int_{4\pi} \bar{\sigma}(\vec{\Omega} \cdot \vec{\Omega}') d\vec{\Omega}' \quad [m^{-1}] \quad (6.18)$$

and defining

$$\begin{aligned} \tau^{-1} &= \bar{v} \bar{\Sigma} \quad [sec^{-1}] \\ \gamma_l &= \left(\frac{\sigma_l}{\sigma_0} \frac{1}{2l+1} - 1 \right) < 0 \quad \text{für } l \geq 1 \quad [/] \end{aligned} \quad (6.19)$$

$$\implies \gamma_0 = 0 \quad (6.20)$$

one gets

$$\boxed{\frac{\partial}{\partial t} h + \bar{v} \vec{\Omega} \cdot \nabla h = \frac{1}{\tau} \sum_{l=1}^{+\infty} \gamma_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega})}. \quad (6.21)}$$

As $\gamma_l \leq 0$, the development components of order l are the more rapidly decaying the l becoming greater. The total derivation in the direction of the velocity $\bar{v} \vec{\Omega}$ leads to

$$\frac{d}{dt} h(\vec{x}, t, \vec{\Omega}) = \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} \frac{d}{dt} h_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}) = \frac{1}{\tau} \sum_{l=1}^{+\infty} \gamma_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}). \quad (6.22)$$

That is why the time behaviour of the single development components result in

$$\frac{d}{dt} h_{lm}(t) = \frac{\gamma_l}{\tau} h_{lm} \quad (6.23)$$

and

$$h_{lm}(t) \sim \exp\left(\frac{\gamma_l}{\tau} \cdot t\right). \quad (6.24)$$

So approximations of first order turn out to approach exact solutions, asymptotically.³

6.3. Brownian motion as Markov Process with natural causality

Defining the transition probability density of directions

$$\boxed{\overline{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') = \sum_{l=0}^{+\infty} \frac{2l+1}{4\pi} e^{+\Upsilon_l \cdot \frac{t_\epsilon}{\tau}} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}')} \quad (6.25)}$$

and determining the following relationships

$$\epsilon = \frac{t_\epsilon}{\tau}, \quad \frac{1}{\tau} = \bar{v} \cdot \bar{\Sigma} = \bar{v} \cdot 4\pi\rho\sigma_0 = \text{const} \quad (6.26)$$

$$\bar{\sigma}(\vec{\Omega} \cdot \vec{\Omega}') = \sum_{l=0}^{+\infty} \sigma_l P_l(\cos(\alpha)) = \sum_{l=0}^{+\infty} \sigma_l \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}') \quad (6.27)$$

$$\Upsilon_l = \left(\frac{\sigma_l}{\sigma_0} \cdot \frac{1}{2l+1} - 1 \right) \quad (6.28)$$

an integral equation of self-diffusion results in dependence of directions of motions, cross sections and locally averaged absolute values of velocities as coefficients,

$$\boxed{h_{t_\epsilon}(\vec{x}, t, \vec{\Omega}) = \int_{4\pi} \overline{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') \cdot h_{t_\epsilon}(\vec{x} - \bar{v}\vec{\Omega}'t_\epsilon, t - t_\epsilon, \vec{\Omega}') d\vec{\Omega}'} \quad (6.29)}$$

from which equation

$$\frac{\partial}{\partial t} h + \bar{v}\vec{\Omega} \cdot \nabla h = \frac{1}{\tau} \sum_{l=1}^{+\infty} \Upsilon_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \quad (6.21)$$

³This is correct in the case of lacking absorption processes. We are only regarding scattering.

may be reconstructed.

Proof:

h_{t_ϵ} developed around \vec{x} and t until first order one gets

$$h_{t_\epsilon}(\vec{x} - \vec{v}'_{t_\epsilon} \cdot t_\epsilon, \vec{\Omega}', t - t_\epsilon) = h_{t_\epsilon}(\vec{x}, \vec{\Omega}', t) - \tau \cdot \epsilon \cdot \left[\frac{\partial h'_{t_\epsilon}}{\partial t} + \vec{v}'_{t_\epsilon} \cdot \vec{\nabla} h'_{t_\epsilon} + O(\epsilon^2) \right] \quad (6.30)$$

with $t_\epsilon = \tau \cdot \epsilon$. Inserted into (6.29) this leads to

$$h_{t_\epsilon} = \int_{4\pi} \overline{W}_{t_\epsilon} h'_{t_\epsilon} d\vec{\Omega}' - \int_{4\pi} \overline{W}_{t_\epsilon} \cdot \tau \cdot \epsilon \cdot \left[\frac{\partial h'_{t_\epsilon}}{\partial t} + \vec{v}'_{t_\epsilon} \cdot \vec{\nabla} h'_{t_\epsilon} + O(\epsilon^2) \right] d\vec{\Omega}' \quad (6.31)$$

and simple conversions give

$$\frac{\int_{4\pi} \overline{W}_{t_\epsilon} h'_{t_\epsilon} d\vec{\Omega}' - h_{t_\epsilon}}{\epsilon} = \int_{4\pi} \overline{W}_{t_\epsilon} \cdot \tau \cdot \left[\frac{\partial h'_{t_\epsilon}}{\partial t} + \vec{v}'_{t_\epsilon} \cdot \vec{\nabla} h'_{t_\epsilon} + O(\epsilon^2) \right] d\vec{\Omega}'. \quad (6.32)$$

Executing the limiting process $t_\epsilon \rightarrow 0$ the transition probability $\overline{W}_{t_\epsilon}$ results in a δ -function and the particle density distribution h_{t_ϵ} achieves the limiting function h .

$$\begin{aligned} \lim_{t_\epsilon \rightarrow 0} \overline{W}_{t_\epsilon} &= \delta(\vec{\Omega}, \vec{\Omega}') \quad \text{:delta-Function} \\ \lim_{t_\epsilon \rightarrow 0} h_{t_\epsilon} &= h \\ \lim_{t_\epsilon \rightarrow 0} \vec{v}_{t_\epsilon} &= \vec{v} = \vec{v} \cdot \vec{\Omega} \\ \lim_{t_\epsilon \rightarrow 0} \vec{\Omega}_{t_\epsilon} &= \vec{\Omega} \end{aligned} \quad (6.33)$$

Executing the limiting process $t_\epsilon \rightarrow 0$ on equation (6.32) the t_ϵ -indexing disappears in accordance with the distribution functions.

Before the limes process is carried out the following integrations lead to

$$\begin{aligned}
 & \int_{4\pi} \bar{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') h_{t_\epsilon}(\vec{x}, t, \vec{\Omega}') d\vec{\Omega}' = \\
 & \int_{4\pi} \left[\sum_{l=0}^{+\infty} \frac{2l+1}{4\pi} e^{+\Upsilon_l \cdot \frac{t_\epsilon}{\tau}} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') \right] h_{t_\epsilon}(\vec{x}, t, \vec{\Omega}') d\vec{\Omega}' \\
 & = \int_{4\pi} \left[\sum_{l=0}^{+\infty} \frac{2l+1}{4\pi} e^{+\Upsilon_l \cdot \frac{t_\epsilon}{\tau}} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') \right] \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} h_{t_\epsilon lm}(\mathbf{x}, t) P_{lm}^*(\vec{\Omega}') d\vec{\Omega}' \\
 & = \sum_{l=0}^{+\infty} e^{+\Upsilon_l \cdot \frac{t_\epsilon}{\tau}} \sum_{m=-l}^{m=+l} h_{t_\epsilon lm}(\mathbf{x}, t) P_{lm}^*(\vec{\Omega}).
 \end{aligned} \tag{6.34}$$

Thus one gets

$$\begin{aligned}
 & \frac{\int_{4\pi} \bar{W}_{t_\epsilon} h'_{t_\epsilon} d\vec{\Omega}' - h_{t_\epsilon}}{\epsilon \cdot \tau} \\
 & = \frac{\int_{4\pi} \bar{W}_{t_\epsilon} h'_{t_\epsilon} d\vec{\Omega}' - \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} h_{t_\epsilon lm}(\mathbf{x}, t) P_{lm}^*(\vec{\Omega})}{\epsilon \cdot \tau} \\
 & = \frac{\sum_{l=0}^{+\infty} (e^{+\Upsilon_l \cdot \frac{t_\epsilon}{\tau}} - 1) \sum_{m=-l}^{m=+l} h_{t_\epsilon lm}(\mathbf{x}, t) P_{lm}^*(\vec{\Omega})}{\epsilon \cdot \tau}.
 \end{aligned} \tag{6.35}$$

Setting

$$\Upsilon_l = \lim_{t_\epsilon \rightarrow 0} \frac{e^{+\Upsilon_l \cdot \frac{t_\epsilon}{\tau}} - 1}{\epsilon}, \quad t_\epsilon = \epsilon \cdot \tau \tag{6.36}$$

creates

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \bar{W}_{t_\epsilon} h'_{t_\epsilon} d\vec{\Omega}' - h_{t_\epsilon}}{\epsilon \cdot \tau} = \frac{1}{\tau} \sum_{l=1}^{+\infty} \Upsilon_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \tag{6.37}$$

and

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \bar{W}_{t_\epsilon} h'_{t_\epsilon} d\vec{\Omega}' - h_{t_\epsilon}}{\epsilon \cdot \tau} = \frac{\partial h}{\partial t} + \vec{v} \cdot \vec{\nabla} h, \tag{6.38}$$

which results in

$$\boxed{\frac{\partial h}{\partial t} + \bar{v} \vec{\Omega} \cdot \vec{\nabla} h = \frac{1}{\tau} \sum_{l=1}^{+\infty} \Upsilon_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega})}. \quad (6.39)}$$

q.e.d.

Extending the transition probability density \bar{W} by the velocity distribution $g(v')$

$$W_{t_\epsilon}(\vec{\Omega}, v' \vec{\Omega}') = g(v') \bar{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}'), \quad (6.40)$$

one gets the Brownian molecular motion under the terms of the described model in the most general form.

$$\boxed{f_{t_\epsilon}(\vec{x}, v_{t_\epsilon} \vec{\Omega}, t) = \int_{4\pi} \int_0^\infty W_{t_\epsilon}(\vec{\Omega}, v' \vec{\Omega}') f_{t_\epsilon}(\vec{x} - v' \vec{\Omega}' t_\epsilon, v' \vec{\Omega}', t - t_\epsilon) dv' d\vec{\Omega}'} \quad (6.41)$$

The transition probabilities are not symmetric in contrary to the differential cross section!

6.4. Approximation formula

An approximation formula of 1. order of the equation

$$\frac{\partial}{\partial t} h + \bar{v} \vec{\Omega} \cdot \nabla h = \frac{1}{\tau} \sum_{l=1}^{\infty} \gamma_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}) \quad (6.42)$$

is beeing looked for. The approach accounts for the methods of the transport theory of nuclear reactor physics [31]. In cartesian coordinates this leads to

$$\frac{\partial}{\partial t} h + \bar{v} \cdot \left(\Omega_x \frac{\partial}{\partial x} h + \Omega_y \frac{\partial}{\partial y} h + \Omega_z \frac{\partial}{\partial z} h \right) = \frac{1}{\tau} \sum_{l=1}^{\infty} \gamma_l \cdot \sum_{m=-l}^{+l} h_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}). \quad (6.43)$$

$$\begin{aligned}
\Omega_x &= \sin(\vartheta)\cos(\varphi) \\
\Omega_y &= \sin(\vartheta)\sin(\varphi) \\
\Omega_z &= \cos(\vartheta)
\end{aligned} \tag{6.44}$$

The spherical harmonics of 0th and 1st Order are

$$\begin{aligned}
P_{00} &= 1 & P_{1-1} &= 2^{-\frac{1}{2}}e^{-i\varphi}\sin\vartheta & P_{10} &= \cos\vartheta & P_{11} &= -2^{-\frac{1}{2}}e^{i\varphi}\sin\vartheta \\
P_{00}^* &= 1 & P_{1-1}^* &= 2^{-\frac{1}{2}}e^{+i\varphi}\sin\vartheta & P_{10}^* &= \cos\vartheta & P_{11}^* &= -2^{-\frac{1}{2}}e^{-i\varphi}\sin\vartheta.
\end{aligned} \tag{6.45}$$

In cartesian coordinates until 1st order this leads to

$$\frac{\partial}{\partial t}h + \bar{v} \cdot \left(\Omega_x \frac{\partial}{\partial x}h + \Omega_y \frac{\partial}{\partial y}h + \Omega_z \frac{\partial}{\partial z}h \right) = \frac{1}{\tau} \gamma_1 \cdot \sum_{m=-1}^{+1} h_{1m}(\vec{x}, t) P_{1m}(\vec{\Omega}) \tag{6.46}$$

The direction vectors in cartesian coordinates expressed by spherical harmonics are written

$$\begin{aligned}
\Omega_x &= 2^{-\frac{1}{2}}[P_{1-1} - P_{11}] \\
\Omega_y &= -i2^{-\frac{1}{2}}[P_{1-1} + P_{11}] \\
\Omega_z &= P_{10}.
\end{aligned} \tag{6.47}$$

The transport equation in 1st approximation is reduced to

$$\begin{aligned}
&\frac{\partial}{\partial t}(h_{00}P_{00} + h_{1-1}P_{1-1} + h_{10}P_{10} + h_{11}P_{11}) \\
&+ \bar{v} \left[2^{-\frac{1}{2}}[P_{1-1} - P_{11}] \frac{\partial}{\partial x}(h_{00}P_{00} + h_{1-1}P_{1-1} + h_{10}P_{10} + h_{11}P_{11}) \right. \\
&- i2^{-\frac{1}{2}}[P_{1-1} + P_{11}] \frac{\partial}{\partial y}(h_{00}P_{00} + h_{1-1}P_{1-1} + h_{10}P_{10} + h_{11}P_{11}) \\
&\left. + P_{10} \frac{\partial}{\partial z}(h_{00}P_{00} + h_{1-1}P_{1-1} + h_{10}P_{10} + h_{11}P_{11}) \right] \\
&= \frac{1}{\tau} \gamma_1 \cdot (h_{1-10}P_{1-1} + h_{10}P_{10} + h_{11}P_{11}).
\end{aligned} \tag{6.48}$$

After integrating $\int (6.48 P_{lm}^*(\vec{\Omega}) d\vec{\Omega})$ for $l = 0, 1$ the evolution equation set until 1st

order

$$\frac{\partial h_{00}}{\partial t} + \frac{\bar{v}}{3} \left[2^{-\frac{1}{2}} \left(-\frac{\partial h_{11}}{\partial x} + \frac{\partial h_{1-1}}{\partial x} \right) - i 2^{-\frac{1}{2}} \left(\frac{\partial h_{11}}{\partial y} + \frac{\partial h_{1-1}}{\partial y} \right) + \frac{\partial h_{10}}{\partial z} \right] = 0 \quad (6.49)$$

$$\frac{\partial h_{10}}{\partial t} + \bar{v} \frac{\partial h_{00}}{\partial z} - \frac{\Upsilon_1}{\tau} h_{10} = 0 \quad (6.50)$$

$$\frac{\partial h_{1-1}}{\partial t} + \bar{v} 2^{-\frac{1}{2}} \left(\frac{\partial h_{00}}{\partial x} + i \frac{\partial h_{00}}{\partial y} \right) - \frac{\Upsilon_1}{\tau} h_{1-1} = 0 \quad (6.51)$$

$$\frac{\partial h_{11}}{\partial t} + \bar{v} 2^{-\frac{1}{2}} \left(-\frac{\partial h_{00}}{\partial x} + i \frac{\partial h_{00}}{\partial y} \right) - \frac{\Upsilon_1}{\tau} h_{11} = 0 \quad (6.52)$$

is approached.

Now we define a vector field \vec{J} .

$$\begin{aligned} J_x &= \frac{4\pi}{3} 2^{-\frac{1}{2}} h_{(1-1} - h_{11}) \\ J_y &= -i \frac{4\pi}{3} 2^{-\frac{1}{2}} (h_{1-1} + h_{11}) \\ J_z &= \frac{4\pi}{3} h_{10} \\ \Phi &= 4\pi h_{00} \end{aligned} \quad (6.53)$$

Insertion (6.53) into (6.49) gives

$$\frac{\partial \Phi}{\partial t} + \bar{v} \cdot \vec{\nabla} \cdot \vec{J} = 0. \quad (6.54)$$

Φ is the particle density of an in a thought experiment assumed small part of the molecular set. Inserting (6.53) into (6.50) until (6.52) leads to

$$\vec{J} = \frac{\tau}{\Upsilon_1} \left[\frac{\bar{v}}{3} \vec{\nabla} \Phi + \frac{\partial \vec{J}}{\partial t} \right] \quad (6.55)$$

with $\Upsilon_1 = \left(\frac{1}{3} \frac{\sigma_1}{\sigma_0} - 1 \right) = -\eta$

So a telegrapher's equation arises⁴

$$\frac{\tau}{\eta} \frac{\partial^2 \Phi}{\partial t^2} + \frac{\partial \Phi}{\partial t} = \frac{\tau}{\eta} \bar{v} \vec{\nabla} \cdot \frac{\bar{v}}{3} \vec{\nabla} \Phi. \quad (6.56)$$

⁴To derive telegrapher's equation relativistic considerations are not necessary as is stated in [6].
The propagation speed is closely connected with the speed of sound.

As η , τ und \bar{v} represent constants, the telegrapher's equation is written

$$\boxed{\begin{aligned} \frac{\tau}{\eta} \frac{\partial^2 \Phi}{\partial t^2} + \frac{\partial \Phi}{\partial t} &= D \Delta \Phi \\ \text{with } D &= \frac{\tau \bar{v}^2}{\eta 3} \end{aligned}} \quad (6.57)$$

D =diffusion coefficient, η dimensionless, $\tau = (\bar{v} \cdot \bar{\Sigma})^{-1}$ = mean free collision time

\bar{v} =mean amount of velocity

Compared to the 1st derivation the term with temporal derivation of 2nd order can normally be neglected.

The dependence of the diffusion coefficient from macroscopic state variables of an ideal gas may happen as follows:

The equation of state of the ideal gas becomes

$$p = \rho R T. \quad (6.58)$$

The mean quadratic velocity of a Maxwellian velocity distribution of particles with mass m is [5]

$$\overline{v^2} = \frac{3kT}{m} \quad (6.59)$$

\Rightarrow

$$\bar{v} = \sqrt{\frac{8}{\pi} \frac{kT}{m}} = \sqrt{\frac{8}{\pi} \frac{p}{\rho}} \quad (6.60)$$

m means the mass of a molecule.

k is the Boltzmann constant.

To get a comparison with the speed of sound at a Gaussian velocity distribution

$$c = \sqrt{\left. \frac{\partial p}{\partial \rho} \right|_T} \quad (6.61)$$

one obtains

$$\bar{v} = \sqrt{\frac{8}{\pi}} c \quad (6.62)$$

and

$$\boxed{D = \frac{1}{\eta \bar{\Sigma}} \frac{8}{3\pi} c^2}. \quad (6.63)$$

The propagation speed for Brownian molecular motion is \bar{v} . This in particular becomes apparent by equation (6.29). In connection with the diffusion approximation an unlimited propagation speed is assigned. This leads to solutions approaching asymptotically to those of the linear Boltzmann Equation. In close proximity to point sources (less than 3 average free lengths afar⁵) one obtains the following characteristics of the exact and the diffusional solution.

$$\Phi \sim \frac{1}{r^2} \quad \text{solution of the transport equation in the proximity of a point source} \quad (6.64)$$

Anticipating this result from an exact theory appears directly plausible.

$$\Phi \sim \frac{1}{r} \quad \text{solution of the diffusion approximation in the proximity of a point source} \quad (6.65)$$

Avoiding such deficiencies it is necessary to take a stochastic velocity distribution into account as root of the diffusion process. Analyzing turbulent particle transport this does not satisfy.

6.5. Appendix: equations for the spherical hamonics components

The general equations arise out of

$$\int (6.43) P_{lm}^*(\vec{\Omega}) d\vec{\Omega} \quad (6.66)$$

\Rightarrow

⁵An experience of the neutron transport theory [31]

$$\begin{aligned}
\frac{\partial h_{lm}(\vec{x}, t)}{\partial t} = & -\bar{v} \left[\frac{\sqrt{(l+2+m)(l+1+m)}}{2l+3} \left(-\frac{1}{2} \frac{\partial h_{l+1,m+1}}{\partial x} - \frac{i}{2} \frac{\partial h_{l+1,m+1}}{\partial y} \right) \right. \\
& + \frac{\sqrt{(l+1-m)(l+2-m)}}{2l+3} \left(\frac{1}{2} \frac{\partial h_{l+1,m-1}}{\partial x} - \frac{i}{2} \frac{\partial h_{l+1,m-1}}{\partial y} \right) \\
& + \frac{\sqrt{(l-1-m)(l-m)}}{2l-1} \left(\frac{1}{2} \frac{\partial h_{l-1,m+1}}{\partial x} + \frac{i}{2} \frac{\partial h_{l-1,m+1}}{\partial y} \right) \\
& + \frac{\sqrt{(l+m)(l+m-1)}}{2l-1} \left(-\frac{1}{2} \frac{\partial h_{l-1,m-1}}{\partial x} + \frac{i}{2} \frac{\partial h_{l-1,m-1}}{\partial y} \right) \\
& \left. + \frac{\sqrt{(l+1+m)(l-m+1)}}{2l+3} \frac{\partial h_{l+1,m}}{\partial z} + \frac{\sqrt{(l+m)(l-m)}}{2l-1} \frac{\partial h_{l-1,m}}{\partial z} \right] - \frac{\Upsilon_l}{\tau} h_{lm}
\end{aligned} \tag{6.67}$$

7. Stochastic transport by longitudinal fluctuations of a continuum

$$\boxed{\begin{aligned} \bar{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, t) &= \int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}') \bar{f}_{t_\epsilon}(\vec{x} - t_\epsilon \cdot \vec{v}' \vec{\Omega}', \vec{\Omega}', t - t_\epsilon) d\vec{\Omega}' \\ &\quad \Updownarrow \\ \frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \cdot \vec{\nabla} \bar{f} &= -\frac{1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \frac{l(l+1)}{2} \bar{f}_{l,m}(\vec{x}, t) P_{l,m}(\vec{\Omega}) \end{aligned}}$$

7.1. Introduction

The motion of passive particles by longitudinal continuum fluctuations is examined. The particles are moved in this field without interaction.¹ In accordance with section 4.3 they perform detailed motions of single fluid elements of fluid continua. The considered velocities of the particles are determined by measure processes. The particles coming from point x_1 and moving further for a time t_ϵ are detected in x_2 . So the velocity \vec{v}_{t_ϵ} may be assigned to

$$\vec{v}_{t_\epsilon} = \frac{\vec{x}_2 - \vec{x}_1}{t_\epsilon} = v_{t_\epsilon} \vec{\Omega}_{t_\epsilon}. \quad (7.1)$$

This corresponds to $(\vec{x}_1, t) \longrightarrow (\vec{x}_2, t + t_\epsilon) = (\vec{x}_1 + \vec{v}_{t_\epsilon} \cdot t_\epsilon, t + t_\epsilon)$.

According to an ensemble consideration (see chapter 4) for every point (\vec{x}, t) a continuously differentiable particle density distribution of velocities \vec{v}_{t_ϵ} is assigned in accordance with

$$f_{t_\epsilon} = f_{t_\epsilon}(\vec{x}, \vec{v}_{t_\epsilon}, t). \quad (7.2)$$

¹Such conditions generally lead to linear equations.

The functions indexed with t_ϵ or ϵ enclose motion quantities \vec{v}_{t_ϵ} or their motion directions $\vec{\Omega}_{t_\epsilon}$ as variables subordinated to an understanding of measurement accuracy. The indexing of motion quantities with t_ϵ or ϵ may be dropped if their functions are indexed. Executing a limiting process, for instance

$$\lim_{t_\epsilon \rightarrow 0} f_{t_\epsilon}(\vec{x}, \vec{v}, t) = f(\vec{x}, \vec{v}, t) \quad (7.3)$$

f and \vec{v} are literally understood as results of exact measurement processes.² Integrating the particle density distribution over the velocity one obtains an expectation value of a particle density not generally coinciding with the actually measured value ρ .

$$\langle \rho_{t_\epsilon}(\vec{x}, t) \rangle = \int_{4\pi} \int_0^\infty f_{t_\epsilon}(\vec{x}, v\vec{\Omega}, t) dv d\vec{\Omega} \neq \rho_{t_\epsilon}(\vec{x}, t) \quad (7.4)$$

This is contradicting the molecular self-diffusion being an inherent stochastic process.

It results into a rigorously derived partial differential equation calculating particle density distributions in dependence on space-time and motion directions. The initially unlimited number of unknown coefficients is reduced to one, a local time-scaling. The initially abstractly formulated transition probabilities obtain their precise functional dependencies alternatively generating an integral equation. For numerical solutions there are always suitable Monte-Carlo methods possible.

Equations of 1st approximation substantially differ from usual diffusion equations.

7.2. Transport by Markov Processes with natural causality

The probability particles at location \vec{x} and time t changing their velocity from $\vec{v}'_{t_\epsilon} = v'\vec{\Omega}'$ to $\vec{v}_{t_\epsilon} = v\vec{\Omega}$ is given by the transition probability

$$W_{t_\epsilon} = W_{t_\epsilon}(\vec{x}, t, v\vec{\Omega}, v'\vec{\Omega}') \quad (7.5)$$

with

$$\int_{4\pi} \int_0^\infty W_{t_\epsilon}(\vec{x}, t, v\vec{\Omega}, v'\vec{\Omega}') dv' d\vec{\Omega}' = 1. \quad (7.6)$$

²With the indexing t_ϵ an assigned measurement process is always understood according to accuracy.

So the following Markov Process is defined by

$$f_{t_\epsilon}(\vec{x}, v\vec{\Omega}, t) = \int_{4\pi} \int_0^\infty W_{t_\epsilon}(\vec{x}, t, v\vec{\Omega}, v'\vec{\Omega}') f_{t_\epsilon}(\vec{x} - t_\epsilon \cdot v'\vec{\Omega}', v'\vec{\Omega}', t - t_\epsilon) dv' d\vec{\Omega}' \quad (7.7)$$

$$\Delta \vec{x} = \vec{x} - \vec{x}_1 = v'_{t_\epsilon} \vec{\Omega}'_{t_\epsilon} \cdot t_\epsilon \quad (7.8)$$

For the transition probability W_{t_ϵ} merely steadiness is required regarding all variables. The sequence of the velocities $\vec{v}'_{t_\epsilon}, \vec{v}_{t_\epsilon}$ means a motion

$$(\vec{x} - \vec{v}'_{t_\epsilon} \cdot t_\epsilon, t - t_\epsilon, \vec{v}'_{t_\epsilon}) \longrightarrow (\vec{x}, t, \vec{v}_{t_\epsilon}). \quad (7.9)$$

At the process $t_\epsilon \rightarrow 0$ the transition probabilities W_{t_ϵ} prove to be physical realisations of test functions of distribution theory.

The passive particles have to reproduce the motions of the fluctuation field, exactly. For the particle density distribution $f_{t_\epsilon}(\vec{x}, t, \vec{v})$ a separation approach is formulated without restriction of generality:

$$f_{t_\epsilon}(\vec{x} - v\vec{\Omega} \cdot t_\epsilon, v\vec{\Omega}, t) = G_{t_\epsilon}(\vec{x} - v\vec{\Omega} \cdot t_\epsilon, v\vec{\Omega}, t) \bar{f}_{t_\epsilon}(\vec{x} - \bar{v}\vec{\Omega} \cdot t_\epsilon, \vec{\Omega}, t) \quad (7.10)$$

$$\int_0^\infty G_{t_\epsilon}(\vec{x}, v\vec{\Omega}, t) dv = 1$$

\implies

$$\bar{f}_{t_\epsilon}(\vec{x} - \bar{v}\vec{\Omega} \cdot t_\epsilon, \vec{\Omega}, t) = \int_0^\infty f_{t_\epsilon}(\vec{x} - v\vec{\Omega} \cdot t_\epsilon, v\vec{\Omega}, t) dv \quad (7.11)$$

This results in

$$\bar{\mathbf{v}} = \bar{\mathbf{v}}(\vec{x}, t, \vec{\Omega}) = \int_0^\infty G_{t_\epsilon}(\vec{x}, v\vec{\Omega}, t) \cdot \mathbf{v} dv. \quad (7.12)$$

I.e. $\bar{\mathbf{v}}$ is dependent on $(\vec{x}, t, \vec{\Omega})$.

A transition probability only in dependence on the directions and space-time \bar{W}_{t_ϵ} is obtained by integration of W_{t_ϵ} over the velocity amounts v'_{t_ϵ} and v_{t_ϵ}

$$\boxed{\overline{W}_{t_\epsilon}(\vec{x}, \vec{\Omega}, \vec{\Omega}', t) = \int_0^\infty \int_0^\infty W_{t_\epsilon}(\vec{x}, t, v\vec{\Omega}, v'\vec{\Omega}') G_{t_\epsilon}(\vec{x} - v'\vec{\Omega}' \cdot t_\epsilon, v'\vec{\Omega}', t - t_\epsilon) dv' dv.} \quad (7.13)$$

Now an integration of $\int_0^\infty (7.7) dv$ is leading to

$$\overline{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, t) = \int_0^\infty \int_0^\infty \int_{4\pi} W_{t_\epsilon} G_{t_\epsilon}(\vec{x} - v'\vec{\Omega}' \cdot t_\epsilon, v'\vec{\Omega}', t - t_\epsilon) \overline{f}_{t_\epsilon}(\vec{x} - \vec{v}'\vec{\Omega}' \cdot t_\epsilon, \vec{\Omega}', t - t_\epsilon) dv' dv d\vec{\Omega}' \quad (7.14)$$

respectively

$$\boxed{\overline{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, t) = \int_{4\pi} \overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}') \overline{f}_{t_\epsilon}(\vec{x} - \vec{v}'\vec{\Omega}' \cdot t_\epsilon, \vec{\Omega}', t - t_\epsilon) d\vec{\Omega}'.} \quad (7.15)$$

$\overline{f}_{t_\epsilon}$ in the integrand is developed about \vec{x} and t until 1st order and one obtains

$$\begin{aligned} \overline{f}_{t_\epsilon}(\vec{x} - \vec{v}'\vec{\Omega}' \cdot t_\epsilon, \vec{\Omega}', t - t_\epsilon) &= \overline{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}', t) - \tau_E \cdot \epsilon \cdot \left[\frac{\partial \overline{f}_{t_\epsilon}'}{\partial t} + \vec{v}'_{t_\epsilon} \cdot \vec{\Omega}'_{t_\epsilon} \cdot \vec{\nabla} \overline{f}_{t_\epsilon}' + O(\epsilon^2) \right] \\ \overline{f}_{t_\epsilon}' &= \overline{f}_{t_\epsilon}'(\vec{x}, \vec{\Omega}', t) \end{aligned} \quad (7.16)$$

with $t_\epsilon = \tau_E \cdot \epsilon$ and $\tau_E = \text{const.}$ Inserted in (7.15) this leads to

$$\overline{f}_{t_\epsilon} = \int_{4\pi} \overline{W}_{t_\epsilon} \overline{f}_{t_\epsilon}' d\vec{\Omega}' - \int_{4\pi} \overline{W}_{t_\epsilon} \cdot \tau_E \cdot \epsilon \cdot \left[\frac{\partial \overline{f}_{t_\epsilon}'}{\partial t} + \vec{v}'_{t_\epsilon} \cdot \vec{\nabla} \overline{f}_{t_\epsilon}' + O(\epsilon^2) \right] d\vec{\Omega}'. \quad (7.17)$$

and simple conversions give

$$\frac{\int_{4\pi} \bar{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' - \bar{f}_{t_\epsilon}}{\epsilon} = \int_{4\pi} \bar{W}_{t_\epsilon} \cdot \tau_E \left[\frac{\partial \bar{f}'_{t_\epsilon}}{\partial t} + \vec{v}'_{t_\epsilon} \cdot \vec{\nabla} \bar{f}'_{t_\epsilon} + O(\epsilon) \right] d\vec{\Omega}'. \quad (7.18)$$

The process $t_\epsilon \rightarrow 0$ applied to the transition probability \bar{W}_{t_ϵ} ceates a δ -function and the particle density distribution \bar{f}_{t_ϵ} results in \bar{f} .

$$\begin{aligned} \lim_{t_\epsilon \rightarrow 0} \bar{W}_{t_\epsilon} &= \delta(\vec{\Omega}, \vec{\Omega}') \quad \text{:delta-Function} \\ \lim_{t_\epsilon \rightarrow 0} \bar{f}_{t_\epsilon} &= \bar{f} \\ \lim_{t_\epsilon \rightarrow 0} \vec{v}_{t_\epsilon} &= \vec{v} = \vec{v} \cdot \vec{\Omega} \\ \lim_{t_\epsilon \rightarrow 0} \vec{\Omega}_{t_\epsilon} &= \vec{\Omega} \end{aligned} \quad (7.19)$$

These relations applied to equation (7.18) give

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \bar{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' - \bar{f}_{t_\epsilon}}{\epsilon \cdot \tau_E} = \frac{\partial \bar{f}}{\partial t} + \vec{v} \cdot \vec{\nabla} \bar{f} \quad (7.20)$$

and

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \bar{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' - \bar{f}_{t_\epsilon}}{\epsilon \cdot \tau_E} \quad (7.21)$$

subsequently called exchange-term.

7.3. Calculation of the exchange-term

The dependencies of the transition probability \bar{W}_{t_ϵ} on the initially uncorrelated movement directions $\vec{\Omega}$ and $\vec{\Omega}'$ may be expressed by the scalar product of the movement directions $\vec{\Omega} \cdot \vec{\Omega}'$ and the simultaneous interchange of the constant time scaling τ_E by a time scaling depending on location, time and direction $t_E(\vec{x}, \vec{\Omega}, t)$, i.e.

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \bar{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' - \bar{f}_{t_\epsilon}}{\epsilon \cdot \tau_E} &= \lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') \bar{f}'_{t_\epsilon} d\vec{\Omega}' - \bar{f}_{t_\epsilon}}{\epsilon \cdot t_E(\vec{x}, \vec{\Omega}, t)} \\ \bar{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}') &\longrightarrow \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') \\ \tau_E &\longrightarrow t_E(\vec{x}, \vec{\Omega}, t). \end{aligned} \quad (7.22)$$

The direction distribution of the particles is developed by complex spherical harmonics P_{lm} , the transition probability by legendre polynomials P_l .

$$\bar{f}(\vec{x}, \vec{\Omega}, t) = \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \bar{f}_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}) = \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \bar{f}_{lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \quad (7.23)$$

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega}' \cdot \vec{\Omega}) = \sum_{l=0}^{+\infty} \widetilde{W}_{t_\epsilon l} P_l(\cos(\alpha)) = \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} \widetilde{W}_{t_\epsilon l} P_{lm}(\vec{\Omega}') P_{lm}^*(\vec{\Omega}) \quad (7.24)$$

The spherical harmonics P_{lm} are

$$P_{lm}(\vec{\Omega}) = e^{im\varphi} \frac{(-\sin(\vartheta))^m}{l!2^l} \cdot \left(\frac{(l-m)!}{(l+m)!} \right)^{\frac{1}{2}} \frac{d^{l+m}(\cos^2\vartheta - 1)^l}{(d\cos\vartheta)^{l+m}} \quad (7.25)$$

The normalisation holds:

$$\int_{4\pi} P_{lm} P_{lm}^* d\vec{\Omega} = \begin{cases} \frac{4\pi}{2l+1} & l = l' \text{ und } m=m' \\ 0 & \text{sonst} \end{cases} \quad (7.26)$$

There is the relation between spherical harmonics P_{lm} and Legendre polynomials P_l :

$$P_l(\vec{\Omega}' \cdot \vec{\Omega}) = P_l(\cos(\alpha)) = \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}') P_{lm}^*(\vec{\Omega}) \quad (7.27)$$

Thus one has

$$\begin{aligned} \int_{4\pi} \widetilde{W}_{t_\epsilon} \bar{f}'_\epsilon d\vec{\Omega}' &= \int_{4\pi} \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} \widetilde{W}_{t_\epsilon l} P_{lm}(\vec{\Omega}') P_{lm}^*(\vec{\Omega}) \cdot \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \bar{f}_{t_\epsilon lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}') d\vec{\Omega}' \\ &= \sum_{l=0}^{+\infty} \widetilde{W}_{t_\epsilon l} \frac{4\pi}{2l+1} \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) \bar{f}_{t_\epsilon lm}(\vec{x}, t). \end{aligned} \quad (7.28)$$

The left side of equation (7.20) results in

$$\begin{aligned}
 \lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \widetilde{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' - \bar{f}_{t_\epsilon}}{t_E \cdot \epsilon} &= \lim_{\epsilon \rightarrow 0} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \frac{(\widetilde{W}_{t_\epsilon l \frac{4\pi}{2l+1}} - 1)}{t_E \cdot \epsilon} \bar{f}_{t_\epsilon lm}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) \\
 &= \frac{1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \Upsilon_l \bar{f}_{t_\epsilon lm}(\vec{x}, t) P_{lm}(\vec{\Omega})
 \end{aligned} \tag{7.29}$$

with

$$\Upsilon_l = \lim_{\epsilon \rightarrow 0} \frac{(\widetilde{W}_{t_\epsilon l \frac{4\pi}{2l+1}} - 1)}{\epsilon} \tag{7.30}$$

as exchange coefficient.

Now equation (7.20) yields

$$\boxed{\frac{1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \Upsilon_l \bar{f}_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}) = \frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \cdot \vec{\nabla} \bar{f}}. \tag{7.31}$$

7.4. Calculation of the exchange-coefficients Υ_l

The transition probability is outlined by Legendre-polynomials respectively spherical harmonics:

$$\begin{aligned}
 \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') &= \sum_{l=0}^{+\infty} \widetilde{W}_{t_\epsilon l} P_l(\cos(\vartheta)) = \sum_{l=0}^{+\infty} \widetilde{W}_{t_\epsilon l} \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}') \\
 \cos(\vartheta) &= \vec{\Omega} \cdot \vec{\Omega}' = \mu.
 \end{aligned} \tag{7.32}$$

On the other hand is

$$\begin{aligned} \lim_{t_\epsilon \rightarrow 0} \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') &= \delta(\vec{\Omega} \cdot \vec{\Omega}') \\ \delta(\vec{\Omega} \cdot \vec{\Omega}') &= \sum_{l=0}^{+\infty} \frac{2l+1}{4\pi} \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}') = \sum_{l=0}^{+\infty} \frac{2l+1}{4\pi} P_l \quad \text{see(9.20)}. \end{aligned} \quad (7.33)$$

$\widetilde{W}_{t_\epsilon}(\mu) \geq 0$ is only in the range $\mu \in [1 - \varepsilon, 1]$ essentially different from 0. So the Legendre polynomials are approximated by

$$\begin{aligned} P_l(\mu) &= 1 - \frac{dP_l}{d\mu}\bigg|_1 \cdot \varepsilon + O(\varepsilon^2) \quad \varepsilon = 1 - \mu \\ \frac{dP_l}{d\mu}\bigg|_1 &= \frac{l(l+1)}{2} \quad \text{see (9.1)} \quad P_0 = 1, P_1 = \mu \\ &\Rightarrow \\ P_l(\mu) &= P_0 - (P_0 - P_1) \frac{l(l+1)}{2} + O(\varepsilon^2). \end{aligned} \quad (7.34)$$

Using

$$\int_{-1}^{+1} P_l P_{l'} d\mu = \delta_{ll'} \frac{2}{2l+1} \quad (7.35)$$

follows

$$\int_{-1}^{+1} \widetilde{W}_{t_\epsilon} P_l d\mu = 2\widetilde{W}_{t_\epsilon 0} - l(l+1)\widetilde{W}_{t_\epsilon 0} + \frac{l(l+1)}{3}\widetilde{W}_{t_\epsilon 1} = \frac{2}{2l+1}\widetilde{W}_{t_\epsilon l}. \quad (7.36)$$

Furthermore is

$$\begin{aligned} \int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') d\vec{\Omega}' &= \int_{4\pi} \widetilde{W}_{t_\epsilon 0} d\vec{\Omega}' = 4\pi \widetilde{W}_{t_\epsilon 0} = 1 \\ &\Rightarrow \quad \widetilde{W}_{t_\epsilon 0} = \frac{1}{4\pi}, \end{aligned} \quad (7.37)$$

as $\widetilde{W}_{t_\epsilon}$ for $t_\epsilon \rightarrow 0$ degenerates to a δ -function. That is why the $\widetilde{W}_{t_\epsilon l}$ are expressed by $\widetilde{W}_{t_\epsilon 1}$ and the determination of $\widetilde{W}_{t_\epsilon 1}$ remains to be calculated. We set

$$\lim_{\epsilon \rightarrow 0} \frac{(\widetilde{W}_{t_{\epsilon}1}^{\frac{4\pi}{3}} - 1)}{\epsilon} = \zeta. \quad (7.38)$$

Multiplying equation (7.36) with 2π leads to

$$\frac{4\pi}{2l+1} \widetilde{W}_{t_{\epsilon}l} = 4\pi \widetilde{W}_{t_{\epsilon}0} - (4\pi) \frac{l(l+1)}{2} \widetilde{W}_{t_{\epsilon}0} + \frac{4\pi}{3} \frac{l(l+1)}{2} \widetilde{W}_{t_{\epsilon}1}. \quad (7.39)$$

I.e.

$$\begin{aligned} \frac{4\pi}{2l+1} \widetilde{W}_{t_{\epsilon}l} - \mathbf{1} &= \frac{l(l+1)}{2} \left(\frac{4\pi}{3} \widetilde{W}_{t_{\epsilon}1} - 1 \right) = -\frac{l(l+1)}{2} \zeta + O(\epsilon^2) = \Upsilon_l + O(\epsilon^2) \\ \Upsilon_l &= -\frac{l(l+1)}{2} \zeta \end{aligned} \quad (7.40)$$

\Rightarrow

$$\boxed{\frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \cdot \vec{\nabla} \bar{f} = -\frac{1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \frac{l(l+1)}{2} \bar{f}_{l,m}(\vec{x}, t) P_{l,m}(\vec{\Omega})} \quad (7.41)$$

This equation only contains the unknown coefficients t_E and \bar{v} principally depending upon the space-time-point (\vec{x}, t) and the fluctuation direction $\vec{\Omega}$

$$\begin{aligned} t_E &= t_E(\vec{x}, t, \vec{\Omega}) \\ \bar{v} &= \bar{v}(\vec{x}, t, \vec{\Omega}). \end{aligned} \quad (7.42)$$

The total derivation of $\bar{f}(\vec{x}, t, \vec{\Omega})$ with respect to t in direction of $\vec{\Omega}$ leads to

$$\frac{d}{dt} \bar{f}(\vec{x}, t, \vec{\Omega}) = \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \frac{d}{dt} \bar{f}_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}) = \frac{1}{t_E} \sum_{l=1}^{+\infty} \gamma_l \sum_{m=-l}^{+l} \bar{f}_{lm}(\vec{x}, t) P_{lm}(\vec{\Omega}). \quad (7.43)$$

The time behavior of the spherical harmonic components is described by the equations

$$\frac{d}{dt} \bar{f}_{lm}(t) = \frac{\gamma_l}{t_E} \bar{f}_{lm} \quad (7.44)$$

and result in


$$\bar{f}_{lm}(t) \sim \exp\left(\frac{\gamma_l}{t_E} \cdot t\right). \quad (7.45)$$

The greater the order l the more powerful is its temporal decay.

7.5. Reconstruction of the transition probabilities

\bar{W}_{t_ϵ}

The Transition probability $\widetilde{W}_{t_\epsilon 0 \rightarrow 1 \rightarrow 2}$, changing the movement direction $\vec{\Omega}$ at the times t_0, t_1, t_2 from $\vec{\Omega}_0$ via $\vec{\Omega}_1$ to $\vec{\Omega}_2$



is the product of the single transition probabilities.

$$\widetilde{W}_{t_\epsilon, 0 \rightarrow 1 \rightarrow 2} = \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1) \cdot \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2). \quad (7.46)$$

On the other side

$$\widetilde{W}_{t_\epsilon l} = (1 + \Upsilon_l \epsilon) \frac{2l+1}{4\pi} + O(\epsilon^2) \quad (7.47)$$

holds and thus arises

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') = \sum_{l=0}^{+\infty} (1 + \Upsilon_l \epsilon) \frac{2l+1}{4\pi} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') + O(\epsilon^2). \quad (7.48)$$

The probability, that a particle changes the direction after an infinitesimal time interval $\epsilon \cdot t_E$ from $\vec{\Omega}_0$ to $\vec{\Omega}_2$ is given by

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega}_0 \cdot \vec{\Omega}_2) = \int_{4\pi} \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1) \cdot \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2) d\vec{\Omega}_1 \quad (7.49)$$

and

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega}_0 \cdot \vec{\Omega}_2) = \sum_{l=0}^{+\infty} \left(1 + \Upsilon_l \frac{\epsilon}{2}\right)^2 \frac{2l+1}{4\pi} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}_0) P_{lm}(\vec{\Omega}_2) + O(\epsilon^2). \quad (7.50)$$

Using n intermediate steps $\widetilde{W}_{t_\epsilon}$ is expressed by an integral over the product of the single transition probabilities.

$$\widetilde{W}_{t_\epsilon, 0 \rightarrow 1 \dots \rightarrow n} = \widetilde{W}_{\frac{t_\epsilon}{n}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1) \cdot \widetilde{W}_{\frac{t_\epsilon}{n}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2) \dots \widetilde{W}_{\frac{t_\epsilon}{n}}(\vec{\Omega}_{n-1} \cdot \vec{\Omega}_n) \quad (7.51)$$

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega}_0 \cdot \vec{\Omega}_n) = \int_{4\pi} \int_{4\pi} \dots \int_{4\pi} \widetilde{W}_{\frac{t_\epsilon}{n}} \cdot \widetilde{W}_{\frac{t_\epsilon}{n}} \dots \widetilde{W}_{\frac{t_\epsilon}{n}} d\vec{\Omega}_1 \dots d\vec{\Omega}_{n-1} \quad (7.52)$$

For $n \rightarrow \infty$ this results in:

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') = \lim_{n \rightarrow \infty} \sum_{l=0}^{+\infty} \left\{ 1 + \frac{\epsilon \Upsilon_l}{n} \right\}^n \frac{2l+1}{4\pi} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') + O(\epsilon^2) \quad (7.53)$$

and finally

$$\boxed{\widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') = \sum_{l=0}^{+\infty} e^{\Upsilon_l \cdot \epsilon} \frac{2l+1}{4\pi} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') + O(\epsilon^2)} \quad (7.54)$$

Selecting $\epsilon = \frac{t_\epsilon}{t_E(\vec{x}, t, \vec{\Omega})}$ the exchange function $\widetilde{W}_{t_\epsilon}$ may be understood in the dependencies

$$\widetilde{W}_{t_\epsilon} = \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}'). \quad (7.55)$$

Therefore

$$\overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}') \approx \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}') \quad (7.56)$$

is calculated, too. \implies

$$\boxed{\begin{aligned} \overline{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, t) &= \int_{4\pi} \overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Omega}') \overline{f}_{t_\epsilon}(\vec{x} - t_\epsilon \cdot \vec{v}' \vec{\Omega}', \vec{\Omega}', t - t_\epsilon) d\vec{\Omega}' \\ \vec{v}' &= \vec{v}'(\vec{x}, \vec{\Omega}', t) \end{aligned}} \quad (7.57)$$

The Transition probability $\overline{W}_{t_\epsilon}$ is unsymmetrical in the direction quantities on account of $\epsilon = \frac{t_\epsilon}{t_E(\vec{x}, t, \vec{\Omega})}$.

7.6. Approximation formula

In 1st approximation a telegrapher's equations is derived out of the linear Boltzmann Equation leading to the known diffusion equation without taking into account the second time derivation. In this case the diffusion equation is proved to be usefull. Subsequent considerations are displaying which relation exists between the 1st approximation of the particle transport by longitudinal continuum fluctuations and the known diffusion equation.

Assuming the simplification

$$\frac{1}{t_E} = \tau(\vec{\mathbf{x}}, \vec{\Omega}, t) = \tau_0 = \text{const} \quad (7.58)$$

the transport equation described in 1st approximation is

$$\frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \cdot \nabla \bar{f} = -\frac{1}{t_E} \cdot \sum_{m=-1}^{+1} \bar{f}_{1m}(\vec{\mathbf{x}}, t) P_{1m}(\vec{\Omega}). \quad (7.59)$$

In cartesian coordinates one gets

$$\frac{\partial \bar{f}}{\partial t} + \bar{v}_x \Omega_x \cdot \frac{\partial \bar{f}}{\partial x} + \bar{v}_y \Omega_y \cdot \frac{\partial \bar{f}}{\partial y} + \bar{v}_z \Omega_z \cdot \frac{\partial \bar{f}}{\partial z} = -\frac{1}{t_E} \cdot \sum_{m=-1}^{+1} \bar{f}_{1m}(\vec{\mathbf{x}}, t) P_{1m}(\vec{\Omega}) \quad (7.60)$$

with

$$\bar{v}_x = \bar{v}(\vec{\mathbf{x}}, t, \vec{\Omega}) \quad \bar{v}_y = \bar{v}(\vec{\mathbf{x}}, t, \vec{\Omega}) \quad \bar{v}_z = \bar{v}(\vec{\mathbf{x}}, t, \vec{\Omega}). \quad (7.61)$$

Subsequently we confine us on

$$\bar{v}_x = \bar{v}_y = \bar{v}_z = \bar{v}(\vec{\mathbf{x}}), \quad (7.62)$$

suggesting an isotropy of fluctuation motions in the statistical ensemble. The conditions are selected such that the further derivations analogous to 6.4 follow until to a telegrapher's equation.

$$t_E \frac{\partial^2 \Phi}{\partial t^2} + \frac{\partial \Phi}{\partial t} = \mathbf{t}_E \bar{v} \vec{\nabla} \cdot \frac{\bar{v}}{3} \vec{\nabla} \Phi \quad (7.63)$$

The usual diffusion coefficient normally contained in $\vec{\nabla} \cdot \mathbf{D} \vec{\nabla} \Phi$ cannot be found. In the equation above $\mathbf{D} = \mathbf{t}_E \cdot \frac{\bar{v}^2}{3}$ is contained partly outside partly between the $\vec{\nabla}$ -operators. This has consequences in inhomogeneous media. Such problems arise unrecognized using the Bousinesque approach . I.e. in an inhomogeneous medium

this approach may be fatal. The term of second derivation by time has nothing to do with relativistic theory. Because of the small size of t_E it may generally be neglected.

8. Stochastic transport by turbulent continuum-fluctuations

$$\begin{aligned} \frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \times \vec{\Theta} \cdot \nabla \bar{f} &= \frac{-1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \frac{l(l+1)}{2} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}) \\ &\Downarrow \\ \bar{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, \vec{\Theta}, t) &= \int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}, \vec{\Omega}', \vec{\Theta}') \bar{f}_{t_\epsilon}(\vec{x} - t_\epsilon \cdot \vec{v}' \vec{\Omega}' \times \vec{\Theta}', \vec{\Omega}', \vec{\Theta}'), t - t_\epsilon) d\vec{\Omega}' d\vec{\Theta}' \end{aligned}$$

8.1. Introduction

The motion of passive particles by turbulent continuum fluctuations is examined. The particles are moved not affecting this field. Their trajectories correspond in every ε -neighborhood of a point to a circle segment passed with the velocity

$$\vec{v}_{t_\epsilon} = \vec{\omega}_{t_\epsilon} \times \vec{r}_{t_\epsilon}. \quad (8.1)$$

The considered motion quantities $\vec{\omega}_{t_\epsilon}$ and \vec{r}_{t_ϵ} are determined by successively detecting a single particle originating from a point \vec{x}_0 after a time t_ϵ moving to \vec{x}_1 and after a further time t_ϵ to \vec{x}_2 . By these 3 points a circle segment is uniquely defined for the point \vec{x}_1 with radius vector \vec{r}_{t_ϵ} and a rotation speed $\vec{\omega}_{t_\epsilon}$.

$$\begin{aligned} \vec{r}_{t_\epsilon} &= \vec{r}_{t_\epsilon} \cdot \vec{\Theta}_{t_\epsilon} \\ \vec{\omega}_{t_\epsilon} &= \vec{\omega}_{t_\epsilon} \cdot \vec{\Omega}_{t_\epsilon} \end{aligned} \quad (8.2)$$

In the special case $\vec{\omega}_{t_\epsilon} \rightarrow \mathbf{0}$ and $\vec{r} \rightarrow +\infty$ the velocity \vec{v}_{t_ϵ} is revealed out of its neighborhood.¹ The particle density distributions are received in a thought experiment by an unlimited number of deterministic ensemble-systems (see chapter 4). In every

¹Applying the deterministic theory this problem must be treated numerically.

point (\vec{x}, t) a continuously differentiable particle density distribution of the motion quantities $\vec{\omega}_{t_\epsilon}$ und \vec{r}_{t_ϵ} is assigned in accordance with

$$f_{t_\epsilon} = f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}). \quad (8.3)$$

The with t_ϵ indexed functions are automatically assumed to contain motion quantities of corresponding measurement accuracies. The indexing of the motion quantities can be omitted if the functions are indexed. After execution of a limiting process for example

$$\lim_{t_\epsilon \rightarrow 0} f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}) = f(\vec{x}, t, \vec{\omega}, \vec{r}) \quad (8.4)$$

f and $(\vec{\omega}, \vec{r})$ are understood according to an exact measuring process. Integrating the particle density distribution over the motion quantities one obtains expectation values of a particle density not conforming with the actual particle density ρ .

$$\langle \rho_{t_\epsilon}(\vec{x}, t) \rangle = \int_{2\pi} \int_{4\pi} \int_0^\infty \int_0^\infty f_{t_\epsilon}(\vec{x}, t, \omega \cdot \vec{\Omega}, r \cdot \vec{\Theta}) d\omega dr d\vec{\Omega} d\vec{\Theta} \neq \rho_{t_\epsilon}(\vec{x}, t) \quad (8.5)$$

A strictly deduced partial differential equation is obtained calculating the development of spatio-temporal particle density distributions. The incipiently unlimited number of unknown coefficients is reduced to a local time-scaling related to the vortex calculation of an associated deterministic theory discussed in further chapters. The initially abstractly formulated transition probabilities get concrete functional dependencies. There are always found suitable Monte-Carlo methods treating them with the help of the deterministic theory described in further chapters.

8.2. The transport as Markov Process with natural causality

A particle at location \vec{x} and time t changing its velocity from $\vec{v}' = (\vec{\omega}' \times \vec{r}')$ to $\vec{v} = (\vec{\omega} \times \vec{r})$ is given by the transition probability

$$W_{t_\epsilon} = W_{t_\epsilon}(\vec{x}, t; \vec{\omega}, \vec{r}; \vec{\omega}', \vec{r}') \quad (8.6)$$

with

$$\int_0^\infty \int_0^\infty \int_{4\pi} \int_{2\pi} W_{t_\epsilon}(\vec{x}, t; \vec{\omega}, \vec{r}; \vec{\omega}', \vec{r}') d\omega' dr' d\vec{\Omega}' d\vec{\Theta}' = 1. \quad (8.7)$$

\Rightarrow

$$f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}) = \int_0^\infty \int_0^\infty \int_{4\pi} \int_{2\pi} W_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}, \vec{\omega}', \vec{r}') f_{t_\epsilon}(\vec{x} - \vec{\omega}' \times \vec{r}' \cdot t_\epsilon, \vec{\omega}', \vec{r}', t - t_\epsilon) d\omega' dr' d\Omega' d\Theta' \quad (8.8)$$

Continuity is required respectively of all variables of the transition probability W_{t_ϵ} . The sequence of velocities $\vec{v}'_{t_\epsilon}, \vec{v}_{t_\epsilon}$ means a motion from

$$(\vec{x} - \vec{\omega}'_{t_\epsilon} \times \vec{r}'_{t_\epsilon} \cdot t_\epsilon, t - t_\epsilon, \vec{\omega}'_{t_\epsilon} \times \vec{r}'_{t_\epsilon}) \quad \text{to} \quad (\vec{x}, t, \vec{\omega}_{t_\epsilon} \times \vec{r}_{t_\epsilon}). \quad (8.9)$$

For the limiting process $t_\epsilon \rightarrow 0$ the transition probabilities W_{t_ϵ} prove to be physical realizations of test functions of the distribution theory.

$$\lim_{t_\epsilon \rightarrow 0} W_{t_\epsilon} = \delta(\vec{\omega}, \vec{r}; \vec{\omega}', \vec{r}'). \quad (8.10)$$

The passive scalar particles precisely reproduce the motions of the fluctuation field. For the particle density distribution $f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r})$ the following separation approach is used without loss of generality:

$$f_{t_\epsilon}(\vec{x} - \vec{\omega} \times \vec{r} \cdot t_\epsilon, t, \vec{\omega}, \vec{r}) = G_{t_\epsilon}(\vec{x} - \vec{\omega} \times \vec{r} \cdot t_\epsilon, t, \vec{\omega}, \vec{r}) \bar{f}_{t_\epsilon}(\vec{x} - \vec{v} \vec{\Omega} \times \vec{\Theta} \cdot t_\epsilon, t, \vec{\Omega}, \vec{\Theta}) \quad (8.11)$$

with

$$\begin{aligned} \int_0^\infty \int_0^\infty G_{t_\epsilon}(\vec{x}, t, \omega \vec{\Omega}, r \vec{\Theta}) d\omega dr &= 1 \\ \int_0^\infty \int_0^\infty G_{t_\epsilon}(\vec{x}, t, \omega \vec{\Omega}, r \vec{\Theta}) \omega r d\omega dr &= \bar{v}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \\ \bar{v}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \bar{\omega}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \cdot \bar{r}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \end{aligned} \quad (8.12)$$

\Rightarrow

$$\bar{f}_{t_\epsilon}(\vec{x} - \vec{v} \vec{\Omega} \times \vec{\Theta} \cdot t_\epsilon, t, \vec{\Omega}, \vec{\Theta}) = \int_0^\infty \int_0^\infty f_{t_\epsilon}(\vec{x} - \vec{\omega} \times \vec{r} \cdot t_\epsilon, t, \omega \cdot \vec{\Omega}, r \cdot \vec{\Theta}) d\omega dr \quad (8.13)$$

One obtains a transition probability \bar{W}_{t_ϵ} only depending on the directions by inte-

grating W_{t_ϵ} over the amounts ω', r', ω, r .

$$\boxed{\overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}, \vec{\Omega}', \vec{\Theta}') = \int_0^\infty \int_0^\infty \int_0^\infty \int_0^\infty W_{t_\epsilon} G_{t_\epsilon}(\vec{x} - \vec{\omega}' \times \vec{r}' \cdot t_\epsilon, t - t_\epsilon, \vec{\omega}', \vec{r}') d\omega' dr' d\omega dr} \quad (8.14)$$

The integration

$$\int_0^\infty \int_0^\infty (8.8) d\omega dr \quad (8.15)$$

gives

$$\bar{f}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) = \int_0^\infty \int_0^\infty \int_0^\infty \int_0^\infty \int_{4\pi} \int_{2\pi} W_{t_\epsilon} f_{t_\epsilon}(\vec{x} - \vec{\omega}' \times \vec{r}' \cdot t_\epsilon, t - t_\epsilon, \vec{\omega}', \vec{r}') d\omega' dr' d\omega dr d\vec{\Omega}' d\vec{\Theta}' \quad (8.16)$$

$$\Rightarrow \boxed{\bar{f}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) = \int_{4\pi} \int_{2\pi} \overline{W}_{t_\epsilon} \bar{f}_{t_\epsilon}(\vec{x} - \vec{v}' \vec{\Omega}' \times \vec{\Theta}' \cdot t_\epsilon, t - t_\epsilon, \vec{\Omega}', \vec{\Theta}') d\vec{\Omega}' d\vec{\Theta}'} \quad (8.17)$$

In the integrand \bar{f}_{t_ϵ} is developed around \vec{x} and t :

$$\bar{f}_{t_\epsilon}(\vec{x} - \vec{\Delta} \vec{x}, t - t_\epsilon, \vec{\Omega}', \vec{\Theta}') = \bar{f}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}', \vec{\Theta}') - \tau_E \cdot \epsilon \cdot \left[\frac{\partial \bar{f}_{t_\epsilon}}{\partial t} + \vec{v}' \vec{\Omega}' \times \vec{\Theta}' \cdot \nabla \bar{f}_{t_\epsilon} + O(\epsilon^2) \right] \quad (8.18)$$

This leads to

$$\begin{aligned} & \frac{\int_{4\pi} \int_{2\pi} \overline{W}_{t_\epsilon} \bar{f}_{t_\epsilon}' d\vec{\Omega}' d\vec{\Theta}' - \bar{f}_{t_\epsilon}}{\epsilon} = \\ & \int_{4\pi} \int_{2\pi} \overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}', \vec{\Theta}', \vec{\Omega}, \vec{\Theta}) \cdot \tau_E \left[\frac{\partial \bar{f}_{t_\epsilon}}{\partial t} + \vec{v}' \vec{\Omega}' \times \vec{\Theta}' \cdot t_\epsilon \cdot \nabla \bar{f}_{t_\epsilon} + O(\epsilon^2) \right] d\vec{\Omega}' d\vec{\Theta}'. \end{aligned} \quad (8.19)$$

As

$$\lim_{t_\epsilon \rightarrow 0} \overline{W}_{t_\epsilon} = \delta(\vec{\Omega}, \vec{\Theta}; \vec{\Omega}', \vec{\Theta}') \quad (8.20)$$

\Rightarrow

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \int_{2\pi} \overline{W}_{t_\epsilon} \bar{f}_{t_\epsilon}' d\vec{\Omega}' d\vec{\Theta}' - \bar{f}_{t_\epsilon}}{\epsilon \cdot \tau_E} = \frac{\partial \bar{f}}{\partial t} + \vec{v} \vec{\Omega} \times \vec{\Theta} \cdot \nabla \bar{f}. \quad (8.21)$$

Furtheron

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \int_{2\pi} \overline{W}_{t_\epsilon} \overline{f}'_{t_\epsilon} d\vec{\Omega}' d\vec{\Theta}' - \overline{f}_{t_\epsilon}}{\epsilon \cdot \tau_E} \quad (8.22)$$

is called exchange-term.

8.3. Calculation of the exchange-term

Exchange term dependencies of scalar products $\vec{\Omega} \cdot \vec{\Omega}'$ and $\vec{\Theta} \cdot \vec{\Theta}'$ are taken into account instead of individually depending directions $\vec{\Omega}$, $\vec{\Omega}'$ and $\vec{\Theta}$, $\vec{\Theta}'$ demanding the following relation

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{2\pi} \int_{4\pi} \overline{W}_{t_\epsilon} \overline{f}'_{t_\epsilon} d\vec{\Omega}' d\vec{\Theta}' - \overline{f}_{t_\epsilon}}{\epsilon \cdot \tau_E} = \lim_{\epsilon \rightarrow 0} \frac{\int_{2\pi} \int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') \overline{f}'_{t_\epsilon} d\vec{\Omega}' d\vec{\Theta}' - \overline{f}_{t_\epsilon}}{\epsilon \cdot t_E}. \quad (8.23)$$

The following transitions

$$\begin{aligned} \tau_E = \text{const} & \longrightarrow t_E = t_E(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) \\ \overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}; \vec{\Omega}', \vec{\Theta}') & \longrightarrow \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') \end{aligned} \quad (8.24)$$

are regarded. Moreover, a separation of $\vec{\Omega} \cdot \vec{\Omega}'$ and $\vec{\Theta} \cdot \vec{\Theta}'$ is assumed:

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') = V_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}') \cdot M_{t_\epsilon}(\vec{\Theta} \cdot \vec{\Theta}'). \quad (8.25)$$

Functions of the unit vectors $\vec{\Omega}$ and $\vec{\Theta}$ are presented by a complete orthogonal function system representing an extension of the spherical harmonics called turbulence functions.

$$\begin{aligned} \overline{f}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \overline{f}_{t_\epsilon lmk}(\vec{x}, t) Q_{lmk}(\vec{\Omega}, \vec{\Theta}) \\ &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \overline{f}_{t_\epsilon lmk}(\vec{x}, t) Q_{lmk}^*(\vec{\Omega}, \vec{\Theta}) \end{aligned} \quad (8.26)$$

$$\int_{2\pi} \int_{4\pi} Q_{lmk}(\vec{\Omega}, \vec{\Theta}) Q_{lmk}^*(\vec{\Omega}', \vec{\Theta}') d\vec{\Omega}' d\vec{\Theta}' = \begin{cases} \frac{8\pi^2}{2l+1} & \text{für } l = l' \text{ und } m=m' \\ 0 & \text{sonst} \end{cases} \quad (8.27)$$

with

$$\begin{aligned}
 Q_{lmk}(\vec{\Omega}, \vec{\Theta}) &= P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}) \\
 \int_{2\pi} H_{k'}(\vec{\Theta}) H_k^*(\vec{\Theta}) d\vec{\Theta} &= \begin{cases} 2\pi & \text{für } k'=k \\ 0 & \text{sonst} \end{cases} \\
 H_k(\vec{\Theta}) &= e^{ik\theta}
 \end{aligned} \tag{8.28}$$

The product $\vec{\Omega} \cdot \vec{\Omega}'$ in the separated exchange function V_{t_ϵ} is developed by spherical harmonics.

$$\begin{aligned}
 V_{t_\epsilon}(\vec{\Omega}' \cdot \vec{\Omega}) &= \sum_{l=0}^{+\infty} V_{t_\epsilon l} P_l(\cos(\alpha)) = \sum_{l=0}^{+\infty} V_{t_\epsilon l} \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}') P_{lm}^*(\vec{\Omega}) \\
 &\quad \text{mit} \\
 \lim_{t_\epsilon \rightarrow 0} V_{t_\epsilon}(\vec{\Omega}' \cdot \vec{\Omega}) &= \delta(\vec{\Omega}, \vec{\Omega}')
 \end{aligned} \tag{8.29}$$

The product $\vec{\Theta} \cdot \vec{\Theta}'$ in the separated exchange function M_{t_ϵ} is developed by functions H_k .

$$M_{t_\epsilon}(\vec{\Theta}' \cdot \vec{\Theta}) = \sum_{k=0}^{+\infty} M_{t_\epsilon k} \cos(k\beta) = \frac{1}{2} \sum_{k=0}^{k=+\infty} M_{t_\epsilon k} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] \tag{8.30}$$

with

$$\begin{aligned}
 \cos(k\beta) &= \frac{1}{2} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] = \frac{1}{2} [e^{ik(\theta' - \theta)} + e^{-ik(\theta' - \theta)}] \\
 \vec{\Theta}' \cdot \vec{\Theta} = \cos(\beta) &= \cos(\theta' - \theta) = \frac{1}{2} [H_1(\vec{\Theta}') H_1^*(\vec{\Theta}) + H_{-1}(\vec{\Theta}') H_{-1}^*(\vec{\Theta})] = \frac{1}{2} [e^{i(\theta' - \theta)} + e^{-i(\theta' - \theta)}] \\
 \lim_{t_\epsilon \rightarrow 0} M_{t_\epsilon}(\vec{\Theta}' \cdot \vec{\Theta}) &= \delta(\vec{\Theta}, \vec{\Theta}')
 \end{aligned} \tag{8.31}$$

\Rightarrow

$$\begin{aligned}
 \int_{4\pi} \int_{2\pi} \widetilde{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' d\vec{\Theta} &= \int_{4\pi} \int_{2\pi} V_{t_\epsilon} (\vec{\Omega}' \cdot \vec{\Omega}) \cdot M_{t_\epsilon} (\vec{\Theta}' \cdot \vec{\Theta}) \bar{f}'_{t_\epsilon} d\vec{\Omega}' d\vec{\Theta}' \\
 &= \int_{4\pi} \int_{2\pi} \left[\sum_{l=0}^{+\infty} V_{t_\epsilon l} \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}') P_{lm}^*(\vec{\Omega}) \cdot \frac{1}{2} \sum_{k=0}^{k=+\infty} M_{t_\epsilon k} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] \right] \\
 &\quad \cdot \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \bar{f}_{t_\epsilon lmk}(\vec{x}, t) P_{lm}^*(\vec{\Omega}') H_k^*(\vec{\Theta}') \right] d\vec{\Omega}' d\vec{\Theta}' \\
 &= \sum_{l=0}^{+\infty} V_{t_\epsilon l} \frac{4\pi}{2l+1} \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) \sum_{k=0}^{+\infty} M_{t_\epsilon k} 2\pi \bar{f}_{t_\epsilon lmk}(\vec{x}, t) H_k^*(\vec{\Theta}).
 \end{aligned} \tag{8.32}$$

Finally the exchange term results in

$$\begin{aligned}
 \lim_{\epsilon \rightarrow 0} \frac{\int_{4\pi} \int_{2\pi} \widetilde{W}_{t_\epsilon} \bar{f}'_{t_\epsilon} d\vec{\Omega}' d\vec{\Theta}' - \bar{f}_{t_\epsilon}}{\epsilon} \\
 &= \lim_{\epsilon \rightarrow 0} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=0}^{+\infty} \frac{(V_{t_\epsilon l} \frac{4\pi}{2l+1} M_{t_\epsilon k} 2\pi - 1)}{\epsilon} \bar{f}_{t_\epsilon lmk}(\vec{x}, t) P_{lm}^*(\vec{\Omega}) H_k^*(\vec{\Theta}) \\
 &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=0}^{+\infty} \Upsilon_{lk} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}).
 \end{aligned} \tag{8.33}$$

With the exchange coefficients

$$\Upsilon_{lk} = \lim_{\epsilon \rightarrow 0} \frac{(V_{t_\epsilon l} \frac{4\pi}{2l+1} M_{t_\epsilon k} 2\pi - 1)}{\epsilon} \tag{8.34}$$

the transport equation

$$\boxed{\frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \times \vec{\Theta} \cdot \nabla \bar{f} = \frac{1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \Upsilon_{lk} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta})} \tag{8.35}$$

is achieved. Further on it is shown that in Υ_{lk} the index k may be skipped.

8.4. Calculation of the exchange-coefficients Υ_l

Considering an overall closed volume range V the particle number in the entire volume remains constant if no absorption is assumed.

$$\text{total number of particles} = \int_{\mathbb{V}} \int_{4\pi} \int_{2\pi} \bar{f} d\vec{\Omega} d\vec{\Theta} d\mathbb{V} = \text{const.} \quad (8.36)$$

\Rightarrow

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{V}} \int_{4\pi} \int_{2\pi} \bar{f} d\vec{\Omega} d\vec{\Theta} d\mathbb{V} = \\ \int_{\mathbb{V}} \int_{4\pi} \int_{2\pi} \left[\frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \times \vec{\Theta} \cdot \nabla \bar{f} \right] d\vec{\Omega} d\vec{\Theta} d\mathbb{V} = \Upsilon_{0,0} \cdot \mathbf{V} = 0 \end{aligned} \quad (8.37)$$

and thus

$$\boxed{\Upsilon_{0,0} = 0}. \quad (8.38)$$

Getting an overview over the exchange function M_{t_ϵ} the essential relations are presented again with the following equations:

$$\begin{aligned} M_{t_\epsilon}(\vec{\Theta}' \cdot \vec{\Theta}) &= \sum_{k=0}^{+\infty} M_{t_\epsilon k} \cos(k\beta) = \frac{1}{2} \sum_{k=0}^{k=+\infty} M_{t_\epsilon k} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] \\ \cos(k\beta) &= \frac{1}{2} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] = \frac{1}{2} [e^{ik(\theta' - \theta)} + e^{-ik(\theta' - \theta)}] \\ \vec{\Theta}' \cdot \vec{\Theta} = \cos(\beta) = \cos(\theta' - \theta) &= \frac{1}{2} [H_1(\vec{\Theta}') H_1^*(\vec{\Theta}) + H_{-1}(\vec{\Theta}') H_{-1}^*(\vec{\Theta})] = \frac{1}{2} [e^{i(\theta' - \theta)} + e^{-i(\theta' - \theta)}] \\ \lim_{\epsilon \rightarrow 0} M_{t_\epsilon}(\vec{x}, t, \vec{\Theta}' \cdot \vec{\Theta}) &= \delta(\vec{\Theta}, \vec{\Theta}') \\ \int_{2\pi} H_{k'}(\vec{\Theta}) H_k^*(\vec{\Theta}) d\vec{\Theta} &= \begin{cases} 2\pi & \text{für } k'=k \\ 0 & \text{else} \end{cases} \end{aligned}$$

$M_{t_\epsilon}(\vec{\Theta}' \cdot \vec{\Theta}) = \sum_{k=0}^{+\infty} M_{t_\epsilon k} \cos(k\beta)$ only takes values essentially different from 0 in an ε -neighborhood of $\beta = 0$, such that $\vec{\Theta}' \cdot \vec{\Theta} = \cos(\beta) = 1 - O(\varepsilon^2)$ is sufficient. \Rightarrow

$$\begin{aligned} 2\pi \cdot M_{t_\epsilon k} &= \\ \int_{-\pi}^{+\pi} M_{t_\epsilon} \cos(k\beta) d\beta &= \int_{-\pi}^{+\pi} M_{t_\epsilon} (1 - O(\epsilon)) d\beta = 2\pi \cdot M_{t_\epsilon 0} - O(\epsilon^2). \end{aligned} \quad (8.39)$$

On the other hand

$$\begin{aligned} \int_{2\pi} M_{t_\epsilon}(\vec{\Theta} \cdot \vec{\Theta}')' d\vec{\Theta}' = \\ \frac{1}{2} \int_{2\pi} \sum_{k=0}^{k=+\infty} M_{t_\epsilon k} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] d\vec{\Theta}' = 2\pi \cdot M_{t_\epsilon 0} = 1. \end{aligned} \quad (8.40)$$

is valid. \implies

$$M_{t_\epsilon k} = M_{t_\epsilon 0} = \frac{1}{2\pi}. \quad (8.41)$$

The calculation of the exchange coefficients is not influenced by M_{t_ϵ} the Υ -values given by

$$\Upsilon_l = \lim_{t_\epsilon \rightarrow 0} \frac{(V_{t_\epsilon l}^{\frac{4\pi}{2l+1}} - 1)}{t_\epsilon}. \quad (8.42)$$

Further calculation of the Υ_l analogously happen to section 7.4 with the result

$$\Upsilon_l = -\frac{l(l+1)}{2}\zeta \quad \zeta = \text{const.} \quad (8.43)$$

Now the equation of turbulent particle transport is written

$$\boxed{\frac{\partial \bar{f}}{\partial t} + \bar{v} \vec{\Omega} \times \vec{\Theta} \cdot \nabla \bar{f} = -\frac{1}{t_E} \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \frac{l(l+1)}{2} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta})} \quad (8.44)$$

the coefficient $\frac{\zeta}{t_E}$ replaced by $\frac{1}{t_E}$. A more complicated dependency of $t_E = t_E(\vec{x}, t, \vec{\Omega})$ possibly remains. Maybe, physically justified simplifications lead to practical solutions. The below presented theory of deterministic turbulence enables the calculation of these coefficients by numerical evaluation.

Die total derivative with respect to time gives

$$\begin{aligned}
 \frac{d}{dt} \bar{f}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{m=+l} \sum_{k=-\infty}^{+\infty} \frac{d}{dt} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}) \\
 &= \frac{1}{t_E} \sum_{l=1}^{+\infty} \gamma_l \cdot \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} \bar{f}_{lmk}(\vec{x}, t) P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}).
 \end{aligned} \tag{8.45}$$

The time behavior of the single modes are obtained by

$$\frac{d}{dt} \bar{f}_{lmk}(t) = \frac{\gamma_l}{t_E} \bar{f}_{lmk} \tag{8.46}$$


$$\bar{f}_{lmk}(t) \sim \exp\left(\frac{\gamma_l}{t_E} \cdot t\right). \tag{8.47}$$

The greater the order l the more powerful is its temporal decay.

8.5. Reconstruction of the transition probabilities

$\overline{W}_{t_\epsilon}$

The transition probability $\widetilde{W}_{t_\epsilon, 0 \rightarrow 1 \rightarrow 2}$, a particle changing its motion pair of directions $(\vec{\Omega}, \vec{\Theta})$ at the times t_0, t_1, t_2 from $(\vec{\Omega}_0, \vec{\Theta}_0)$ via $(\vec{\Omega}_1, \vec{\Theta}_1)$ to $(\vec{\Omega}_2, \vec{\Theta}_2)$,



results out of the product of the single probabilities of the pairs of directions (vortex vector and radius vector direction of motion in a circle segment). The grafical presentation is meant symbolically because such a pair of directions does not compose to an overall direction. $\vec{\Omega}_i$ is always orthogonal to $\vec{\Theta}_i$. A vectorial overall direction of $\vec{\Omega}_i$ and $\vec{\Theta}_i$ has no physical meaning in the 3 dimensional space. ²

$$\widetilde{W}_{t_\epsilon, 0 \rightarrow 1 \rightarrow 2} = \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1, \vec{\Theta}_0 \cdot \vec{\Theta}_1) \cdot \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2, \vec{\Theta}_1 \cdot \vec{\Theta}_2) \tag{8.48}$$

² $\vec{\Omega}, \vec{\Theta}$ would make a single direction vector in a 4-dimensional space. The longitudinal fluctuations in the 4-dimensional space should accord to turbulence in the 3-dimensional space.

The probability, that a particle changes its pair of directions within a time $t_\epsilon = \epsilon \cdot t_E$ from $(\vec{\Omega}_0, \vec{\Theta}_0)$ to $(\vec{\Omega}_2, \vec{\Theta}_2)$, is obtained by

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega}_0 \cdot \vec{\Omega}_2, \vec{\Theta}_0 \cdot \vec{\Theta}_2) = \int_{2\pi} \int_{4\pi} \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1, \vec{\Theta}_0 \cdot \vec{\Theta}_1) \cdot \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2, \vec{\Theta}_1 \cdot \vec{\Theta}_2) d\vec{\Omega}_1 d\vec{\Theta}_1. \quad (8.49)$$

The evolution coefficients of the transition probability are

$$\widetilde{W}_{\frac{t_\epsilon}{2}l} = \left\{ 1 + \frac{\epsilon \cdot \Upsilon_l}{2} \right\} \frac{2l+1}{4\pi} \cdot \frac{1}{2\pi} \quad (8.50)$$

and therefore

$$\begin{aligned} \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1, \vec{\Theta}_0 \cdot \vec{\Theta}_1) &= \sum_{l=0}^{+\infty} \left\{ 1 + \frac{\epsilon \cdot \Upsilon_l}{2} \right\} \frac{2l+1}{4\pi} \cdot \frac{1}{2\pi} \sum_{m=-l}^{+l} P_{lm}(\vec{\Omega}_1) P_{lm}^*(\vec{\Omega}_0) \\ &\quad \cdot \frac{1}{2} \sum_{k=-\infty}^{+\infty} [H_k(\vec{\Theta}_1) H_k^*(\vec{\Theta}_0) + H_{-k}(\vec{\Theta}_1) H_{-k}^*(\vec{\Theta}_0)]. \end{aligned} \quad (8.51)$$

respectively

$$\begin{aligned} \widetilde{W}_{\frac{t_\epsilon}{2}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2, \vec{\Theta}_1 \cdot \vec{\Theta}_2) &= \sum_{l=0}^{+\infty} \left\{ 1 + \frac{\epsilon \cdot \Upsilon_l}{2} \right\} \frac{2l+1}{4\pi} \cdot \frac{1}{2\pi} \sum_{m=-l}^{+l} P_{lm}(\vec{\Omega}_2) P_{lm}^*(\vec{\Omega}_1) \\ &\quad \cdot \frac{1}{2} \sum_{k=-\infty}^{+\infty} [H_k(\vec{\Theta}_2) H_k^*(\vec{\Theta}_1) + H_{-k}(\vec{\Theta}_2) H_{-k}^*(\vec{\Theta}_1)]. \end{aligned} \quad (8.52)$$

Integrating (8.49) one obtains

$$\begin{aligned} \widetilde{W}_{t_\epsilon}(\vec{\Omega}_0 \cdot \vec{\Omega}_2, \vec{\Theta}_0 \cdot \vec{\Theta}_2) &= \sum_{l=0}^{+\infty} \left\{ 1 + \frac{\epsilon \cdot \Upsilon_l}{2} \right\}^2 \frac{2l+1}{4\pi} \cdot \frac{1}{2\pi} \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}_0) P_{lm}(\vec{\Omega}_2) \\ &\quad \cdot \frac{1}{2} \sum_{k=-\infty}^{+\infty} [H_k(\vec{\Theta}_0) H_k^*(\vec{\Theta}_2) + H_{-k}(\vec{\Theta}_0) H_{-k}^*(\vec{\Theta}_2)]. \end{aligned} \quad (8.53)$$

Using n intermediate stages $\widetilde{W}_{t_\epsilon}$ is expressed by an integral over the product of the single transition probabilities.

$$\widetilde{W}_{t_\epsilon, 0 \rightarrow 1 \dots \rightarrow n} = \widetilde{W}_{\frac{t_\epsilon}{n}}(\vec{\Omega}_0 \cdot \vec{\Omega}_1, \vec{\Theta}_0 \cdot \vec{\Theta}_1) \cdot \widetilde{W}_{\frac{t_\epsilon}{n}}(\vec{\Omega}_1 \cdot \vec{\Omega}_2, \vec{\Theta}_1 \cdot \vec{\Theta}_2) \dots \widetilde{W}_{\frac{t_\epsilon}{n}}(\vec{\Omega}_{n-1} \cdot \vec{\Omega}_n, \vec{\Theta}_{n-1} \cdot \vec{\Theta}_n) \quad (8.54)$$

$$\widetilde{W}_{t_\epsilon}(\vec{\Omega}_0 \cdot \vec{\Omega}_n, \vec{\Theta}_0 \cdot \vec{\Theta}_n) = \int_{2\pi} \int_{4\pi} \int_{2\pi} \int_{4\pi} \dots \int_{2\pi} \int_{4\pi} \widetilde{W}_{\frac{t_\epsilon}{n}} \cdot \widetilde{W}_{\frac{t_\epsilon}{n}} \dots \widetilde{W}_{\frac{t_\epsilon}{n}} d\vec{\Omega}_1 d\vec{\Theta}_1 \dots d\vec{\Omega}_{n-1} d\vec{\Theta}_{n-1} \quad (8.55)$$

$$\begin{aligned} \widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') &= \lim_{n \rightarrow \infty} \sum_{l=0}^{+\infty} \left\{ 1 + \frac{\epsilon \cdot \Upsilon_l}{2} \right\}^n \frac{2l+1}{4\pi} \\ &\cdot \frac{1}{2\pi} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') \cdot \frac{1}{2} \sum_{k=-\infty}^{+\infty} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] \end{aligned} \quad (8.56)$$

For $n \rightarrow \infty$ arises

$$\lim_{n \rightarrow \infty} \left\{ 1 + \frac{\epsilon \cdot \Upsilon_l}{2} \right\}^n = e^{\Upsilon_l \epsilon} \quad (8.57)$$

and using (8.55)

\Rightarrow

$$\boxed{\begin{aligned} &\widetilde{W}_{t_\epsilon}(\vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') = \\ &\sum_{l=0}^{+\infty} e^{\Upsilon_l \epsilon} \frac{2l+1}{4\pi} \cdot \frac{1}{2\pi} \cdot \sum_{m=-l}^{+l} P_{lm}^*(\vec{\Omega}) P_{lm}(\vec{\Omega}') \cdot \frac{1}{2} \sum_{k=-\infty}^{+\infty} [H_k(\vec{\Theta}') H_k^*(\vec{\Theta}) + H_{-k}(\vec{\Theta}') H_{-k}^*(\vec{\Theta})] \end{aligned}} \quad (8.58)$$

Choosing $\epsilon = \frac{t_\epsilon}{t_E(\vec{x}, t, \vec{\Omega})}$ the exchange function $\widetilde{W}_{t_\epsilon}$ may be understood in the dependencies

$$\widetilde{W}_{t_\epsilon} = \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') \quad (8.59)$$

and

$$\overline{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}, \vec{\Omega}', \vec{\Theta}') \approx \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega} \cdot \vec{\Omega}', \vec{\Theta} \cdot \vec{\Theta}') \quad (8.60)$$

is given, too. \implies

$$\boxed{\begin{aligned} \bar{f}_{t_\epsilon}(\vec{x}, \vec{\Omega}, \vec{\Theta}, t) &= \int_{4\pi} \widetilde{W}_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}, \vec{\Omega}', \vec{\Theta}') \bar{f}_{t_\epsilon}(\vec{x} - t_\epsilon \cdot \vec{v}' \vec{\Omega}' \times \vec{\Theta}', \vec{\Omega}', \vec{\Theta}'), t - t_\epsilon) d\vec{\Omega}' d\vec{\Theta}' \\ \vec{v}' &= \vec{v}'(\vec{x}, \vec{\Omega}', \vec{\Theta}', t) = \vec{\omega}'(\vec{x}, \vec{\Omega}', \vec{\Theta}', t) \cdot \vec{r}'(\vec{x}, \vec{\Omega}', \vec{\Theta}', t) \end{aligned}}$$

(8.61)

9. Appendix

9.1. Legendre-Polynomials

The Legendre-polynomials are defined within the interval $[-1, +1]$ by

$$P_n = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n, n \in N. \quad (9.1)$$

They represent a complete orthogonal function system with

$$\int_{-1}^{+1} P_n(x) P_m(x) dx = \begin{cases} \frac{2}{2m+1} & \text{für } m = n \\ 0 & \text{sonst.} \end{cases} \quad (9.2)$$

Every continuously differentiable function $f(x)$ defined within $[-1, +1]$ can be developed by Legendre-polynomials according to

$$f(x) = \sum_{l=0}^{\infty} f_l P_l(x). \quad (9.3)$$

The f_l are the evolution coefficients. A presentation of the δ -function by Legendre-polynomials is obtained by

$$\delta(x, x') = \sum_{l=0}^{\infty} \frac{2l+1}{2} P_l(x) P_l(x'). \quad (9.4)$$

Important recurrence equations are [30]

$$\begin{aligned} (n+1)P_{n+1} &= (2n+1)xP_n(x) - nP_{n-1}(x) \\ P'_{n+1}(x) - xP'_n(x) &= (n+1)P_n(x), n = 0, 1, 2, \dots \\ (1-x^2)P'_n(x) &= nP_{n-1}(x) - nxP_n(x). \end{aligned} \quad (9.5)$$

An integral representation of the Legendre-polynomials is obtained by

$$P_n(x) = \frac{1}{\pi} \int_0^\pi (x + \sqrt{x^2 - 1} \cos(\varphi))^n d\varphi. \quad (9.6)$$

Owing to $|x + \sqrt{x^2 - 1} \cos(\theta)| = |\cos(\theta) + i \sin(\theta) \cos(\theta)| \leq 1$

$$|P_n(x)| \leq 1 \quad (9.7)$$

follows. These polynomials have their maximum for $x = 1$, particularly

$$P_n(1) = 1. \quad (9.8)$$

$$\boxed{\frac{dP_l(x)}{dx} \Big|_1 = \frac{l(l+1)}{2}} \quad (9.9)$$

is proved by complete induction.

Proof :

$$1. P'_0(1) = 0$$

Assumption:

$$2. P'_n(1) = \frac{n(n+1)}{2}$$

\implies

$$3. P'_{n+1}(1) = \frac{(n+2)(n+1)}{2} \quad \text{wegen} \quad (9.5) \quad P'_{n+1}(1) - P'_n(1) = (n+1)P_n(1) \\ \text{q.e.d.}$$

9.2. Spherical harmonics

The Spherical harmonics [[31] page 224] represent a complete orthogonal, complex function system on the spherical surface

$$\begin{aligned} P_{lm}(\vec{\Omega}) &= e^{im\varphi} \frac{(-\sin(\vartheta))^m}{l!2^l} \cdot \left(\frac{(l-m)!}{(l+m)!} \right)^{\frac{1}{2}} \frac{d^{l+m}(\cos^2\vartheta - 1)^l}{(d\cos\vartheta)^{l+m}} \\ &= e^{im\varphi} \frac{(\sin(\vartheta))^{-m}}{l!2^l} \cdot \left(\frac{(l+m)!}{(l-m)!} \right)^{\frac{1}{2}} \frac{d^{l-m}(\cos^2\vartheta - 1)^l}{(d\cos\vartheta)^{l-m}} \end{aligned} \quad (9.10)$$

with

$$P_{l,-m}(\vec{\Omega}) = (-)^m P_{lm}^*(\vec{\Omega}) \quad (9.11)$$

and

$$\int_{4\pi} d\vec{\Omega} P_{l'm'}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}) = \delta_{l'l} \delta_{m'm} \frac{4\pi}{2l+1} . \quad (9.12)$$

All continuously differentiable functions on the spherical surface $f(\Omega) = f(\theta, \phi)$ can be developed according to

$$f(\vec{\Omega}) = \sum_{l=0}^{\infty} \sum_{m=-l}^{m=+l} f_{lm} P_{lm}(\vec{\Omega}) \quad (9.13)$$

the f_{lm} representing the evolution coefficients. The $P_{lm}^*(\vec{\Omega})$ being complex to $P_{lm}(\vec{\Omega})$ $f(\vec{\Omega})$ can be alternatively considered

$$f(\vec{\Omega}) = \sum_{l=0}^{\infty} \sum_{m=-l}^{m=+l} f_{lm} P_{lm}^*(\vec{\Omega}). \quad (9.14)$$

The spherical harmonics for $l = 0, 1$ are

$$\begin{aligned} P_{00} &= P_{00}^* = 1 \\ P_{1,-1}(\vec{\Omega}) &= 2^{-\frac{1}{2}} e^{-i\varphi} \sin(\vartheta), \quad P_{1,-1}^* = 2^{-\frac{1}{2}} e^{i\varphi} \sin(\vartheta) \\ P_{1,0}(\vec{\Omega}) &= P_{1,0}^*(\vec{\Omega}) = \cos(\vartheta) = P_1(\vec{\Omega}) \\ P_{1,1}(\vec{\Omega}) &= -2^{-\frac{1}{2}} e^{i\varphi} \sin(\vartheta), \quad P_{1,1}^*(\vec{\Omega}) = -2^{-\frac{1}{2}} e^{-i\varphi} \sin(\vartheta) . \end{aligned} \quad (9.15)$$

The connection of spherical harmonics and Legendre-polynomials is obtained by

$$P_{l0} = P_{l0}^* = P_l. \quad (9.16)$$

Furthermore the addition theorem

$$P_l(\cos(\vartheta)) = \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}') P_{lm}^*(\vec{\Omega}) \quad (9.17)$$

matters with

$$\cos(\vartheta) = \vec{\Omega}' \cdot \vec{\Omega}. \quad (9.18)$$

The δ -function depending on the spherical harmonics may be stated by

$$\delta(\vec{\Omega}, \vec{\Omega}') = \sum_{l=0}^{\infty} \frac{2l+1}{4\pi} \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}') \quad (9.19)$$

and

$$\delta(\vec{\Omega}, \vec{\Omega}') = \sum_{l=0}^{\infty} \frac{2l+1}{4\pi} P_l(\vec{\Omega} \cdot \vec{\Omega}'). \quad (9.20)$$

9.3. Turbulence-functions

Functions of the unit direction vectors $\vec{\Omega}$ and $\vec{\Theta}$ are represented by a complete orthogonal function system meaning an extension of the spherical harmonics. We call them turbulence functions.

$$\begin{aligned} Q_{lmk}(\vec{\Omega}, \vec{\Theta}) &= P_{lm}(\vec{\Omega}) H_k(\vec{\Theta}) \\ &\quad P_{lm}(\vec{\Omega}) \quad \text{spherical harmonics} \\ \int_{2\pi} H_{k'}(\vec{\Theta}) H_k^*(\vec{\Theta}) d\vec{\Theta} &= \begin{cases} 2\pi & \text{for } k'=k \\ 0 & \text{else} \end{cases} \\ H_k(\vec{\Theta}) &= e^{ik\theta} \end{aligned} \quad (9.21)$$

$$\cos(\vartheta) = \vec{\Omega}' \cdot \vec{\Omega}. \quad (9.22)$$

with

$$\int_{2\pi} \int_{4\pi} Q_{lmk}(\vec{\Omega}, \vec{\Theta}) Q_{lmk}^*(\vec{\Omega}', \vec{\Theta}') d\vec{\Omega}' d\vec{\Theta}' = \begin{cases} \frac{8\pi^2}{2l+1} & \text{for } l = l'; m = m'; k = k' \\ 0 & \text{else} \end{cases} \quad (9.23)$$

Such, suitable distribution functions are described by

$$\begin{aligned} f_{t_\epsilon}(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-\infty}^{+\infty} f_{lmk}(\vec{x}, t) Q_{lmk}(\vec{\Omega}, \vec{\Theta}) \\ f(\vec{x}, t, \vec{\Omega}, \vec{\Theta}) &= \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} P_{lm}(\vec{\Omega}) \sum_{k=-\infty}^{+\infty} f_{lmk}(\vec{x}, t) H_k(\vec{\Theta}). \end{aligned} \quad (9.24)$$

The δ -function depending on the turbulence functions is expressed

$$\delta(\vec{\Omega}, \vec{\Omega}'; \vec{\Theta}, \vec{\Theta}') = \sum_{l=0}^{\infty} \frac{2l+1}{4\pi} \sum_{m=-l}^{m=+l} P_{lm}(\vec{\Omega}) P_{lm}^*(\vec{\Omega}') \sum_{k=-\infty}^{+\infty} \frac{1}{2\pi} H_k(\vec{\Theta}) H_k^*(\vec{\Theta}') \quad (9.25)$$

and such

$$\delta(\vec{\Omega}, \vec{\Omega}'; \vec{\Theta}, \vec{\Theta}') = \frac{1}{8\pi^2} \sum_{l=0}^{\infty} (2l+1) P_l(\vec{\Omega} \cdot \vec{\Omega}') \sum_{k=-\infty}^{+\infty} \exp(ik(\Theta - \Theta')). \quad (9.26)$$

9.4. Euler-angles as fluctuation properties of the turbulent particle transport

The angles respectively unit direction vectors $\vec{\Omega}$ and $\vec{\Theta}$ of turbulent motions are applied using the turbulence functions. The unit vector $\vec{\Omega} \times \vec{\Theta}$ with $\vec{\Omega} \perp \vec{\Theta}$ depending on the angles θ , φ and ϑ is determined. Initially, the direction vector $\vec{\Omega}$

$$\vec{\Omega}^0 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (9.27)$$

may be given before a rotation. The orthogonal direction vector $\vec{\Theta}^0$ may be described in this starting situation by

$$\vec{\Theta}^0 = \begin{pmatrix} \sin\theta \\ \cos\theta \\ 0 \end{pmatrix} \quad (9.28)$$

The rotation $\mathbf{T} = \mathbf{T}_2 \cdot \mathbf{T}_1$ with

$$\mathbf{T}_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\vartheta & \sin\vartheta \\ 0 & -\sin\vartheta & \cos\vartheta \end{pmatrix} \quad (9.29)$$

and

$$\mathbf{T}_2 = \begin{pmatrix} \cos\varphi & \sin\varphi & 0 \\ -\sin\varphi & \cos\varphi & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (9.30)$$

results in

$$\mathbf{T} = \mathbf{T}_2 \cdot \mathbf{T}_1 = \begin{pmatrix} \cos\varphi & \sin\varphi\cos\vartheta & \sin\varphi\sin\vartheta \\ -\sin\varphi & \cos\varphi\cos\vartheta & \cos\varphi\sin\vartheta \\ 0 & -\sin\vartheta & \cos\vartheta \end{pmatrix} \quad (9.31)$$

with the unit vectors

$$\begin{aligned} \vec{\Theta} &= \mathbf{T} \cdot \vec{\Theta}^0 = \begin{pmatrix} \cos\varphi\sin\theta + \sin\varphi\cos\vartheta\cos\theta \\ -\sin\varphi\sin\theta + \cos\varphi\cos\vartheta\cos\theta \\ -\sin\vartheta\cos\theta \end{pmatrix} \\ \vec{\Omega} &= \mathbf{T} \cdot \vec{\Omega}^0 = \begin{pmatrix} \sin\varphi\sin\vartheta \\ \cos\varphi\sin\vartheta \\ \cos\vartheta \end{pmatrix} \end{aligned} \quad (9.32)$$

and

$$\begin{aligned} \vec{\Omega} \times \vec{\Theta} &= \begin{pmatrix} \sin\varphi\sin\vartheta \\ \cos\varphi\sin\vartheta \\ \cos\vartheta \end{pmatrix} \times \begin{pmatrix} \cos\varphi\sin\theta + \sin\varphi\cos\vartheta\cos\theta \\ -\sin\varphi\sin\theta + \cos\varphi\cos\vartheta\cos\theta \\ -\sin\vartheta\cos\theta \end{pmatrix} \\ &= \begin{pmatrix} -\cos\varphi\cos\theta + \sin\varphi\cos\vartheta\sin\theta \\ \cos\varphi\cos\vartheta\sin\theta + \sin\varphi\cos\theta \\ -\sin\vartheta\sin\theta \end{pmatrix}. \end{aligned} \quad (9.33)$$

Part III.

Deterministic
continuum-fluctuations and their
stochastic view within the meaning
of an ensemble-theory

10. Introduction

In part II a probabilistic theory of turbulent particle transport is developed from a stochastic ensemble consideration of an unlimited number of parallelly existent, deterministic continuum fluctuations. In this part III the relation of partial differential equations of deterministic continuum fluctuations to the stochastic ensemble-counterpart is established. The causal Markov Process (section 4.5) matters, essentially. Its local description leads to two vector fields with a dual pair of coupled partial, quasilinear differential vector equations distinguishing between mass transport and transport of pure motion quantities $\partial\vec{\mathbf{A}}/\partial t$ and $\vec{\nabla} \times \vec{\mathbf{A}}$ of fluctuating vector fields $\vec{\mathbf{A}}$.

Chapter 11: Turbulent motions have the local velocities $\vec{\mathbf{v}} = \vec{\boldsymbol{\omega}} \times \vec{\mathbf{r}}$ resulting in a dual equation system of a vortex field $\vec{\boldsymbol{\omega}}$ and a curvature vector field $\vec{\mathbf{b}}$. Including the underlying momentum equations (not Navier-Stokes-equations) this system is not yet complete.

Chapter 12: The deterministic transport of pure motion quantities of sufficiently often continuously differentiable fields $\partial\vec{\mathbf{A}}/\partial t$ and $\vec{\nabla} \times \vec{\mathbf{A}}$ is examined leading to a pair of dual coupled vector equations. Depending on interpretation they may be viewed as deformation fluctuation-equations, as generalisations of the Maxwell Equations of vacuum or applied as equations of $\partial\vec{\mathbf{v}}/\partial t$ and $\vec{\nabla} \times \vec{\mathbf{v}}$ of the turbulent velocity field $\vec{\mathbf{v}}$.

Chapter 13: After a discussion of possible momentum equations as foundation for turbulence-calculations the results of chapter 11 and 12 are combined to a complete turbulence-equation system. This system consists of 12 equations with 12 unknowns.¹ From an initial velocity field $\vec{\mathbf{v}}(\vec{\mathbf{x}}, t_0)$ and its partial, temporal derivation $\frac{\partial}{\partial t}\vec{\mathbf{v}}(\vec{\mathbf{x}}, t)|_{t_0}$ the further evolution of the velocity field, its related vortex- and curvature fields as well as the acceleration field $\vec{\mathbf{q}}$ operating in the turbulence field may be calculated.² The acceleration field generally is not conservative meaning $\vec{\nabla} \times \vec{\mathbf{q}} \neq \mathbf{0}$. Matter density distributions may be determined via the continuum-equation in the frame

¹The Einstein Equations of General Relativity consist of 10 equations. Suitable evolution equations with initial- and possibly boundary conditions remain troublesome in a 3+1-geometry.

²As there are only motion quantities in this equation system, it is successfully used for evolution problems in General Relativity.

of a subsequent evaluation in consequence of thermodynamic state quantities being computable (as far as a local thermodynamic equilibrium is existent).

11. Deterministic turbulent mass-transport and its stochastic formulation

$$\begin{aligned}
 f_{t_\epsilon}(t, \vec{x}, \vec{\omega}, \vec{r}) &= \int_{\vec{\omega}'} \int_{\vec{r}'} W_{t_\epsilon}(t, \vec{x}, \vec{\omega}, \vec{r}, \vec{\omega}', \vec{r}') \cdot f_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon, \vec{\omega}', \vec{r}') d\vec{\omega}' d\vec{r}' \\
 &\quad \Updownarrow \\
 \frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{a} - \frac{1}{2} \vec{\nabla} \times \vec{q} &= 0 \\
 \frac{\partial}{\partial t} \vec{b} + \vec{\nabla} \times \vec{\omega} - \frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] &= 0
 \end{aligned}$$

11.1. Introduction

A stochastic ensemble-consideration of deterministic fields is understood as the examination of an unlimited number of comparable, parallelly existent systems, analogously to chapter 4. In this case turbulently moved fluids are examined considering statistical deliberations and its deterministic counterparts. That a linking of deterministic and stochastic theory may be available and further more that out of this connection additionally important (sometimes otherwise not known) relations arise for deterministic formulations, is shown in the following. This is discussed for a turbulent mass transport.

11.2. The transition: stochastic theory \longleftrightarrow deterministic theory

Every space-time-point (\vec{x}, t) a continuously differentiable fluid element distribution over the motion quantities $\vec{\omega}_{t_\epsilon}$ and \vec{r}_{t_ϵ} is assigned according to

$$f_{t_\epsilon} = f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}). \quad (11.1)$$

Indexing functions with t_ϵ it is automatically assumed that the included motion quantities $(\vec{\omega}, \vec{r})$ are assigned to a t_ϵ -measurement accuracy. The indexing of the motion quantities may be omitted in the functions if the functions are accordingly indexed.

After an execution of a $\lim t_\epsilon \rightarrow 0$ process, such as

$$\lim_{t_\epsilon \rightarrow 0} f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}) = f(\vec{x}, t, \vec{\omega}, \vec{r}) \quad (11.2)$$

f and $(\vec{\omega}, \vec{r})$ are understood as results of an exact measuring process.

The change of motion quantities in point (\vec{x}, t)

$$\left(\vec{\omega}'_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon), \vec{r}'_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon) \right) \longrightarrow \left(\vec{\omega}_{t_\epsilon}(\vec{x}, t), \vec{r}_{t_\epsilon}(\vec{x}, t) \right)$$

is controlled by the transition probability density $W_{t_\epsilon} = W_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}, \vec{\omega}', \vec{r}')$.¹ with

$$\begin{aligned} \lim_{t_\epsilon \rightarrow 0} W_{t_\epsilon} &= \delta(\vec{\omega}, \vec{r}; \vec{\omega}', \vec{r}') \\ f_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}) &= \int_{\vec{r}} \int_{\vec{\omega}} W_{t_\epsilon}(\vec{x}, t, \vec{\omega}, \vec{r}, \vec{\omega}', \vec{r}') \cdot f_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon, \vec{\omega}', \vec{r}') d\vec{\omega}' d\vec{r}' \\ \Delta\vec{x} &= t_\epsilon \cdot \vec{\omega}' \times \vec{r}' \end{aligned} \quad (11.3)$$

These equations characterize stochastic turbulence of the continuum in the frame of an ensemble theory and represent a Markov Process with natural causality.

f_{t_ϵ} is developed in (11.3) until the 1st order around $(\vec{x}, t) \implies$

$$f_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon, \vec{\omega}', \vec{r}') = f_{t_\epsilon}(\vec{x}, t, \vec{\omega}', \vec{r}') - \frac{\partial f'_{t_\epsilon}}{\partial t} \cdot t_\epsilon - \Delta\vec{x} \cdot \vec{\nabla} f_{t_\epsilon}(\vec{x}, t, \vec{\omega}', \vec{r}') + \mathcal{O}(t_\epsilon^2) \quad (11.4)$$

¹The otherwise in distribution theory used test functions in this connection have an immediate physical meaning with the formulation of the transition probability density.

with $f'_{t_\epsilon} = f_{t_\epsilon}(\vec{x}, t, \vec{\omega}', \vec{r}')$ and one obtains

$$\int_{\vec{r}} \int_{\vec{\omega}} W_{t_\epsilon} \left[\frac{\partial f'_{t_\epsilon}}{\partial t} + \vec{\omega}' \times \vec{r}' \cdot \vec{\nabla} f'_{t_\epsilon} \right] d\vec{\omega}' d\vec{r}' + O(t_\epsilon^2) = \frac{\int_{\vec{r}} \int_{\vec{\omega}} W_{t_\epsilon} f'_{t_\epsilon} d\vec{\omega}' d\vec{r}' - f_{t_\epsilon}}{t_\epsilon}. \quad (11.5)$$

$\lim_{t_\epsilon \rightarrow 0}$ applied to (11.5) leads to

$$\frac{\partial f}{\partial t} + \vec{\omega} \times \vec{r} \cdot \vec{\nabla} f = \lim_{t_\epsilon \rightarrow 0} \frac{\int_{\vec{r}} \int_{\vec{\omega}} W_{t_\epsilon} f'_{t_\epsilon} d\vec{\omega}' d\vec{r}' - f_{t_\epsilon}}{t_\epsilon}. \quad (11.6)$$

The right side must contain the characteristics of the turbulent fluid.

$$\lim_{t_\epsilon \rightarrow 0} \frac{\int_{\vec{r}} \int_{\vec{\omega}} W_{t_\epsilon} f'_{t_\epsilon} d\vec{\omega}' d\vec{r}' - f_{t_\epsilon}}{t_\epsilon} = F \quad (11.7)$$

F has to be chosen such, that the deterministic vortex equations result under the influence of the assumed acceleration field. Further on the ansatz

$$F = \frac{1}{2} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] f \quad (11.8)$$

is shown precisely fulfilling this condition. Thus one obtains

$$\frac{\partial f}{\partial t} + \vec{\omega} \times \vec{r} \cdot \vec{\nabla} f = \frac{1}{2} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] f. \quad (11.9)$$

Limiting ourselves to one system of the ensemble the distribution function f degenerates to a δ -function.

$$f \rightarrow \delta(\vec{\omega}_{(\vec{x}, t)}, \vec{r}_{(\vec{x}, t)}; \vec{\omega}, \vec{r}) \quad (11.10)$$

The indexing of quantities like $\vec{\omega}_{(\vec{x}, t)}$ by (\vec{x}, t) means the vector $\vec{\omega}$ in the space-time point (\vec{x}, t) ² whereas $\vec{\omega}(\vec{x}, t)$ represents the spatiotemporal field $\vec{\omega}$ in dependence on (\vec{x}, t) .

It results in the key equation for the transition stochastic-deterministic

$$\boxed{\frac{\partial}{\partial t} \delta + \vec{\omega}_{(\vec{x}, t)} \times \vec{r}_{(\vec{x}, t)} \cdot \vec{\nabla} \delta = \frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x}, t)}}{\omega_{(\vec{x}, t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x}, t)} \right] \delta}. \quad (11.11)$$

²That is the situation considering stochastically.

Definition of the operator $\Xi[...]$:

From the vector $\vec{\mathbf{A}}_{(\vec{x},t)}$ respectively the scalar function value $\mathbf{f}_{(\vec{x},t)}$ existing in the space-time-point (\vec{x}, t) of the system a vector function respectively a scalar function arises by the operator Ξ

$$\Xi\left[\vec{\mathbf{A}}_{(\vec{x},t)}\right] = \vec{\mathbf{A}}(\vec{x}, t) \quad (11.12)$$

respectively

$$\Xi\left[\mathbf{f}_{(\vec{x},t)}\right] = \mathbf{f}(\vec{x}, t) \quad (11.13)$$

an appropriate field existing around the point (\vec{x}, t) . The Operator $\Xi[...]$ evokes this functionality to “life“.

Accordingly the following relationships are noted:

$$\begin{aligned} \Xi\left[\int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x},t)}, \vec{r}_{(\vec{x},t)}; \vec{\omega}, \vec{r}) d\vec{\omega} d\vec{r}\right] &= 1 \\ \Xi\left[\int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x},t)}, \vec{r}_{(\vec{x},t)}; \vec{\omega}, \vec{r}) \vec{\omega} d\vec{\omega} d\vec{r}\right] &= \Xi\left[\vec{\omega}_{(\vec{x},t)}\right] = \vec{\omega}(\vec{x}, t) \\ \Xi\left[\int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x},t)}, \vec{r}_{(\vec{x},t)}; \vec{\omega}, \vec{r}) \vec{r} d\vec{\omega} d\vec{r}\right] &= \Xi\left[\vec{r}_{(\vec{x},t)}\right] = \vec{r}(\vec{x}, t) \end{aligned} \quad (11.14)$$

or

$$\Xi\left[\int_{\vec{r}} \int_{\vec{\omega}} \delta(\vec{\omega}_{(\vec{x},t)}, \vec{r}_{(\vec{x},t)}; \vec{\omega}, \vec{r}) \omega^2 \vec{r} d\vec{\omega} d\vec{r}\right] = \Xi\left[\omega_{(\vec{x},t)}^2 \vec{r}_{(\vec{x},t)}\right] = \omega^2(\vec{x}, t) \vec{r}(\vec{x}, t). \quad (11.15)$$

11.3. The deterministic equations of turbulence

From the general momentum equation

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = \vec{q} \quad (11.16)$$

the vortex equation may be developed using the $\vec{\nabla} \times$ -operator

$$\frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times (\vec{v} \times \vec{\omega}) - \frac{1}{2} \vec{\nabla} \times \vec{q} = 0. \quad (11.17)$$

The relations of deterministic and stochastic description are established the same vortex equation opening up from the above key equation. In the following the method

is presented designing the dual pair of deterministic vector equations from the key equation

$$\frac{\partial}{\partial t}\delta + \vec{\omega}_{(\vec{x},t)} \times \vec{r}_{(\vec{x},t)} \cdot \vec{\nabla}\delta = \frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right] \delta. \quad (11.18)$$

In this situation the vectors of the motion quantities may be pushed before and after the differential operators. The Term

$$\frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right] \delta \quad (11.19)$$

guarantees the finding of equation (11.17) and its dual one. It is

$$\vec{v} \perp \vec{\omega} \perp \vec{r}. \quad (11.20)$$

and setting

$$\vec{a} = \vec{v} \times \vec{\omega} \quad (11.21)$$

this results in

$$\vec{r} \parallel \vec{a}. \quad (11.22)$$

Such \vec{a} and \vec{r} are linked as follows³

$$\vec{r} = \frac{\vec{a}}{\omega^2}. \quad (11.23)$$

\Rightarrow

$$\text{with} \quad \delta = \delta(\vec{\omega}_{(\vec{x},t)}, \vec{r}_{(\vec{x},t)}; \vec{\omega}, \vec{r})$$

$$\begin{aligned} \vec{\omega}_{(\vec{x},t)} \times \vec{r}_{(\vec{x},t)} \cdot \vec{\nabla}\delta &= -\vec{r}_{(\vec{x},t)} \times \vec{\omega}_{(\vec{x},t)} \cdot \vec{\nabla}\delta \\ &= -\vec{\omega}_{(\vec{x},t)} \cdot \vec{\nabla} \times \vec{r}_{(\vec{x},t)} \delta \\ &= -\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{a}_{(\vec{x},t)} \delta. \end{aligned}$$

³Symbols as ω, r, a, v etc. always mean amounts of the corresponding vectors.

Inserting in (11.18) gives

$$\begin{aligned}
 & \frac{\partial}{\partial t} \left(\frac{\vec{\omega}_{(\vec{x},t)} \cdot \vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \delta \right) - \frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times (\vec{a}_{(\vec{x},t)} \delta) - \frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right] \delta = 0 \\
 & \implies \frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \left[\frac{\partial}{\partial t} (\vec{\omega}_{(\vec{x},t)} \delta) - \vec{\nabla} \times (\vec{a}_{(\vec{x},t)} \delta) - \frac{1}{2} [\vec{\nabla} \times \vec{q}_{(\vec{x},t)}] \delta \right] = 0 \quad (11.24) \\
 & \implies \frac{\partial}{\partial t} (\vec{\omega}_{(\vec{x},t)} \delta) - \vec{\nabla} \times (\vec{a}_{(\vec{x},t)} \delta) - \frac{1}{2} [\vec{\nabla} \times \vec{q}_{(\vec{x},t)}] \delta = 0
 \end{aligned}$$

One obtains

$$\Xi \left[\int_{\vec{r}} \int_{\vec{\omega}} \left[\frac{\partial}{\partial t} (\vec{\omega}_{(\vec{x},t)} \delta) - \vec{\nabla} \times (\vec{a}_{(\vec{x},t)} \delta) - \frac{1}{2} [\vec{\nabla} \times \vec{q}_{(\vec{x},t)}] \delta = 0 \right] d\vec{\omega} d\vec{r} \right] \quad (11.25)$$

because integration and differentiation beeing exchangeable follows

$$\left[\frac{\partial}{\partial t} \Xi [\vec{\omega}_{(\vec{x},t)}] - \vec{\nabla} \times \Xi [\vec{a}_{(\vec{x},t)}] - \frac{1}{2} \vec{\nabla} \times \Xi [\vec{q}_{(\vec{x},t)}] \right] = 0 \quad (11.26)$$

and we have the first of the dual turbulence equations

$$\frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{a} - \frac{1}{2} \vec{\nabla} \times \vec{q} = 0 \quad (11.27)$$

accordingly

$$\frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times (\vec{v} \times \vec{\omega}) - \frac{1}{2} \vec{\nabla} \times \vec{q} = 0.$$

Hereby the connection of stochastics and deterministics is achieved. From the key-equation above a second equation, the dual one, may be derived.

Back to the initial equation (11.18)

$$\frac{\partial}{\partial t} \delta + \vec{\omega}_{(\vec{x},t)} \times \vec{r}_{(\vec{x},t)} \cdot \vec{\nabla} \delta = \frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right] \delta$$

Simple conversions give

$$\begin{aligned}
 \frac{\partial}{\partial t} \left(\vec{r}_{(\vec{x},t)} \cdot \frac{\vec{r}_{(\vec{x},t)}}{r_{(\vec{x},t)}^2} \delta \right) + \vec{r}_{(\vec{x},t)} \cdot \vec{\nabla} \times (\vec{\omega}_{(\vec{x},t)} \delta) - \frac{\vec{r}_{(\vec{x},t)} \cdot \vec{r}_{(\vec{x},t)}}{r_{(\vec{x},t)}^2} \frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right] \delta = 0 \\
 \longrightarrow \vec{r}_{(\vec{x},t)} \left[\frac{\partial}{\partial t} \frac{\vec{r}_{(\vec{x},t)}}{r_{(\vec{x},t)}^2} \delta + \vec{\nabla} \times (\vec{\omega}_{(\vec{x},t)} \delta) - \frac{\vec{r}_{(\vec{x},t)}}{r_{(\vec{x},t)}^2} \frac{1}{2} \left[\frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right] \delta \right] = 0
 \end{aligned} \tag{11.28}$$

Using the curvature vector field of the fluid trajectories $\vec{b} = \frac{\vec{r}}{r^2}$ the equation is written

$$\frac{\partial}{\partial t} (\vec{b}_{(\vec{x},t)} \delta) + \vec{\nabla} \times (\vec{\omega}_{(\vec{x},t)} \delta) - \frac{1}{2} \vec{b}_{(\vec{x},t)} \frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \delta = 0 \tag{11.29}$$

and applying the operators Ξ arises

$$\Xi \left[\int_{\vec{r}} \int_{\vec{\omega}} \left[\frac{\partial}{\partial t} (\vec{b}_{(\vec{x},t)} \delta) + \vec{\nabla} \times (\vec{\omega}_{(\vec{x},t)} \delta) - \frac{1}{2} \vec{b}_{(\vec{x},t)} \frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \delta = 0 \right] d\vec{\omega} d\vec{r} \right] \tag{11.30}$$

respectively

$$\frac{\partial}{\partial t} \Xi[\vec{b}_{(\vec{x},t)}] + \vec{\nabla} \times \Xi[\vec{\omega}_{(\vec{x},t)}] - \frac{1}{2} \Xi \left[\left(\vec{b}_{(\vec{x},t)} \frac{\vec{\omega}_{(\vec{x},t)}}{\omega_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{q}_{(\vec{x},t)} \right) \right] = 0. \tag{11.31}$$

Such the second of the dual turbulence equations is approached

$$\frac{\partial}{\partial t} \vec{b} + \vec{\nabla} \times \vec{\omega} - \frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] = 0. \tag{11.32}$$

Closing this dual equation system

$$\boxed{
 \begin{aligned}
 \frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{a} - \frac{1}{2} \vec{\nabla} \times \vec{q} &= 0 \\
 \frac{\partial}{\partial t} \vec{b} + \vec{\nabla} \times \vec{\omega} - \frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] &= 0 \\
 \vec{v} = \vec{\omega} \times \frac{\vec{b}}{b^2}, \quad \vec{a} = \vec{v} \times \vec{\omega}
 \end{aligned}
 } \tag{11.33}$$

further equations are necessary besides the momentum equations. In the case of the Navier-Stokes-equations

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\frac{1}{\rho} \vec{\nabla} p + \vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3}\right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v})$$

i.e.

$$\vec{q} = -\frac{1}{\rho} \vec{\nabla} p + \vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3}\right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v})$$

this could happen by simultaneously using the known continuity, energy as well as state equation. But this proves not to be expedient. In chapter 13 the complete equation system is presented and it is shown that the usual Navier-Stokes-equations are not warranting the correct momentum balancing in turbulence.

The term

$$-\frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right]$$

may lead to removable singularities in space-time-points (\vec{x}, t) when turning points occur in the fluid element trajectories $\vec{\omega} = 0$ and $\vec{b} = \mathbf{0}$ arising simultaneously. In this case the whole term is calculated from its surroundings. The same shall apply for the calculation of the velocity \vec{v} . In such cases there is an alternative way shown in chapter 13, too.

12. Stochastic and deterministic general vector fields

$$\begin{aligned}
 f_{t_\epsilon}(\vec{x}, t, \vec{E}, \vec{B}) &= \int_{\vec{B}} \int_{\vec{E}} W_{t_\epsilon}(\vec{x}, t, \vec{E}, \vec{B}, \vec{E}', \vec{B}') \cdot f_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon, \vec{E}', \vec{B}') d\vec{E}' d\vec{B}' \\
 &\quad \Downarrow \\
 \frac{\partial}{\partial t} \vec{B} - \vec{\nabla} \times \vec{E} &= 0 \\
 \frac{\partial}{\partial t} \left(\frac{B^2}{E^2} \cdot \vec{E} \right) + \vec{\nabla} \times \vec{B} &= 0
 \end{aligned}$$

12.1. Introduction

Subsequently continuum fluctuations of general 3 dimensional vector fields $\vec{A}(\vec{x}, t)$ with $\vec{\nabla} \times \vec{A} \neq \mathbf{0}$ are analysed. They have to be sufficiently often continuously differentiable. Defining the vector fields \vec{E} and \vec{B} by

$$\begin{aligned}
 \vec{E} &= \partial \vec{A} / \partial t \neq 0 \\
 \vec{B} &= \vec{\nabla} \times \vec{A} \neq 0
 \end{aligned} \tag{12.1}$$

and owing to the exchangeability of the operators $\partial/\partial t$ und $\vec{\nabla} \times$

$$\frac{\partial \vec{B}}{\partial t} = \vec{\nabla} \times \vec{E} \tag{12.2}$$

follows. This is a necessary consequence of the condition of the continuous differentiability of $\vec{A}(\vec{x}, t)$. This relation is known according to the Maxwell Equations. The for this purpose dual equation is subsequently beeing looked for. In an analogous approach derivating the turbulence equations a stochastic continuum process in the frame of an ensemble theory is formulated such that according to a deterministic

theory the already known as well as the related dual equation arise with fluctuating quantities $\vec{\mathbf{E}}$ und $\vec{\mathbf{B}}$.

12.2. The Transition: stochastic theory \longleftrightarrow deterministic theory

This transition takes place in the same way as the derivation of the dual turbulence equation pair. Every space-time-point $(\vec{\mathbf{x}}, t)$ a continuously differentiable distribution density f_{t_ϵ} is assigned to the motion quantities $\vec{\mathbf{E}}_{t_\epsilon} = \partial \vec{\mathbf{A}}_{t_\epsilon} / \partial t$ and $\vec{\mathbf{B}}_{t_\epsilon} = \vec{\nabla} \times \vec{\mathbf{A}}_{t_\epsilon}$ with

$$f_{t_\epsilon} = f_{t_\epsilon}(\vec{\mathbf{x}}, t, \vec{\mathbf{E}}, \vec{\mathbf{B}}). \quad (12.3)$$

In the with t_ϵ or ϵ indexed functions f_{t_ϵ} it is automatically assumed that the included motion quantities $(\vec{\mathbf{E}}, \vec{\mathbf{B}})$ are assigned to a t_ϵ -measurement accuracy. The indexing of the motion quantities may be omitted in functions appropriately indexed themselves.

After the execution of a $\lim t_\epsilon \rightarrow 0$ -process

$$\lim_{t_\epsilon \rightarrow 0} f_{t_\epsilon}(\vec{\mathbf{x}}, t, \vec{\mathbf{E}}, \vec{\mathbf{B}}) = f(\vec{\mathbf{x}}, t, \vec{\mathbf{E}}, \vec{\mathbf{B}}) \quad (12.4)$$

f and $(\vec{\mathbf{E}}, \vec{\mathbf{B}})$ are understood in the sense of an exact measurement process.

The stochastic transport of the fluctuation quantities

$$\left(\vec{\mathbf{E}}'_{t_\epsilon}(\vec{\mathbf{x}} - \Delta \vec{\mathbf{x}}, t - t_\epsilon), \vec{\mathbf{B}}'_{t_\epsilon}(\vec{\mathbf{x}} - \Delta \vec{\mathbf{x}}, t - t_\epsilon) \right) \longrightarrow \left(\vec{\mathbf{E}}_{t_\epsilon}(\vec{\mathbf{x}}, t), \vec{\mathbf{B}}_{t_\epsilon}(\vec{\mathbf{x}}, t) \right)$$

happens by the transition probability density $W_{t_\epsilon} = W_{t_\epsilon}(\vec{x}, t, \vec{E}, \vec{B}, \vec{E}', \vec{B}')$ with

$$\begin{aligned} \lim_{t_\epsilon \rightarrow 0} W_{t_\epsilon} &= \delta(\vec{E}, \vec{B}; \vec{E}', \vec{B}') \\ f_{t_\epsilon}(\vec{x}, t, \vec{E}, \vec{B}) &= \int \int_{\vec{B}' \vec{E}'} W_{t_\epsilon}(\vec{x}, t, \vec{E}, \vec{B}, \vec{E}', \vec{B}') \cdot f_{t_\epsilon}(\vec{x} - \Delta\vec{x}, t - t_\epsilon, \vec{E}', \vec{B}') d\vec{E}' d\vec{B}' \\ \Delta\vec{x} &= t_\epsilon \cdot \vec{E}' \times \frac{\vec{B}'}{B'^2} \quad \text{and} \quad \vec{E}' \times \frac{\vec{B}'}{B'^2} = \text{velocity of propagation.} \end{aligned} \tag{12.5}$$

These equations define stochastic continuum fluctuations of the quantities \vec{E} und \vec{B} in the sense of an ensemble-theory and represent a Markov Process of natural causality. The test-functions of distribution theory obtain by this formulation of a transition probability density W_{t_ϵ} an immediate physical meaning.

f_{t_ϵ} is developed until the 1st order about $(\vec{x}, t) \Rightarrow$

$$\begin{aligned} f_{t_\epsilon}(t - t_\epsilon, \vec{x} - \Delta\vec{x}, \vec{E}', \vec{B}') &= f'_{t_\epsilon} - \frac{\partial f'_{t_\epsilon}}{\partial t} \cdot t_\epsilon - \Delta\vec{x} \cdot \vec{\nabla} f'_{t_\epsilon} + O(t_\epsilon^2) \\ f'_{t_\epsilon} &= f_{t_\epsilon}(\vec{x}, t, \vec{E}', \vec{B}') \end{aligned} \tag{12.6}$$

und one gets

$$\int_{\vec{E}} \int_{\vec{B}} W_{t_\epsilon} \left[\frac{\partial f'_{t_\epsilon}}{\partial t} + \vec{E}' \times \frac{\vec{B}'}{B'^2} \cdot \vec{\nabla} f'_{t_\epsilon} \right] d\vec{E}' d\vec{B}' + O(t_\epsilon^2) = \frac{\int_{\vec{B}} \int_{\vec{E}} W_{t_\epsilon} f'_{t_\epsilon} d\vec{E}' d\vec{B}' - f_{t_\epsilon}}{t_\epsilon}. \tag{12.7}$$

By the process $t_\epsilon \rightarrow 0$ W_{t_ϵ} degenerates to a δ -function:

$$\lim_{t_\epsilon \rightarrow 0} W_{t_\epsilon} = \delta(\vec{E}, \vec{B}; \vec{E}', \vec{B}') \tag{12.8}$$

$\lim_{t_\epsilon \rightarrow 0}$ applied leads to

$$\frac{\partial f}{\partial t} + \vec{E} \times \frac{\vec{B}}{B^2} \cdot \vec{\nabla} f = \lim_{t_\epsilon \rightarrow 0} \frac{\int_{\vec{E}} \int_{\vec{B}} W_{t_\epsilon} f'_{t_\epsilon} d\vec{E}' d\vec{B}' - f_{t_\epsilon}}{t_\epsilon}. \tag{12.9}$$

Recovering equation (12.2) after the transition to deterministic consideration the exchange term has to vanish, in this case.

$$\lim_{t_\epsilon \rightarrow 0} \frac{\int_{\vec{\mathbf{B}}} \int_{\vec{\mathbf{E}}} W_{t_\epsilon} f'_{t_\epsilon} d\vec{\mathbf{E}}' d\vec{\mathbf{B}}' - f_{t_\epsilon}}{t_\epsilon} = \mathbf{0}. \quad (12.10)$$

This link is an integral part of the considered stochastic process.

Limiting ourselves to one system of the ensemble the function $f(\vec{\mathbf{x}}, t, \vec{\mathbf{E}}, \vec{\mathbf{B}})$ in the space-time-point $(\vec{\mathbf{x}}, t)$ degenerates to a δ -function

$$f(\vec{\mathbf{x}}, t, \vec{\mathbf{E}}, \vec{\mathbf{B}}) \longrightarrow \delta(\vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)}, \vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)}; \vec{\mathbf{E}}, \vec{\mathbf{B}})\text{-function}. \quad (12.11)$$

From equation (12.9) arises the key-equation

$$\boxed{\frac{\partial}{\partial t} \delta + \vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)} \times \frac{\vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)}}{B_{(\vec{\mathbf{x}}, t)}^2} \cdot \vec{\nabla} \delta = \mathbf{0}}. \quad (12.12)$$

Respectively section 11.2 the $\Xi[\dots]$ -operator is inserted as follows

$$\begin{aligned} \Xi \left[\int_{\vec{\mathbf{E}}} \int_{\vec{\mathbf{B}}} \delta(\vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)}, \vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)}; \vec{\mathbf{B}}, \vec{\mathbf{E}}) \vec{\mathbf{B}} d\vec{\mathbf{B}} d\vec{\mathbf{E}} \right] &= \Xi[\vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)}] = \vec{\mathbf{B}}(\vec{\mathbf{x}}, t) \\ \Xi \left[\int_{\vec{\mathbf{E}}} \int_{\vec{\mathbf{B}}} \delta(\vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)}, \vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)}; \vec{\mathbf{B}}, \vec{\mathbf{E}}) \vec{\mathbf{E}} d\vec{\mathbf{B}} d\vec{\mathbf{E}} \right] &= \Xi[\vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)}] = \vec{\mathbf{E}}(\vec{\mathbf{x}}, t) \end{aligned} \quad (12.13)$$

or

$$\Xi \left[\int_{\vec{\mathbf{E}}} \int_{\vec{\mathbf{B}}} \delta(\vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)}, \vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)}; \vec{\mathbf{b}}, \vec{\mathbf{E}}) \left(\frac{B^2}{E^2} \cdot \vec{\mathbf{E}} \right) d\vec{\mathbf{B}} d\vec{\mathbf{E}} \right] = \Xi \left[\frac{B_{(\vec{\mathbf{x}}, t)}^2}{E_{(\vec{\mathbf{x}}, t)}^2} \cdot \vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)} \right] = \frac{B^2(\vec{\mathbf{x}}, t)}{E^2(\vec{\mathbf{x}}, t)} \cdot \vec{\mathbf{E}}(\vec{\mathbf{x}}, t), \quad (12.14)$$

developing the deterministic equations from the key equation.

12.3. The deterministic fluctuation-equations

The key-equation (12.12) represents the interface for the transition of stochastic to deterministic consideration. From the perspective of statistics over the states of movement of the parallelly assumed deterministic processes in the respective point $(\vec{\mathbf{x}}, t)$ one is confined to a single system and such to a single state of motion $(\vec{\mathbf{E}}_{(\vec{\mathbf{x}}, t)}, \vec{\mathbf{B}}_{(\vec{\mathbf{x}}, t)})$.

In this situation the vectors of the motion quantities may be pushed before and behind the differential operators

$$\begin{aligned}\vec{\mathbf{E}}_{(\vec{x},t)} \times \frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \cdot \vec{\nabla} \delta &= -\frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \times \vec{\mathbf{E}}_{(\vec{x},t)} \cdot \vec{\nabla} \delta \\ &= -\frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times \vec{\mathbf{E}}_{(\vec{x},t)} \delta\end{aligned}$$

Further more there is

$$\begin{aligned}\frac{\partial}{\partial t} \left(\frac{\vec{\mathbf{B}}_{(\vec{x},t)} \cdot \vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \delta \right) - \frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \cdot \vec{\nabla} \times (\vec{\mathbf{E}}_{(\vec{x},t)} \delta) &= 0 \\ \implies \frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \cdot \left[\frac{\partial}{\partial t} (\vec{\mathbf{B}}_{(\vec{x},t)} \delta) - \vec{\nabla} \times (\vec{\mathbf{E}}_{(\vec{x},t)} \delta) \right] &= 0 \\ \implies \frac{\partial}{\partial t} (\vec{\mathbf{B}}_{(\vec{x},t)} \delta) - \vec{\nabla} \times (\vec{\mathbf{E}}_{(\vec{x},t)} \delta) &= 0.\end{aligned}\tag{12.15}$$

Now the vector fields of the motion quantities $(\vec{\mathbf{E}}_{(\vec{x},t)}, \vec{\mathbf{B}}_{(\vec{x},t)})$ of the one deterministic system are created about the point (\vec{x}, t) and such the transition to the deterministic equations of the one system has succeeded.

One obtains

$$\Xi \left[\int_{\vec{\mathbf{B}}} \int_{\vec{\mathbf{E}}} \left[\frac{\partial}{\partial t} (\vec{\mathbf{B}}_{(\vec{x},t)} \delta) - \vec{\nabla} \times (\vec{\mathbf{E}}_{(\vec{x},t)} \delta) = 0 \right] d\vec{\mathbf{E}} d\vec{\mathbf{B}} \right].\tag{12.16}$$

As integration and differentiation are exchangeable \implies

$$\frac{\partial}{\partial t} \Xi[\vec{\mathbf{B}}_{(\vec{x},t)}] - \vec{\nabla} \times \Xi[\vec{\mathbf{E}}_{(\vec{x},t)}] = 0\tag{12.17}$$

and it results in the 1.st of the dual fluctuation equations

$$\frac{\partial}{\partial t} \vec{\mathbf{B}} - \vec{\nabla} \times \vec{\mathbf{E}} = 0.\tag{12.18}$$

Hereby the stochastic-deterministic connection is established.

Back to the key-equation (12.12)

$$\frac{\partial}{\partial t} \delta + \vec{\mathbf{E}}_{(\vec{x},t)} \times \frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \cdot \vec{\nabla} \delta = 0$$

one obtains by simple conversion

$$\begin{aligned} \frac{\partial}{\partial t} \left(\vec{\mathbf{E}}_{(\vec{x},t)} \cdot \frac{\vec{\mathbf{E}}_{(\vec{x},t)}}{E_{(\vec{x},t)}^2} \delta \right) + \vec{\mathbf{E}}_{(\vec{x},t)} \cdot \vec{\nabla} \times \left(\frac{\vec{\mathbf{B}}_{(\vec{x},t)}}{B_{(\vec{x},t)}^2} \delta \right) &= 0 \\ \frac{\partial}{\partial t} \left(\frac{B_{(\vec{x},t)}^2}{E_{(\vec{x},t)}^2} \cdot \vec{\mathbf{E}}_{(\vec{x},t)} \delta \right) + \vec{\nabla} \times (\vec{\mathbf{B}}_{(\vec{x},t)} \delta) &= 0 \end{aligned} \quad (12.19)$$

and

$$\Xi \left[\int_{\vec{\mathbf{B}}} \int_{\vec{\mathbf{E}}} \left[\frac{\partial}{\partial t} \left(\frac{B_{(\vec{x},t)}^2}{E_{(\vec{x},t)}^2} \cdot \vec{\mathbf{E}}_{(\vec{x},t)} \delta \right) + \vec{\nabla} \times (\vec{\mathbf{B}}_{(\vec{x},t)} \delta) = 0 \right] d\vec{\mathbf{E}} d\vec{\mathbf{B}} \right] \quad (12.20)$$

respectively

$$\frac{\partial}{\partial t} \Xi \left[\frac{B_{(\vec{x},t)}^2}{E_{(\vec{x},t)}^2} \cdot \vec{\mathbf{E}}_{(\vec{x},t)} \right] + \vec{\nabla} \times \Xi[\vec{\mathbf{B}}_{(\vec{x},t)}] = 0. \quad (12.21)$$

So we have the second of the two dual equations

$$\frac{\partial}{\partial t} \left(\frac{B^2}{E^2} \cdot \vec{\mathbf{E}} \right) + \vec{\nabla} \times (\vec{\mathbf{B}}) = 0. \quad (12.22)$$

The result is recapitulated by the following equation system:

$$\boxed{\begin{aligned} \frac{\partial}{\partial t} \vec{\mathbf{B}} - \vec{\nabla} \times \vec{\mathbf{E}} &= 0 \\ \frac{\partial}{\partial t} \left(\frac{B^2}{E^2} \cdot \vec{\mathbf{E}} \right) + \vec{\nabla} \times \vec{\mathbf{B}} &= 0 \\ \vec{\mathbf{E}} \times \frac{\vec{\mathbf{B}}}{B^2} &= \text{propagation speed} \end{aligned}} \quad (12.23)$$

with $|\vec{\mathbf{E}} \times \frac{\vec{\mathbf{B}}}{B^2}| \leq |\vec{\mathbf{E}}| \cdot |\frac{\vec{\mathbf{B}}}{B^2}|$. I.e. $\frac{E^2}{B^2}$ is not the quadratic propagation speed. Interestingly, this only becomes clear after the involvement of the stochastic ensemble theory.

The equation system (12.23) is in such general terms that the physical significance depends on the interpretation of the starting field $\vec{\mathbf{A}}$, the boundary conditions as well as on the initial conditions. Hereunder, a deformation vector field, the velocity vector field of turbulence motions or the fluctuations of any other continuously differentiable

vector field may be understood. These equations possess with boundary- and suitable initial conditions exactly one solution after the theorem of Cauchy-Kowalewskaja [4]. This statement is at first restricted to the calculation of the fields $\vec{\mathbf{E}}$ and $\vec{\mathbf{B}}$. Calculating the field $\vec{\mathbf{A}}$ with the mere knowledge of

$$\frac{\partial \vec{\mathbf{A}}}{\partial t} = \vec{\mathbf{E}} \quad (12.24)$$

is not possible in all cases. A negative example is the calculation of $\vec{\mathbf{v}}$ with the knowledge of $\frac{\partial \vec{\mathbf{v}}}{\partial t}$ related to turbulent velocity fluctuations as shown in chapter 13. However, in this case these relations are applied completing the turbulence equations. The particular definition of turbulence fluctuation elements (chapter 3) makes this problem almost vividly comprehensible.

Considering turbulent motions this can be done from a different perspective. With the equation system (12.23) the motion quantities

$$\vec{\mathbf{E}} = \frac{\partial}{\partial t} \vec{\mathbf{v}} \quad \text{and} \quad \vec{\mathbf{B}} = \vec{\nabla} \times \vec{\mathbf{v}}$$

are transported with the propagation speed

$$\vec{\mathbf{v}} = \vec{\mathbf{E}} \times \frac{\vec{\mathbf{B}}}{B^2}.$$

The equation system (11.33) describes the mass transport by the velocity $\vec{\mathbf{v}}$. In consideration of $\vec{\mathbf{b}} = \frac{\vec{\mathbf{a}}}{\vec{v}^2}$ (11.33) may be formulated omitting the viscosity and assuming $\vec{\nabla} \times \vec{\mathbf{q}} = 0$ as follows:

$$\begin{aligned} \frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{\mathbf{a}} &= 0 \\ \frac{\partial}{\partial t} \left(\frac{\vec{\mathbf{a}}}{v^2} \right) + \vec{\nabla} \times \vec{\omega} &= 0 \\ \vec{\mathbf{v}} = \vec{\omega} \times \frac{\vec{\mathbf{b}}}{b^2}, \quad \vec{\mathbf{a}} = \vec{\mathbf{v}} \times \vec{\omega}, \quad \vec{\mathbf{v}} &= \text{propagation speed} \end{aligned} \quad (12.25)$$

In doing so $\vec{\mathbf{v}} \perp \vec{\omega} \perp \vec{\mathbf{a}}$ holds. The equations (12.23) and (12.25) do not formally differ apart from orthogonality conditions. But it is not expected, that the fluctuations generated by a conservative acceleration field ($\vec{\nabla} \times \vec{\mathbf{q}} = 0$) may describe

hydrodynamic turbulences. This is discussed in chapter 13.

12.3.1. The vacuum Maxwell Equations

The propagation speed having the constant amount of light velocity one obtains the known equations of vacuum-electrodynamics in the coordinate system of the observer:

$$\begin{aligned} \frac{\partial}{\partial t} \vec{B} - \vec{\nabla} \times \vec{E} &= 0 \\ \frac{1}{c^2} \frac{\partial}{\partial t} \vec{E} + \vec{\nabla} \times \vec{B} &= 0 \quad \text{mit} \quad \vec{E} \perp \vec{B} \\ \vec{E} \times \frac{\vec{B}}{B^2} &= \vec{c} = \text{propagation speed of light.} \end{aligned} \tag{12.26}$$

Hereby a formal analogy is established between electrodynamics and turbulent fluid dynamics. It is only based on the analogy of the propagation of the motion quantities (\vec{E}, \vec{B}) and $(\frac{\partial}{\partial t} \vec{v}, \vec{\nabla} \times \vec{v})$. But a turbulent mass transport with the local velocity \vec{v} cannot be sufficiently described in this way as stated in chapter 13.

So the electrodynamic equations of vacuum are generally derived, too. Usually, they are seen in the above equations with $-\vec{E}$. It is more than pure supposition, that they describe properties of space-time without a unification of General Relativity and electromagnetic field in vacuum having succeeded, though many physicists not least Einstein [7], Jordan [15] and many others having endeavoured.

There is still the explanation of the associated initial field \vec{A} it generally happening in the frame of vector potential considerations, without recognizing \vec{A} as definite physical object. Only by a direct comprehension of the vector potential the electromagnetic field may be explained without means of mechanical quantities.¹

¹Electrodynamics is introduced in physics via mechanical effects.

12.4. Surfacelike deformation-fluctuations in 3-dimensional space

Let \vec{d} be a continuously differentiable deformation vector field defining an area and \vec{b} und \vec{e} the derived fields

$$\vec{e} = \frac{\partial}{\partial t} \vec{d}, \quad \vec{b} = \vec{\nabla} \times \vec{d} \quad (12.27)$$

with

$$\begin{aligned} \vec{d}(x, y, t) &= \left(d_x(x, y, t), d_y(x, y, t), d_z(x, y, t) \right) \\ \vec{e}(x, y, t) &= \left(e_x(x, y, t), e_y(x, y, t), e_z(x, y, t) \right) \\ \vec{b}(x, y, t) &= \left(b_x(x, y, t), b_y(x, y, t), b_z(x, y, t) \right). \end{aligned} \quad (12.28)$$

Then the deformation is without loss of generality seen as deformation of the $\mathbf{x} - \mathbf{y}$ -area. The equations of motion formally equal the equations of 3-dimensional fluctuations

$$\begin{aligned} \frac{\partial}{\partial t} \vec{b} - \vec{\nabla} \times \vec{e} &= 0 \\ \frac{\partial}{\partial t} \left(\frac{b^2}{e^2} \cdot \vec{e} \right) + \vec{\nabla} \times \vec{b} &= 0 \\ \vec{e} \times \frac{\vec{b}}{b^2} &= \text{propagation speed}, \end{aligned} \quad (12.29)$$

only, the operator $\vec{\nabla} \times$ corresponds to

$$\vec{\nabla} \times \vec{d} = \begin{pmatrix} \partial d_z / \partial y \\ -\partial d_z / \partial x \\ \partial d_y / \partial x - \partial d_x \partial y \end{pmatrix}. \quad (12.30)$$

The solution uniquely succeeds by the initial conditions $\vec{b}(x, y, t_0)$ and $\vec{e}(x, y, t_0)$ according to the theorem of Cauchy-Kowalewskaya [4]. The solution for this area corresponds to a partial solution of a 3-dimensional complete solution. Physical material properties are not explicitly included in these equations. They have to be implicitly

considered by initial and boundary conditions. Sole precondition is that the appropriate materials act continuously. It also means that the physical process has to be clarified enabling the corresponding initial and border conditions.

12.5. 1-dimensional deformation-fluctuations in 3-dimensional space

Let \vec{d} be a continuously differentiable deformation vector field defining a trajectory and \vec{b} und \vec{e} the derived fields

$$\vec{e} = \frac{\partial}{\partial t} \vec{d}, \quad \vec{b} = \vec{\nabla} \times \vec{d} \quad (12.31)$$

with

$$\begin{aligned} \vec{d}(x, t) &= (d_x(x, t), d_y(x, t), d_z(x, t)) \\ \vec{e}(x, t) &= (e_x(x, t), e_y(x, t), e_z(x, t)) \\ \vec{b}(x, t) &= (b_x(x, t), b_y(x, t), b_z(x, t)). \end{aligned} \quad (12.32)$$

Then the deformation is without loss of generality seen as deformation of the \mathbf{x} -coordinate. The equations of motion formally equal the equations of 3-dimensional fluctuations

$$\begin{aligned} \frac{\partial}{\partial t} \vec{b} - \vec{\nabla} \times \vec{e} &= 0 \\ \frac{\partial}{\partial t} \left(\frac{b^2}{e^2} \cdot \vec{e} \right) + \vec{\nabla} \times \vec{b} &= 0 \\ \vec{e} \times \frac{\vec{b}}{b^2} &= \text{propagation speed}, \end{aligned} \quad (12.33)$$

only, the operator $\vec{\nabla} \times$ corresponds to

$$\vec{\nabla} \times \vec{d} = \begin{pmatrix} 0 \\ -\partial d_z / \partial x \\ \partial d_y / \partial x \end{pmatrix}. \quad (12.34)$$

This leads to the component equations

$$\begin{aligned}
 \partial b_y / \partial t &= - \partial e_z / \partial x \\
 \partial b_z / \partial x &= \partial e_y / \partial x \\
 \partial [(b^2/e^2) \cdot e_y] / \partial t &= - \partial b_z / \partial x \\
 \partial [(b^2/e^2) \cdot e_z] / \partial t &= \partial b_y / \partial x
 \end{aligned} \tag{12.35}$$

$$\vec{e} \times \vec{b} / b^2 = \text{propagation speed.}$$

The x-component remains constant. The solution uniquely results from the initial conditions $\vec{b}(\mathbf{x}, t_0)$ and $\vec{e}(\mathbf{x}, t_0)$ according to the theorem of Cauchy-Kowalewskaya [4]. The solution for this 1-dimensional trajectory corresponds to a partial solution of a 3-dimensional complete solution. Physical material properties are not explicitly included in these equations. They have to be implicitly considered by initial and boundary conditions. Sole precondition is that the appropriate materials act continuously. It also means that the physical process has to be clarified enabling the corresponding initial and border conditions.

13. Geometrodynamics of turbulence

$$\begin{aligned} \vec{E} + \frac{1}{2} \vec{\nabla} \vec{v}^2 - 2 \vec{v} \times \vec{\omega} &= \vec{q} \\ \frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{a} &= \frac{1}{2} \vec{\nabla} \times \vec{q} \\ \frac{\partial}{\partial t} \vec{b} + \vec{\nabla} \times \vec{\omega} &= \frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] \\ \frac{\partial}{\partial t} \vec{F} &= -2 \vec{\nabla} \times \vec{\omega} \text{ mit } \vec{F} = \frac{4\omega^2}{E^2} \cdot \vec{E} \end{aligned}$$

13.1. Introduction

For a fluctuating continuum field

$$\frac{d}{dt} \vec{v}(\vec{x}, t) = \frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = \vec{q}(\vec{x}, t) \quad (13.1)$$

may be formally comprehended as a momentum equation. As soon as hydrodynamics is involved where a local thermodynamic balance is assumed, the Eulerian equations

$$\vec{q} \stackrel{?}{=} -\frac{1}{\rho} \vec{\nabla} p \quad (13.2)$$

are noted with the indication of the 2nd Newtonian law. They are only justified under restrictive rules like incompressibility of fluids or $\frac{1}{\rho} \vec{\nabla} p = \vec{\nabla} h$ (h=spec. enthalpy) and or negligible rubbing viscosity. So only limiting cases of fluid dynamics are characterized.

But generally, $\vec{\nabla} \times \vec{q} \neq \mathbf{0}$ is to be presumed. \vec{q} is in contrast to Newtonian mechanics a non-conservative acceleration field. \vec{q} has transversal and longitudinal parts

$$\vec{q} = \vec{q}_{\perp} + \vec{q}_{\parallel}. \quad (13.3)$$

The same applies for the velocity field \vec{v}

$$\vec{v} = \vec{v}_\perp + \vec{v}_\parallel = \vec{\omega} \times \vec{R}. \quad (13.4)$$

The disassembly of the velocity field is adequately taken into account by the development of the dual turbulence equation system. In the momentum equation (13.1) 12 unknowns are “hiddenly“ contained and with the turbulence equation only 9 coupled equations are available. For the field $\rho\vec{q}$ a disassembly in longitudinal und transversal part has to be considered, too.

$$\rho \frac{d}{dt} \vec{v}(\vec{x}, t) = \rho\vec{q} = (\rho\vec{q})_\perp + (\rho\vec{q})_\parallel \quad (13.5)$$

Using the Navier-Stokes-equations (Appendix A) leads to

$$\rho\vec{q} = (\rho\vec{q})_\perp + (\rho\vec{q})_\parallel \stackrel{?}{=} -\vec{\nabla} p + \rho \cdot \vec{g} + \eta \Delta \vec{v} + \left(\xi + \frac{\eta}{3}\right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v})$$

\Rightarrow ¹

$$(\rho\vec{q})_\perp \stackrel{?}{=} -\eta \vec{\nabla} \times \vec{\nabla} \times \vec{v} \quad (13.6)$$

and

$$(\rho\vec{q})_\parallel \stackrel{?}{=} -\vec{\nabla} p + \rho \cdot \vec{g} + \left(\xi + \eta \frac{4}{3}\right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}). \quad (13.7)$$

\vec{g} = earth acceleration

As turbulent motions of sufficiently high reynolds number create negligible viscosity effects and on the other hand \vec{q}_\perp represents the decisive propulsion of the vortex motions turbulences are not correctly calculated by the usual equation system consisting of Navier-Stokes-equations, equation of continuity and energy equation. Equation (13.6) can not be correct. \vec{q}_\parallel contributes nothing to the propulsion of the vortex motions. The turbulent dissipation can not be attributed to viscosity but to the matter exchange of the fluid elements and involved thermodynamic changes of state, if a local thermodynamic state is possible. Then the turbulent dissipation decisively decomposes the kinetic energy. \Rightarrow

$$\rho\vec{q} = (\rho\vec{q})_\perp + (\rho\vec{q})_\parallel \neq -\vec{\nabla} p + \rho \cdot \vec{g} + \eta \Delta \vec{v} + \left(\xi + \frac{\eta}{3}\right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \quad (13.8)$$

¹ $\Delta \vec{v} = \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) - \vec{\nabla} \times \vec{\nabla} \times \vec{v}$

The equations, often called conservation laws [2](Navier-Stokes-equations, equation of continuity and energy equation), do not meet these requirements for turbulence with the exception of the equation of continuity.

13.2. The complete set of turbulence-equations

In the turbulence equations (11.33) the viscous terms according to high reynolds numbers may be omitted whereas for sufficiently small reynolds numbers (laminar motions) they obtain significance.

The equation system

$$\frac{\partial \vec{v}}{\partial t} + \frac{1}{2} \vec{\nabla} \vec{v}^2 - 2\vec{v} \times \vec{\omega} = \vec{q} \quad (13.9)$$

$$\frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{a} = \frac{1}{2} \vec{\nabla} \times \vec{q} \quad (13.10)$$

$$\frac{\partial}{\partial t} \vec{b} + \vec{\nabla} \times \vec{\omega} = \frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] \quad (13.11)$$

with

$$\vec{v} = \vec{\omega} \times \frac{\vec{b}}{b^2}, \quad \vec{a} = \vec{v} \times \vec{\omega}, \quad \vec{\nabla} \times \vec{v} \perp \vec{v} \quad (13.12)$$

is not complete and as the Navier-Stokes-equations as momentum balance are refuted, the usual energy equation, derived from Navier-Stokes-equations and equation of continuity, is rejected, too. So the customarily for completion used energy equation, equation of continuity and state equation can not fill this gap.

There is the possibility observing the evolution of the velocity field not only by mass transport via the equations (13.9), (13.10) and (13.11) but via the progress of their fluctuation quantities $\frac{\partial \vec{v}}{\partial t}$ and $\vec{\nabla} \times \vec{v}$, too. Assuming the equation system (12.25)

$$\frac{\partial}{\partial t} \vec{B} - \vec{\nabla} \times \vec{E} = 0 \quad (13.13)$$

$$\frac{\partial}{\partial t} \left(\frac{B^2}{E^2} \cdot \vec{E} \right) + \vec{\nabla} \times \vec{B} = 0 \quad (13.14)$$

$$\vec{E} \times \frac{\vec{B}}{B^2} = \text{propagationspeed} \quad (13.15)$$

with

$$|\vec{E} \times \frac{\vec{B}}{B^2}| \leq |\vec{E}| \cdot \left| \frac{\vec{B}}{B^2} \right|$$

and

$$\vec{\mathbf{E}} = \frac{\partial \vec{\mathbf{v}}}{\partial t} \text{ and } \vec{\mathbf{B}} = \vec{\nabla} \times \vec{\mathbf{v}}, \text{ as well as } \vec{\mathbf{F}} = \frac{B^2}{E^2} \cdot \vec{\mathbf{E}},$$

one obtains the further equation

$$\boxed{\frac{\partial}{\partial t} \vec{\mathbf{F}} + 2 \vec{\nabla} \times \vec{\omega} = 0}. \quad (13.16)$$

Equation 13.13 with $\vec{\mathbf{B}} = \vec{\nabla} \times \vec{\mathbf{v}} = 2\vec{\omega}$ results in

$$\frac{\partial}{\partial t} 2\vec{\omega} - \vec{\nabla} \times \frac{\partial \vec{\mathbf{v}}}{\partial t} = 0.$$

It corresponds to (13.10) on account of

$$\vec{\nabla} \times \frac{\partial \vec{\mathbf{v}}}{\partial t} = 2 \cdot \left(\vec{\nabla} \times \vec{\mathbf{a}} + \frac{1}{2} \vec{\nabla} \times \vec{\mathbf{q}} \right) = 2 \cdot \frac{\partial \vec{\omega}}{\partial t}$$

with

$$\begin{aligned} \vec{\mathbf{v}} &= \vec{\omega} \times \frac{\vec{\mathbf{b}}}{b^2}, \\ \vec{\mathbf{a}} &= \vec{\mathbf{v}} \times \vec{\omega}, \\ \vec{\mathbf{v}} &\perp \vec{\nabla} \times \vec{\mathbf{v}} \\ \vec{\mathbf{E}} &= \frac{\partial \vec{\mathbf{v}}}{\partial t} \\ \vec{\mathbf{E}} &= 4\omega^2 \vec{\mathbf{F}}^{-1}. \end{aligned}$$

The invers vector respectively the scalar product means $\vec{\mathbf{F}}^{-1} = \vec{\mathbf{F}}/\vec{\mathbf{F}}^2 \implies \vec{\mathbf{F}}^{-1} \cdot \vec{\mathbf{F}} = \mathbf{1}$. This corresponds to the relation of a curvature vector $\vec{\mathbf{b}}$ and its associated radius vector $\vec{\mathbf{r}}$ of a continuously differentiable trajectory in one point $(\vec{\mathbf{x}}, t)$ with $\vec{\mathbf{b}} \cdot \vec{\mathbf{r}} = \mathbf{1}$.

The motion of a turbulence field is characterised by a vortex field $\vec{\omega}(\vec{\mathbf{x}}, t)$ and a curvature vector field² $\vec{\mathbf{b}}(\vec{\mathbf{x}}, t)$.

²Generally, one meets in physics curvature tensor fields at least of 2nd degree as in deformation theory or General Relativity.

So one obtains the complete equation system

$$\begin{aligned}
 & \vec{E} + \frac{1}{2} \vec{\nabla} \vec{v}^2 - 2\vec{v} \times \vec{\omega} = \vec{q} \\
 & \frac{\partial}{\partial t} \vec{\omega} - \vec{\nabla} \times \vec{a} = \frac{1}{2} \vec{\nabla} \times \vec{q} \\
 & \frac{\partial}{\partial t} \vec{b} + \vec{\nabla} \times \vec{\omega} = \frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right] \\
 & \frac{\partial}{\partial t} \vec{F} = -2\vec{\nabla} \times \vec{\omega} \text{ with } \vec{F} = \frac{4\omega^2}{E^2} \cdot \vec{E}
 \end{aligned} \tag{13.17}$$

At this a pairwise orthogonality of the vectors $(\vec{v}, \vec{\omega}, \vec{b})$ i.e.: $\vec{v} \perp \vec{\omega}$, $\vec{v} \perp \vec{b}$, $\vec{b} \perp \vec{\omega}$ exists. Pursueing the trajectory of a fluid element beeing possible only after the calculation of the deterministic turbulence field the trajectory is accompanied by a frame of $\vec{v}, \vec{\omega}$ and \vec{b} except in points where $\vec{\omega} = \mathbf{0}$ and $\vec{b} = \mathbf{0}$ (turning points). Nevertheless, in this case $\vec{v} \neq \mathbf{0}$ has to be otherwise the turbulence has come to an end.

13.3. Comments on the application of the complete equation system

On account of the theorem of Cauchy-Kowalewskaja [4] a unique solution is existing. The equation system may be numerically solved for the fields $\vec{\omega}$, \vec{b} , \vec{q} and $\vec{E} = \frac{\partial \vec{v}}{\partial t}$ (this is treated as an independent field as well as $\vec{\omega}, \vec{b}$ und \vec{q}) simultaneously obtaining the fields \vec{a} and \vec{v} . The special approach of [28] enables 2 times continuously differentiable solutions not meaning analytical results. The order of differentiability may be principally driven forward. This particularly goes at the expense of the calculation effort.

Numerically solving this equation system [28] inflexible difference schemes are forbidden as beeing usual according to DNS-calculations (Direct Numerical Simulations related to Navier-Stokes-, continuum- and energy equation), as in the above equation system from the field environment removable singularities of $\vec{v} = \vec{\omega} \times \frac{\vec{b}}{b^2}$, $\frac{1}{2} \vec{b} \left[\frac{\vec{\omega}}{\omega^2} \cdot \vec{\nabla} \times \vec{q} \right]$ and $(2\vec{\omega})^2 \vec{F}^{-1} = \frac{\partial \vec{v}}{\partial t}$ in different space-time-points (\vec{x}, t) are to be recognized. This outcome is a result of possible turning points of the fluid element trajectories leading to simultaneous values of $\vec{\omega} = 0$ and $\vec{b} = 0$. Die fineness of the time discretisations is determined by the vortex field $\vec{\omega}$.

The in some turbulence models mentioned space- and time-scaling in this theory is

led back to the fluctuations of curvature fields $\vec{\mathbf{b}}$ and vortex fields $\vec{\omega}$. Quantitative dependencies become accessible through numerical calculations.

Though friction losses according to heavy turbulent motions (high reynolds numbers) may be omitted the kinetic energy density may significantly decrease. Thus a part has to be converted into inner energy of thermodynamics if a local thermodynamic balance is existent. It is recalled (chapter 3.4), that turbulent fluid motions are characterized the surroundings of fluid elements continuously exchanging their matter and thus their thermodynamic state quantities, too.

The equation system (13.17) stands out only consisting of motion quantities, i.e. velocities and their temporal and spatial differentiations, a vector curvature field, its assigned vortex field and an abstract acceleration field $\vec{\mathbf{q}}$. Mass distributions respectively densities and thermodynamic quantities as pressure and inner energy do not appear. This fact finds its application in the general-relativistic considerations, too. The density distributions may be calculated by subsequent evaluation via the known velocity fields and the equation of continuity

$$\frac{\partial}{\partial t} \rho = -\vec{\nabla} \cdot (\rho \vec{\mathbf{v}}). \quad (13.18)$$

The complete turbulence equation system may be solved even if no local thermodynamics is existent. Then the subsequent evaluation is limited to density calculations. One obtains the thermodynamic pressure distribution if existent by the subsequently calculated density field ρ and the acceleration field $\vec{\mathbf{q}}$ assuming

$$(\rho \vec{\mathbf{q}})_{\parallel} = -\vec{\nabla} \mathbf{p} + \rho \vec{\mathbf{g}} + (\xi + \eta \frac{4}{3}) \vec{\nabla} (\vec{\nabla} \cdot \vec{\mathbf{v}}). \quad (13.19)$$

via Poisson-equation ³ :

$$\Delta \mathbf{p} = -\vec{\nabla} \cdot (\rho \vec{\mathbf{q}}) + \vec{\nabla} \cdot \rho \vec{\mathbf{g}} + \vec{\nabla} \cdot (\xi + \eta \frac{4}{3}) \vec{\nabla} (\vec{\nabla} \cdot \vec{\mathbf{v}}). \quad (13.20)$$

At high reynolds numbers

$$\Delta \mathbf{p} = -\vec{\nabla} \cdot (\rho \vec{\mathbf{q}}) + \vec{\nabla} \cdot \rho \vec{\mathbf{g}} \quad (13.21)$$

is certainly sufficient. But it is not obvious, whether $(\rho \vec{\mathbf{q}})_{\parallel}$ may be represented this way. Upon positive comparison density- and pressure evolution are determined without knowledge of a related state equation. Knowing the state equation all desired thermodynamic state quantities of a single-phase system result. On the other hand a physical process is to be found to create the used initial conditions.

³The transversal part $(\rho \vec{\mathbf{q}})_{\perp}$ disappears with divergence formation

The Turbulence depends on an initially assumed motion field

$$\left(\vec{\omega}(\vec{x}, t_0), \vec{b}(\vec{x}, t_0), \frac{\partial \vec{v}}{\partial t} \Big|_{t_0} \right) \Rightarrow \vec{q}(\vec{x}, t_0),^4 \quad (13.22)$$

determining the further course, alone. Evaluating $\vec{q}(\vec{x}, t_0)$ happens by summation of the terms in the momentum equation. An interaction of geometrodynamics and thermodynamics, maybe assumed in accordance with the Navier-Stokes-equations, does not apply.

13.4. The impossibility of calculating turbulence fields only knowing $\frac{\partial \vec{v}}{\partial t}(\vec{x}, t)$

The impression may arise applying turbulence calculations that it is sufficient to use the equation system

$$\begin{aligned} \frac{\partial}{\partial t} \vec{B} - \vec{\nabla} \times \vec{E} &= 0 \\ \frac{\partial}{\partial t} \left(\frac{B^2}{E^2} \cdot \vec{E} \right) + \vec{\nabla} \times \vec{B} &= 0 \\ \vec{E} \times \frac{\vec{B}}{B^2} &= \text{propagation speed,} \\ |\vec{E} \times \frac{\vec{B}}{B^2}| &\leq |\vec{E}| \cdot \left| \frac{\vec{B}}{B^2} \right| \end{aligned}$$

and

$$\vec{E} = \frac{\partial \vec{v}}{\partial t} \text{ und } \vec{B} = \vec{\nabla} \times \vec{v}, \text{ sowie } \vec{F} = \frac{B^2}{E^2} \cdot \vec{E}$$

to determinate the velocity field numerically from the knowledge of $\frac{\partial \vec{v}}{\partial t}|_i$ by

$$\vec{v}(\vec{x}, t)_{i+1} = \frac{\partial \vec{v}(\vec{x}, t)}{\partial t} \Big|_i \Delta t_i + \vec{v}(\vec{x}, t_0)_i. \quad (13.23)$$

Usually numerical time-integrations via $\Delta \vec{v} = \frac{\partial \vec{v}}{\partial t} \cdot \Delta t$ lead in relation to turbulence calculations firstly to error accumulation for a $\vec{v}(\vec{x}, t)$ evaluation (refined methods of numerical mathematics integrating such vector functions do not help) and secondly achieve $\vec{\nabla} \times \vec{v} \not\approx \vec{v}$ with progressing time evolution. 1st is one reason why weather

⁴Inserting in equation (13.9)

forecasts at meteorology are difficult (besides the principally faults of the used momentum equations). The forecasts are limited to few days. The choice of shorter time steps does not help. This difficulty does not exist regarding laminar fluid dynamics. The reason for this fundamental problem of turbulence is explained as follows:

Solving the equation system (13.17) numerically the pairwise orthogonality of the vectors $\vec{v}, \vec{\omega}, \vec{b}$ ($\vec{v} \perp \vec{\omega}$, $\vec{v} \perp \vec{b}$, $\vec{b} \perp \vec{\omega}$) has to be considered as constraint. For analytic solutions, which cannot be formulated, these conditions should be fulfilled by the initial values, alone.

Calculating $\vec{v}(\vec{x}, t)$ by $\vec{\omega}$ and \vec{r}

$$\vec{v}(\vec{x}, t) = \vec{\omega}(\vec{x}, t) \times \vec{r}(\vec{x}, t) \text{ with } \vec{r}(\vec{x}, t) = \frac{\vec{b}(\vec{x}, t)}{b^2}$$

there is a time integration of the velocity field of higher accuracy. It holds

$$\frac{\partial \vec{v}}{\partial t} = \frac{\partial \vec{\omega}}{\partial t} \times \vec{r} + \vec{\omega} \times \frac{\partial \vec{r}}{\partial t}.$$

The numerical time evolution of $\vec{v}_i \Rightarrow \vec{v}_{i+1}$ arises calculating $\vec{v}_i = \vec{\omega}_i \times \vec{r}_i$ by means of

$$\vec{\omega}_{i+1} = \frac{\partial \vec{\omega}}{\partial t}|_i \cdot \Delta t_i + \vec{\omega}_i + \dots$$

and

$$\vec{r}_{i+1} = \frac{\partial \vec{r}}{\partial t}|_i \cdot \Delta t_i + \vec{r}_i + \dots$$

to

$$\vec{v}_{i+1} = \left(\frac{\partial \vec{\omega}}{\partial t}|_i \cdot \Delta t_i + \vec{\omega}_i \right) \times \left(\frac{\partial \vec{r}}{\partial t}|_i \cdot \Delta t_i + \vec{r}_i \right) + \dots$$

i.e.

$$\vec{v}_{i+1} = \left(\vec{\omega}_i \times \vec{r}_i \right) + \left(\frac{\partial \vec{\omega}}{\partial t}|_i \cdot \Delta t_i \times \vec{r}_i + \vec{\omega}_i \times \frac{\partial \vec{r}}{\partial t}|_i \cdot \Delta t_i \right) + \left(\frac{\partial \vec{\omega}}{\partial t}|_i \cdot \Delta t_i \times \frac{\partial \vec{r}}{\partial t}|_i \cdot \Delta t_i \right) + \dots$$

respectively

$$\boxed{\vec{v}_{i+1} = \vec{v}_i + \frac{\partial \vec{v}}{\partial t}|_i \cdot \Delta t_i + \left(\frac{\partial \vec{\omega}}{\partial t}|_i \times \frac{\partial \vec{r}}{\partial t}|_i \right) \cdot (\Delta t_i)^2 + \dots} \quad (13.24)$$

$\frac{\partial \vec{r}}{\partial t}$ is derived as follows:

$$\vec{b} = \vec{r} \cdot (\vec{b} \cdot \vec{b}) \quad (13.25)$$

\Rightarrow

$$\frac{\partial \vec{\mathbf{b}}}{\partial t} = \mathbf{b}^2 \frac{\partial \vec{\mathbf{r}}}{\partial t} + 2\vec{\mathbf{r}} \left(\frac{\partial \vec{\mathbf{b}}}{\partial t} \cdot \vec{\mathbf{b}} \right)$$

\Rightarrow

$$\frac{\partial \vec{\mathbf{r}}}{\partial t} = \left[\frac{\partial \vec{\mathbf{b}}}{\partial t} - 2 \frac{\vec{\mathbf{b}}}{b^2} \left(\frac{\partial \vec{\mathbf{b}}}{\partial t} \cdot \vec{\mathbf{b}} \right) \right] / b^2.$$

In particular space-time points $(\vec{\mathbf{x}}, t)$ fluid elements may be in the proximity or direct in a turning point, in which $\vec{\omega}(\vec{\mathbf{x}}, t) = \mathbf{0}$ as well as $\vec{\mathbf{b}}(\vec{\mathbf{x}}, t) = \mathbf{0}$ and such $\vec{\mathbf{r}}(\vec{\mathbf{x}}, t) = \vec{\mathbf{b}}/b^2 = \infty$ holds. So the temporal evolution term of 2nd order is vital for turbulence calculations. That is why the with (13.23) mentioned velocity integration is not expedient. Considering the complete turbulence equation set the temporal velocity integration automatically results in the desired order.

13.5. Summary

With the installation of the equation system (13.17) a geometrodynamics of turbulence is expressed only obtaining motion quantities i.e. it only consists of velocities and their time and space derivatives. A corresponding statement is made for their initial- and boundary conditions. Special material properties of a fluid may only influence solutions via initial- and boundary conditions. Therefore it may be important to formulate the suitable process of the genesis of such conditions.

13.6. Appendix A: The basic equations of laminar fluid-dynamics

In the following, the known and established fundamental equations of laminar fluid-dynamics are derived and presented.

13.6.1. The equation of continuity

Being $\rho = \rho(\vec{\mathbf{x}}, t)$ the mass density and $\vec{\mathbf{v}} = \vec{\mathbf{v}}(\vec{\mathbf{x}}, t)$ the velocity of the fluid elements the mass density passing through the surface of a volume per unit time is

$$\oint \rho \cdot \vec{\mathbf{v}} \cdot d\vec{\mathbf{f}}.$$

On the other hand this corresponds to the mass density decrease of the volume per time of

$$-\frac{\partial}{\partial t} \int_V \rho \cdot dV.$$

So one obtains

$$\frac{\partial}{\partial t} \int_V \rho \cdot dV = - \oint \rho \cdot \vec{v} \cdot d\vec{f}.$$

Using Green's formula

$$\oint \rho \cdot \vec{v} \cdot d\vec{f} = \int_V \vec{\nabla} \cdot (\rho \cdot \vec{v}) \cdot dV$$

there is

$$\int_V \left(\frac{\partial}{\partial t} \rho + \vec{\nabla} \cdot (\rho \cdot \vec{v}) \right) \cdot dV = 0$$

respectively

$$\boxed{\frac{\partial}{\partial t} \rho + \vec{\nabla} \cdot (\rho \cdot \vec{v}) = 0}. \quad (13.26)$$

Originating from a constant density $\rho = \text{const}$ one obtains

$$\vec{\nabla} \cdot \vec{v} = 0. \quad (13.27)$$

That means, a divergence-free velocity field is the necessary condition of a constant mass-distribution, but this condition is not sufficient! The equation of continuity proves to be the only correct conservation equation of the three fundamental fluid-dynamics equations: equation of continuity, Navier-Stokes-equations and energy equation. The structure of this equation of continuity may be the formal standard for the categorisation of scalar conservation quantities at least in continuum physics. In the following it is shown, that this can not be achieved for the energy equation.

13.6.2. The Navier-Stokes-equations

Considering the influence of viscose friction one usually [17] takes as its starting point the momentum density conservation. Expediently, the Ricci Calculus is used as formulation. \vec{g} means an additional acceleration field, such as the earth acceleration. These equations finding their empirical confirmation in the laminar or stationary case are based directly on

$$\rho \cdot \left(\frac{\partial v_i}{\partial t} + v_k \cdot \frac{\partial v_i}{\partial x_k} \right) = - \frac{\partial}{\partial x_k} (p \delta_{ik} - \sigma_{ik}) + \rho \cdot g_i. \quad (13.28)$$

The stress tensor σ_{ik} must consist of a symmetrical tensor whose trace vanishes in the situation $\vec{v} = \vec{\omega} \times \vec{r}$ with $\vec{\omega} \neq \mathbf{0}$ plus a diagonal tensor. Such a linear composition of the tensors

$$\frac{\partial v_i}{\partial x_k} + \frac{\partial v_k}{\partial x_i} \text{ and } \delta_{ik} \frac{\partial v_l}{\partial x_l} \quad (13.29)$$

is needed with coefficients determined experimentally [17]. σ_{ik} is the most general tensor of linear, spatial velocity derivations of 1st order within these conditions:

$$\sigma_{ik} = \eta \left(\frac{\partial v_i}{\partial x_k} + \frac{\partial v_k}{\partial x_i} - \frac{2}{3} \delta_{ik} \frac{\partial v_l}{\partial x_l} \right) + \xi \delta_{ik} \frac{\partial v_l}{\partial x_l}. \quad (13.30)$$

The coefficients η und ξ not essentially changing in the fluid one obtains

$$\begin{aligned} \frac{\partial}{\partial x_k} \sigma_{ik} &= \eta \left(\frac{\partial^2 v_i}{\partial x_k^2} + \frac{\partial}{\partial x_i} \frac{\partial v_k}{\partial x_k} - \frac{2}{3} \frac{\partial}{\partial x_i} \frac{\partial v_l}{\partial x_l} \right) + \xi \frac{\partial}{\partial x_i} \frac{\partial v_l}{\partial x_l} \\ &= \eta \frac{\partial^2 v_i}{\partial x_k^2} + \left(\xi + \frac{1}{3} \eta \right) \frac{\partial}{\partial x_i} \frac{\partial v_l}{\partial x_l}. \end{aligned} \quad (13.31)$$

Because of

$$\sum_{k=1}^3 \frac{\partial^2 v_i}{\partial x_k^2} = \Delta v_i \quad (13.32)$$

and

$$\sum_{l=1}^3 \frac{\partial v_l}{\partial x_l} = \vec{\nabla} \cdot \vec{v} \quad (13.33)$$

the known Navier-Stokes equations

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\frac{1}{\rho} \vec{\nabla} p + \vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \quad (13.34)$$

are finally achieved with $\nu = \frac{\eta}{\rho}$ and $\zeta = \frac{\xi}{\rho}$. This demonstrates the coefficients ν and ζ not beeing constant coefficients at heavy density fluctuations. On the other hand the question arises, if the the coefficients η und ξ may be seen as essentially constant (prudently not!). Whereby the above derivation of the viscose terms of the Navier-Stokes-equations for turbulence problems of sufficiently high reynolds numbers appears dubious. On the other side, numerical examinations [28] show that the viscous

friction in case of heavy turbulence problems is irrelevant. The known considerations of similarity by the Reynolds law of similarity do not have a sufficient sustainability.

13.6.3. The equation of energy

With the energy-equation of laminar hydrodynamics (not the hydrodynamics of turbulent motion) the kinetic energy of the fluid and its inner thermodynamic energy density are balanced. The derivation essentially follows the presentation of [19]. Here non adiabatic fluid motions and the friction viscosity are additionally considered. It is

$$\mathbf{e}_{\text{sum}} = \mathbf{e}_{\text{kin}} + \mathbf{e}_{\text{in}} \quad (13.35)$$

defined with

$$\boxed{\frac{\partial}{\partial t} \mathbf{e}_{\text{sum}} = -\vec{\nabla} \cdot (\vec{v}(\mathbf{e}_{\text{sum}} + p)) + \rho \cdot \vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) + \vec{\nabla} \lambda \vec{\nabla} T} \quad (13.36)$$

In the following this connection is derived. Heat radiation is ignored significantly arising above 500°C . The following specifications are made:

Symbol	Physical Quantity
$\mathbf{e}_{\text{kin}} = \frac{\rho}{2} v^2$	kinetic energy density
ε	spec. inner energy
$\mathbf{e}_{\text{in}} = \rho \varepsilon$	inner energy density
$\mathbf{e}_{\text{sum}} = \mathbf{e}_{\text{kin}} + \mathbf{e}_{\text{in}}$	sum of kinetic energy density and inner energy density
$h = \varepsilon + \frac{p}{\rho}$	spec. enthalpy
ρh	enthalpy density
s	spec. entropy
ρs	entropy density
p	pressure
T	temperature
ρ	density
Q	spec. heat
λ	heat conduction coefficient
\vec{g}	external mass acceleration field

Examined is:

$$\frac{\partial}{\partial t} \mathbf{e}_{\text{sum}} = \frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 \right) + \frac{\partial}{\partial t} (\rho \varepsilon) \quad (13.37)$$

First $\frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 \right) :$

$$\frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 \right) = \frac{v^2}{2} \frac{\partial}{\partial t} \rho + \rho \vec{v} \cdot \frac{\partial}{\partial t} \vec{v} \quad (13.38)$$

$$\frac{\partial}{\partial t} \rho = -\vec{\nabla} \cdot (\rho \vec{v}) \quad \text{equation of continuity} \quad (13.39)$$

Navier-Stokes-equations:

$$\frac{\partial \vec{v}}{\partial t} = -(\vec{v} \cdot \vec{\nabla}) \vec{v} - \frac{1}{\rho} \vec{\nabla} p + \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) \implies (13.40)$$

$$\frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 \right) = -\frac{v^2}{2} \vec{\nabla} \cdot (\rho \vec{v}) - \rho \vec{v} \cdot (\vec{v} \cdot \vec{\nabla}) \vec{v} - \vec{v} \cdot \vec{\nabla} p + \rho \vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) \quad (13.41)$$

$$\begin{aligned} \rho \vec{v} \cdot (\vec{v} \cdot \vec{\nabla}) \vec{v} &= \rho \vec{v} \cdot \left[\frac{1}{2} \vec{\nabla} v^2 - \vec{v} \times \vec{\nabla} \times \vec{v} \right] \\ &= \frac{1}{2} \rho \vec{v} \cdot \vec{\nabla} v^2 \quad \text{on account of} \quad \vec{v} \cdot \vec{v} \times \vec{\nabla} \times \vec{v} = 0 \end{aligned} \quad (13.42)$$

$$dh = T ds + \frac{1}{\rho} dp \quad \text{from thermodynamics} \quad (13.43)$$

$$\vec{\nabla} p = \rho \vec{\nabla} h - \rho T \vec{\nabla} s \quad (13.44)$$

$$\vec{\nabla} \cdot (\rho \vec{v} \cdot \frac{v^2}{2}) = \frac{v^2}{2} \vec{\nabla} \cdot (\rho \vec{v}) + \rho \vec{v} \cdot \frac{1}{2} \vec{\nabla} v^2 \quad (13.45)$$

$$\begin{aligned} \frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 \right) &= \\ &- \vec{\nabla} \cdot (\rho \vec{v} \cdot \frac{v^2}{2}) - \rho \vec{v} \cdot [\vec{\nabla} h - T \vec{\nabla} s] + \rho \vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) \end{aligned}$$

(13.46)

Now $\frac{\partial}{\partial t} (\rho \varepsilon) :$

$$\frac{\partial}{\partial t}(\rho\varepsilon) = \varepsilon \cdot \frac{\partial \rho}{\partial t} + \rho \frac{\partial \varepsilon}{\partial t} \quad (13.47)$$

on account of

$$\begin{aligned} d\varepsilon &= T ds + \frac{p}{\rho^2} d\rho \quad \text{from thermodynamics and} \\ \frac{\partial}{\partial t}\rho &= -\vec{\nabla} \cdot (\rho\vec{v}) \quad \text{equation of continuity} \end{aligned}$$

$$\frac{\partial}{\partial t}(\rho\varepsilon) = h \cdot \frac{\partial \rho}{\partial t} + \rho T \frac{\partial s}{\partial t}. \quad (13.48)$$

Adiabatic transport means

$$\frac{ds}{dt} = \frac{\partial s}{\partial t} + \vec{v} \cdot \vec{\nabla} s = 0 \quad (13.49)$$

Considering non adiabatic transport heat is withdrawn from the Fluid by heat conduction. I.e.

$$\rho T \frac{ds}{dt} = \rho T \frac{\partial s}{\partial t} + \rho T \vec{v} \cdot \vec{\nabla} s = \vec{\nabla} \cdot \lambda \vec{\nabla} T \quad (13.50)$$

$$\frac{\partial s}{\partial t} = -\vec{v} \cdot \vec{\nabla} s + \frac{1}{\rho T} \vec{\nabla} \cdot \lambda \vec{\nabla} T \quad (13.51)$$

$$\boxed{\frac{\partial}{\partial t}(\rho\varepsilon) = -h \cdot \vec{\nabla} \cdot (\rho\vec{v}) - \rho T \vec{v} \cdot \vec{\nabla} s + \vec{\nabla} \cdot \lambda \vec{\nabla} T} \quad (13.52)$$

On account of

$$\begin{aligned} \frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 \right) &= -\vec{\nabla} \cdot (\rho\vec{v} \cdot \frac{v^2}{2}) - \rho\vec{v} \cdot [\vec{\nabla} h - T \vec{\nabla} s] \\ &\quad + \rho\vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) \\ \frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 + \rho\varepsilon \right) &= -\vec{\nabla} \cdot (\rho\vec{v} \cdot \frac{v^2}{2}) - h \cdot \vec{\nabla} \cdot (\rho\vec{v}) - \rho\vec{v} \cdot \vec{\nabla} h \\ &\quad + \rho\vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) + \vec{\nabla} \cdot \lambda \vec{\nabla} T. \end{aligned} \quad (13.53)$$

follows. Respectively

$$\vec{\nabla} \cdot (\rho\vec{v}h) = h \cdot \vec{\nabla} \cdot (\rho\vec{v}) + \rho\vec{v} \cdot \vec{\nabla} h \quad (13.54)$$

follows

$$\begin{aligned} \frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 + \rho \varepsilon \right) = & - \vec{\nabla} \cdot (\rho \vec{v} \cdot \frac{v^2}{2}) - \vec{\nabla} \cdot (\rho \vec{v} h) \\ & + \rho \vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) + \vec{\nabla} \cdot \lambda \vec{\nabla} T \end{aligned} \quad (13.55)$$

and

$$\frac{\partial}{\partial t} \left(\frac{\rho}{2} v^2 + \rho \varepsilon \right) = - \vec{\nabla} \cdot \rho \vec{v} \cdot \left(\frac{v^2}{2} + h \right) + \rho \vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) + \vec{\nabla} \cdot \lambda \vec{\nabla} T \quad (13.56)$$

and finally

$$\boxed{\frac{\partial}{\partial t} \mathbf{e}_{\text{sum}} = - \vec{\nabla} \cdot (\vec{v}(\mathbf{e}_{\text{sum}} + \mathbf{p})) + \rho \cdot \vec{v} \cdot \left(\vec{g} + \nu \Delta \vec{v} + \left(\zeta + \frac{\nu}{3} \right) \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) \right) + \vec{\nabla} \cdot \lambda \vec{\nabla} T.} \quad (13.57)$$

Even without external field \vec{g} and without consideration of heat conduction there is no equation of conservation for the sum of kinetic and inner energy density \mathbf{e}_{sum} ! I.e. \mathbf{e}_{sum} is not a conservation quantity. Energy is lost by fluid motions against non conservative local forces. A potential representing the seemingly lost energy density does not exist! But this is necessary to formulate a real conservation equation. Non conservative acceleration fields do not permit energy-density conservation. Apart from this the energy in a point (\vec{x}, t) is zero in pure field theories, anyway.

$$\frac{\partial}{\partial t} \mathbf{e}_{\text{sum}} + \vec{\nabla} \cdot (\mathbf{e}_{\text{sum}} \vec{v}) \neq 0 \quad (13.58)$$

A conservation of the sum of kinetic and inner energy density had to be represented by a suitable equation of continuity. In many cases the energy equation is described in sufficient approximation by

$$\frac{\partial}{\partial t} \mathbf{e}_{\text{sum}} = - \vec{\nabla} \cdot (\vec{v}(\mathbf{e}_{\text{sum}} + \mathbf{p})). \quad (13.59)$$

Generally, heat conduction can usually be neglected contrary to the energy convection (adiabatic convection), unless the velocities are small enough. (Example: thermics in the atmosphere)

13.7. Appendix B

13.7.1. The Helmholtz-decomposition-theorem

It is shown, that every continuously differentiable vector field may be uniquely disassembled in a longitudinal and a transversal part, i.e.

$$\vec{q}(\vec{x}) = \vec{\nabla}\Phi(\vec{x}) + \vec{\nabla} \times \vec{A}(\vec{x}) = \vec{q}(\vec{x})_{\parallel} + \vec{q}(\vec{x})_{\perp}. \quad (13.60)$$

Proof:

Without loss of generality

$$\begin{aligned} \vec{q}(\vec{x}) &= \int_V \vec{q}(\vec{x}') \delta(\vec{x}, \vec{x}') dV' \\ &= \int_V \vec{q}(\vec{x}') \Delta G(\vec{x}, \vec{x}') dV' \quad \text{with the Green function } G, \quad \Delta G = \delta(\vec{x}, \vec{x}') \\ &= \Delta \int_V \vec{q}(\vec{x}') G(\vec{x}, \vec{x}') dV' \quad \Delta \text{ only differentiated by the unslashed } \vec{x}. \end{aligned} \quad (13.61)$$

can be written. Because of

$$\Delta \vec{Q} = \vec{\nabla}(\vec{\nabla} \cdot \vec{Q}) - \vec{\nabla} \times (\vec{\nabla} \times \vec{Q}) \quad (13.62)$$

follows

$$\vec{q}(\vec{x}) = \vec{\nabla} \left(\vec{\nabla} \cdot \int_V \vec{q}(\vec{x}') G(\vec{x}, \vec{x}') dV' \right) - \vec{\nabla} \times \left(\vec{\nabla} \times \int_V \vec{q}(\vec{x}') G(\vec{x}, \vec{x}') dV' \right). \quad (13.63)$$

and

$$\vec{q}(\vec{x}) = \vec{\nabla} \left(\int_V \vec{q}(\vec{x}') \vec{\nabla} \cdot G(\vec{x}, \vec{x}') dV' \right) + \vec{\nabla} \times \left(\int_V \vec{q}(\vec{x}') \times \vec{\nabla} G(\vec{x}, \vec{x}') dV' \right). \quad (13.64)$$

As⁵

$$\vec{\nabla} G(\vec{x}, \vec{x}') = -\vec{\nabla}' G(\vec{x}, \vec{x}') \quad (13.65)$$

⁵respectively $G(\vec{x}, \vec{x}') = -\frac{1}{4\pi} \frac{1}{|\vec{x} - \vec{x}'|}$

follows

$$\vec{q}(\vec{x}) = -\vec{\nabla} \left(\int_V \vec{q}(\vec{x}') \cdot \vec{\nabla}' G(\vec{x}, \vec{x}') dV' \right) - \vec{\nabla} \times \left(\int_V \vec{q}(\vec{x}') \times \vec{\nabla}' G(\vec{x}, \vec{x}') dV' \right). \quad (13.66)$$

So it results in the longitudinal part

$$\vec{q}_{\parallel}(\vec{x}) = -\vec{\nabla} \left(\int_V \vec{q}(\vec{x}') \cdot \vec{\nabla}' G(\vec{x}, \vec{x}') dV' \right) \quad (13.67)$$

and the transversal part

$$\vec{q}_{\perp}(\vec{x}) = -\vec{\nabla} \times \left(\int_V \vec{q}(\vec{x}') \times \vec{\nabla}' G(\vec{x}, \vec{x}') dV' \right) \quad (13.68)$$

respectively

$$\Phi(\vec{x}) = - \int_V \vec{q}(\vec{x}') \cdot \vec{\nabla}' G(\vec{x}, \vec{x}') dV' \quad (13.69)$$

and the vector potential


$$\vec{A}(\vec{x}) = - \int_V \vec{q}(\vec{x}') \times \vec{\nabla}' G(\vec{x}, \vec{x}') dV'. \quad (13.70)$$

14. Summary

The most important innovations for turbulence research of this treatise are composed in what follows:

1. *-though the matter is discontinuously distributed and moved, it can be considered as a continuum of abstract moved points as fluid elements not representing special atoms or molecules. Such orthogonality of \vec{v} and $\text{rot}(\vec{v})$ is derived from 1 times continuous differentiability on account of continuum property and parallelism of fluid-velocities in close proximity of a space point as necessary condition. Experiments may contradict only, if they are wrong.*
2. *-derivation of the transport equation of molecular self-diffusion as partial differential equation as well as integral equation. These formulations do not constitute continuum fluctuations and are used to compare the intrinsic stochastics with the stochastic ensemble consideration of deterministic individual systems in connection with a passive particle transport. The transition probability of velocities is calculated, explicitly.*
3. *-derivation of the transport equation of a passive particle transport by longitudinal continuum-fluctuations as partial differential equation as well as integral equation. The transition probability of velocities is calculated, explicitly.*
4. *-derivation of the transport equation of a passive particle transport by turbulent continuum-fluctuations as partial differential equation as well as integral equation. At this the local velocity is decomposed in vectorial rotation as well as vectorial curvature of the rotating movement. The transition probability of vortex and curvature fluctuations is calculated, explicitly.*
5. *-derivation of the relation of deterministic fluid-motions and an associated stochastic ensemble consideration resulting in deterministic turbulence equations.*
6. *-derivation of the relation of general deterministic 3-dimensional continuum vectorfield fluctuations and a stochastic ensemble consideration as counterpart. Herefrom result for example general equations of continuum deformations as well as Maxwell-Equations assuming constant propagation speed.*
7. *-establishment of a complete equation system for turbulence combining ideas of 5. and 6.*

OVERVIEW: FLUCTUATIONS of the CONTINUUM



Bibliography

- [1] Bychkov V.L.: Hydrodynamic Analogies between the Equations of Classical Hydrodynamics and Electrodynamics in Electrochemistry-CHEMICAL PHYSICS OF POLYMER MATERIALS
- [2] Chorin Marsden: A Mathematical Introduction to Fluid Mechanics-Springer 1997
- [3] Chandrasekhar S.: Stochastic Problems in Physics and Astronomy Reviews of Modern Physics 15, no. 1 (1943): 1-89
- [4] Dieudonne J.: Grundzüge der modernen Analysis Band 4 1976 page 67
- [5] Döring Werner: EINFÜHRUNG IN DIE THEORETISCHE PHYSIK V Statistische Mechanik Sammlung Götschen 1957 page 23
- [6] Dunkel Jörn: Relativistic Brownian Motion and Diffusion Processes-Dissertation 2008 Universität Augsburg
- [7] Einstein Albert: The Meaning of relativity 1950
- [8] Einstein A.: Über die von der molekularinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen-Annalen der Physik 17, 549 (1905)
- [9] Feynman Richard P.: The Feynman LECTURES ON PHYSICS Volume I Addison-Wesley Publishing Company, Inc 1964 page 3-9
- [10] George William K.: Lectures in Turbulence for the 21st Century- 16 January 2013
- [11] Grad Harold: Principles of the Kinetic Theory of Gases , Handbuch der Physik,Flügge Band 12, 1958, page 205
- [12] Greiner Walter/Reinhardt Joachim: Theoretische Physik Band 7: Quantenelektrodynamik Verlag Harri Deutsch . Thun und Frankfurt am Main 1984 page 18
- [13] Gröbner W., Lesky P.: MATHEMATISCHE METHODEN DER PHYSIK I BI Hochschultaschebücher
- [14] Grosche C. Steiner: Handbook of Feynman Path Integrals 1998 Springer page 2

- [15] Jordan Pascual: SCHWERKRAFT UND WELTALL 1952
- [16] Khanh Tuoc, Trinh: A THEORY OF TURBULENCE-K.T.Trinh@massey.ac.nz
page 43
- [17] Landau L. D., Lifshitz E. M.: FLUID MECHANICS-Volume 6 1963 Seite 44
- [18] Landau L. D., Lifshitz E. M.: The Classical Theory of Fields Volume 2 Fourth
Revised English Edition
- [19] Lesch Prof. Dr. Harald, Dr. G.T. Birk, Prof. Dr. Hartmut Zohm: Theoretische
Hydrodynamik
- [20] Marmier Prof. Dr.- KERNPHYSIK I: Vorlesung 1965 Verlag des Vereins der
Mathematiker und Physiker an der ETH Zürich
- [21] Marmanis: Analogy of electrodynamics and hydrodynamic Equations-
Application to Turbulence-M. Sc. University of Illinois
- [22] McDonough J. M.: Introductionary LECTURES on TURBULENCE- 2007-
University of Kentucky page 5
- [23] McDonough J. M.: LECTURES IN ELEMENTARY fluid dynamics- 2009-
University of Kentucky
- [24] Penrose Roger: The Road to Reality Jonathan Cape 2004
- [25] Rossmann Wulf: Lectures on Differential Geometry Oktober 2003
- [26] Sen R. N.: Causality, Measurement Theory and the Differentiable Struc-
ture of Space-Time CAMBRIDGE MONOGRAPHS ON MATHEMATICAL
PHYSICS 2010 page 11
- [27] Von Smoluchowski M.(1914): Studien über Molekularstatistik von Emulsionen
und deren Zusammenhang mit der Brown'schen Bewegung. Sitzungsberichte.
Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Klasse.
CXXIII Band. X Heft. Dez. 1914. Abteilung II A. Wien. pp.2381-2405.
- [28] Warnemünde, Gunnar: BERECHNUNG TURBULENTER KONTINUUMS-
FLUKTUATIONEN 2019 to be published
- [29] Warnemünde, Rolf: Untersuchung von Anregungsfunktionen für Neutronen-
reaktionen nach der Aktivierungsmethode-Universität Hamburg 1969 Diplomar-
beit
- [30] Whittaker E.T. & Watson G.N. A: Course of Modern Analysis Camebridge
Press 1965 page 308
- [31] Weinberg Alwin M., Wigner Eugene P.: THE PHYSICAL THEORY of NEU-
TRON CHAIN REACTORS The University of Chicago Press 1958

BIBLIOGRAPHY

- [32] Wilczek M.: Theory for the single-point velocity statistics of fully developed turbulence- A. Daitche and R. Friedrich February 2011