

# An interesting property of Euler's totient function

Moreno Borrallo, Juan

March 9, 2020

e-mail: [juan.morenoborrallo@gmail.com](mailto:juan.morenoborrallo@gmail.com)

*"Entia non sunt multiplicanda praeter necessitatem" (Ockam, W.)*

*"Dios no juega a los dados con el Universo" (Einstein, Albert)*

*"Te doy gracias, Padre, porque has ocultado estas cosas a los sabios y entendidos y se las has revelado a la gente sencilla" (Mt 11,25)*

## Abstract

In this brief paper it is proved that, for some positive integer  $n$  and some prime number  $q < n$  such that  $\gcd(q, n) = 1$ , it holds that the set  $S = \{x : 0 \leq x \leq n, \gcd(x, qn) = 1\}$  has no less than  $\frac{\varphi(qn)}{q}$  elements for  $n$  being some prime number, and no less than  $\frac{\varphi(qn) + \omega(n) + 1}{q} - \omega(n)$  elements for  $n$  being some composite number.

**2010MSC:** 11A99

# 1 Theorem

Let  $\varphi(n) = n \prod_{p|n} \left(\frac{p-1}{p}\right)$  denote the Euler's totient function, which counts the number of elements of the set  $\{x : 0 \leq x \leq n, \gcd(x, n) = 1\}$ . In this paper it is proved the following

**Theorem.** Let it be some positive integer  $n$ , and some prime number  $q < n$  such that  $\gcd(q, n) = 1$ . Then, it holds that  $S = \{x : 0 \leq x \leq n, \gcd(x, qn) = 1\}$  has no less than  $\frac{\varphi(qn)}{q}$  elements for  $n$  being some prime number, and no less than  $\frac{\varphi(qn) + \omega(n) + 1}{q} - \omega(n)$  elements for  $n$  being some composite number.

## 1.1 Proof for $n$ being some prime number

If  $n = p$ , where  $p$  is some prime number, and  $q < p$ , then to get the elements of  $S$  we need to subtract from  $\varphi(p)$  those numbers that are multiples of  $q$ ; as there are only  $\lfloor \frac{p}{q} \rfloor$  numbers less than  $p$  are relatively prime to  $p$  and not relatively prime to  $qp$ , we have that

$$|S| = \varphi(p) - \lfloor \frac{p}{q} \rfloor$$

As  $q \nmid p$ , we can affirm that

$$\lfloor \frac{p}{q} \rfloor \leq \frac{p-1}{q} = \frac{\varphi(p)}{q}$$

And subsequently we get that

$$|S| \geq \varphi(p) - \frac{\varphi(p)}{q}$$

Operating, we get that

$$|S| \geq \varphi(p) \left(1 - \frac{1}{q}\right)$$

$$|S| \geq \varphi(p) \left(\frac{q-1}{q}\right)$$

As  $\gcd(q, p) = 1$ , and applying the multiplicative properties of  $\varphi(n)$ , we get that

$$\varphi(p) \left(\frac{q-1}{q}\right) = \frac{\varphi(p) \varphi(q)}{q} = \frac{\varphi(qn)}{q}$$

Therefore, for  $n$  being some prime number,

$$|S| \geq \frac{\varphi(qn)}{q}$$

And the theorem is proved for this particular case.

## 1.2 Proof for $n$ being some composite number

If  $n$  is some composite number, then less than  $\lfloor \frac{n}{q} \rfloor$  numbers less than  $n$  are relatively prime to  $n$  and not relatively prime to  $qn$ ; concretely, the multiples of  $q$  and each prime factor of  $n$  could be double-excluded by  $\varphi(n)$  and  $\frac{n}{q}$ , and therefore need to be added once if necessary. Therefore,

$$|S| = \varphi(n) - \lfloor \frac{n}{q} \rfloor + \sum_{p|n} \binom{\lfloor \frac{n}{qp} \rfloor}{1}$$

Where  $\sum_{p|n} \binom{\lfloor \frac{n}{qp} \rfloor}{1}$  counts the common multiples of  $q$  and each prime factor of  $n$ , which already are double excluded by  $\varphi(n)$  and  $\frac{n}{q}$ .

We have that

$$\lfloor \frac{n}{q} \rfloor < \frac{n}{q}$$

$$\sum_{p|n} \binom{\lfloor \frac{n}{qp} \rfloor}{1} \geq \sum_{p|n} \left( \frac{n - (q-1)p}{qp} \right)$$

As

$$\sum_{p|n} \left( \frac{n - (q-1)p}{qp} \right) = \sum_{p|n} \left( \frac{n}{qp} - 1 + \frac{1}{q} \right)$$

Thus, we can affirm that

$$|S| > \varphi(n) - \frac{n}{q} + \sum_{p|n} \binom{\lfloor \frac{n}{qp} \rfloor}{1} - \omega(n) + \frac{\omega(n)}{q}$$

Where  $\omega(n)$  counts the number of distinct prime divisors of  $n$ .

Operating, we get that

$$|S| > \varphi(n) - \frac{n}{q} \left( 1 - \sum_{p|n} \binom{\lfloor \frac{n}{qp} \rfloor}{1} \right) - \omega(n) + \frac{\omega(n)}{q}$$

For  $\omega(n) > 1$ , it is easy to show that

$$\prod_{p|n} \left( \frac{p-1}{p} \right) - \frac{1}{n} \geq 1 - \sum_{p|n} \left( \frac{1}{p} \right)$$

Therefore,

$$|S| > \varphi(n) - \frac{n}{q} \left( \prod_{p|n} \left( \frac{p-1}{p} \right) - \frac{1}{n} \right) - \omega(n) + \frac{\omega(n)}{q}$$

As  $\varphi(n) = n \prod_{p|n} \left( \frac{p-1}{p} \right)$ , we have that

$$|S| > \varphi(n) - \frac{\varphi(n)}{q} + \frac{1}{q} - \omega(n) \left( 1 - \frac{1}{q} \right)$$

Operating,

$$|S| > \varphi(n) \left( 1 - \frac{1}{q} \right) + \frac{1}{q} - \omega(n) + \frac{\omega(n)}{q}$$

As  $\gcd(q, n) = 1$ , and applying the multiplicative properties of  $\varphi(n)$ , we have that

$$\varphi(qn) = \varphi(n) \varphi(q) = \varphi(n) (q-1)$$

Thus,

$$\varphi(n) \left( 1 - \frac{1}{q} \right) + \frac{1}{q} = \frac{\varphi(qn) + 1}{q}$$

Therefore, for  $n$  being some composite number,

$$|S| > \frac{\varphi(qn) + \omega(n) + 1}{q} - \omega(n)$$

And the theorem is proved.