

On the continuity of a functor

Marcelo Carvalho*

*Departamento de Matemática
Universidade Federal de Santa Catarina
Florianópolis, 88.040-900, SC, Brasil*

Abstract

We examine three definitions of continuity of a functor and search for conditions that may fix one of them as general in the sense of including the previous ones. The first definition conceives continuity in terms of inverse systems and their inverse limit. The second definition was developed in the context of abstract Shape theory and considers continuity relative to a fixed functor K and in a general framework that doesn't need the concept of limit. The third definition also considers continuity relative to a fixed functor K , but employs concurrently the concept of limit of a functor. We show how a modification of the third definition allow us to set it as the most general one.

1 Introduction

In category theory there is a standard definition for the limit of a functor $T : \mathcal{C} \rightarrow \mathcal{D}$ that consists of a pair $(\lim T, \lambda^T)$ with $\lim T \in \text{Obj } \mathcal{D}$ and $\lambda^T : (\lim T)_{\mathcal{C}} \rightarrow T$ a natural transformation satisfying a certain universal property (see §4.3). In contrast with the limit concept, continuity has been conceived in different ways through definitions which are not apparently equivalent. In what follows we will focus our attention on a brief discussion on some of those definitions in order to illustrate which elements they have in common and how they differ from each other.

It seems the first attempt to define continuity goes back to the work of Eilenberg and Steenrod [1] (from now on referred to as ES). Considering the category \mathcal{A} of topological pairs (X, A) and maps (between pairs) they characterized a continuous homology functor H_q as the one satisfying $H_q \lim \{(X_m, A_m), p_{mn}\} \simeq \lim H_q \{(X_m, A_m), p_{mn}\}$ for every inverse system $\{(X_m, A_m), p_{mn}\}$ in \mathcal{A} (there is a similar definition for the cohomology functor H^q). Here, by \simeq it is understood the existence of a natural equivalence. A carefull notice on this definition reveals that the limit is being taken on the inverse systems $\{(X_m, A_m), p_{mn}\}$ and $\{(H_q X_m, H_q A_m), H_q(p_{mn})\}$ rather than on the functor H_q itself, then we do not attribute to H_q a pair like $(\lim H_q, \lambda^{H_q})$.

*e-mail: m.carvalho@ufsc.br

Another definition of continuity similar to the ES definition was employed by W. Holsztynski in his construction of a purely categorical version of Borsuk’s Shape theory [2]. Like the ES approach, Holsztynski’s construction still relies on inverse systems, but it is not restricted to the categories of topological pairs and maps as used by ES. It assumes quite arbitrary categories (except for some technical aspects that restrict the type of categories being used, which plays an essential role in Holsztynski’s Shape theory).

It was also in the context of Shape theory, more precisely in the categorical construction given by Bacon [3] and by Cordier and Porter [4], that the concept of continuity became free from the framework of inverse systems. Their treatment of continuity doesn’t use the definition of limit and it is valid in the context of arbitrary categories. In order to establish the continuity of a functor $T : \mathcal{C} \rightarrow \mathcal{D}$ they relied on a class of functors $K : \mathcal{B} \rightarrow \mathcal{C}$ in terms of which it is developed the concept of K –continuity for the functor T . The continuity of T is seen as a particular case of K –continuity for $K = 1_{\mathcal{C}}$.

It was only with K. Hofmann study on the categorical foundations of topological algebras [5] that the concept of continuity was built upon the standard concept of the limit of a functor. Like Bacon’s approach, given a functor $T : \mathcal{C} \rightarrow \mathcal{D}$ Hofmann develops the concept of continuity relative to a functor $K : \mathcal{B} \rightarrow \mathcal{C}$ and assumes it exists the limits of K and TK . Then K –continuity of T is given by means of an isomorphism $T(\lim K) \simeq \lim TK$, which has a form resembling the ES continuity condition, except that here $\lim K$ and $\lim TK$ refers to the standard definition of the limit of a functor rather than to the limit of inverse systems.

The purpose of our study is to examine the definitions of continuity given by Holsztynski, Bacon-Cordier-Porter and Hofmann and search if one of them may be considered as the most general one in the sense of including the two previous ones as particular cases. One should notice that the concept of continuity we consider is a kind of “intrinsic” concept since it doesn’t rely on any topology one may ascribe to the sets $\text{Morf}_{\mathcal{C}}(C, C')$, $\text{Morf}_{\mathcal{D}}(D, D')$ that would allow us to characterize $T : \text{Morf}_{\mathcal{C}}(C, C') \rightarrow \text{Morf}_{\mathcal{D}}(TC, TC')$ as continuous in the topological sense.

Our work is organized as follows. In section 2 we give a brief description of Holsztynski’s definition of continuity of a functor relative to inverse systems. In section 3 we present the elements that were used by Bacon, Cordier and Porter to develop their concept of continuity, which doesn’t depend on any previous limit concept. This approach is able to include the Holsztynski’s treatment if we restrict our attention to inverse systems. In section 4 we present the construction given by Hofmann, who defined continuity using the concept of limit of a functor. This definition also includes Holsztynski’s treatment when restricted to inverse systems. In addition, we show how this definition allows for a modification that includes the Bacon-Cordier-Porter definition.

A word about notation. All functors we deal with are covariant. Given a functor $F : \mathcal{B} \rightarrow \mathcal{C}$ sometimes we write F_{ob} and F_{mo} to denote its action on the objects and morphisms of \mathcal{B} . A morphism $u \in \text{Morf}_{\mathcal{B}}(B, B')$ is written as $u : B \xrightarrow{B} B'$. Whenever we treat with inverse systems $\{X_{\alpha}, p_{\alpha\beta}\}_M, M$

is a pre-ordered set where the indexes run. When we write relations like $p_\alpha = p_{\alpha\beta} p_\beta$, $u_\alpha = p_\alpha h, \dots$ it is assumed they are valid $\forall \alpha \in M, \forall \beta \in M$, observing that $\alpha \leq \beta$ whenever it appears in $p_{\alpha\beta}$, therefore, for ease of notation we omit this information. Finally, we follow the convention to write natural transformations putting a dot over the arrow, e.g. $u : F \dot{\rightarrow} G$ denotes a natural transformation between functors F and G .

2 Holsztynski's definition of continuity

We review the concept of continuity developed by Holsztynski that is restricted to inverse systems over a category. Its content is enounced in the following definition:

2.1 Def.: Continuity for inverse systems

Let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. T is said *continuous for inverse systems* iff

$$T \varprojlim \{X_\alpha, p_{\alpha\beta}\}_M \simeq \varprojlim T\{X_\alpha, p_{\alpha\beta}\}_M \quad (1)$$

$\forall \{X_\alpha, p_{\alpha\beta}\}_M$ inverse system on \mathcal{C} . ■

2.2 Remark: We recall that the limit of an inverse system $\{X_\alpha, p_{\alpha\beta}\}_M$ on a category \mathcal{C} , denoted by $\varprojlim \{X_\alpha, p_{\alpha\beta}\}_M$, is a terminal object in the category $\text{inv}\{X_\alpha, p_{\alpha\beta}\}_M$, i.e. it is an object $\{p_\alpha : X_\infty \xrightarrow{\mathcal{C}} X_\alpha\}_M \in \text{Obj}_{\text{inv}\{X_\alpha, p_{\alpha\beta}\}_M}$ satisfying

i. $p_\alpha = p_{\alpha\beta} p_\beta$ (2)

ii. *The universal property for inverse systems:*

$$\forall \{u_\alpha : W \xrightarrow{\mathcal{C}} X_\alpha\}_M \in \text{Obj}_{\text{inv}\{X_\alpha, p_{\alpha\beta}\}_M} \text{ with } u_\alpha = p_{\alpha\beta} u_\beta, \exists ! h : W \rightarrow X_\infty \text{ with } u_\alpha = p_\alpha h \quad (3)$$

We summarize conditions i, ii in the commutative diagram below

$$\begin{array}{ccc}
 X_\beta & \xrightarrow{p_{\alpha\beta}} & X_\alpha \\
 & \swarrow p_\beta & \searrow p_\alpha \\
 & X_\infty & \\
 u_\beta \swarrow & & \searrow u_\alpha \\
 & W & \\
 & \uparrow h &
 \end{array}$$

Denoting $\{q_\alpha : Y_\infty \xrightarrow{\mathcal{D}} TX_\alpha\}_M = \varprojlim T\{X_\alpha, p_{\alpha\beta}\}_M$, condition (1) establishes an equivalence $\{T(p_\alpha) : TX_\infty \xrightarrow{\mathcal{D}} TX_\alpha\}_M \simeq \{q_\alpha : Y_\infty \xrightarrow{\mathcal{D}} TX_\alpha\}_M$ between terminal objects in $\text{inv}\{TX_\alpha, T(p_{\alpha\beta})\}_M$.

For further use in sections 3 and 4, we write down Holzstynski's definition of projection

2.3: Def.: Projection

Let \mathcal{B} and \mathcal{C} be categories with $\text{Obj}_{\mathcal{B}} = \text{Obj}_{\mathcal{C}}$. A functor $K : \mathcal{B} \rightarrow \mathcal{C}$ is called a projection iff

$K(B) = B, \forall B \in \text{Obj}_{\mathcal{B}}$, and $K : \text{Morf}_{\mathcal{B}}(B, B') \rightarrow \text{Morf}_{\mathcal{C}}(B, B')$ is surjective $\forall B, B' \in \text{Obj}_{\mathcal{B}}$. ■

Then, when $K : \mathcal{B} \rightarrow \mathcal{C}$ is a projection it becomes implicit that we are dealing with categories \mathcal{B} and \mathcal{C} with $\text{Obj}_{\mathcal{B}} = \text{Obj}_{\mathcal{C}}$.

3 Bacon-Cordier-Porter's definition of continuity

Here we review the concept of continuity developed by Bacon, Cordier and Porter. The concept is referred to a fixed functor K and for this reason is called K -continuity. Before presenting it we need to introduce some preliminary concepts. For the definition of comma category and the other concepts see [4].

3.1 Def.: Codomain functor $\delta^{C \downarrow K}$

Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a functor, $C \in \text{Obj}_{\mathcal{C}}$ and $C \downarrow K$ be the comma category of K -objects under C . We define the codomain functor $\delta^{C \downarrow K} : C \downarrow K \rightarrow \mathcal{B}$ as follows

$$\begin{aligned} \delta_{ob}^{C \downarrow K} : \text{Obj}_{C \downarrow K} &\rightarrow \text{Obj}_{\mathcal{B}} \\ (f, B) &\rightarrow \delta_{ob}^{C \downarrow K}(f, B) := B \end{aligned}$$

$$\begin{aligned} \delta_{mo}^{C \downarrow K} : \text{Morf}_{C \downarrow K}((f, B), (f', B')) &\rightarrow \text{Morf}_{\mathcal{B}}(B, B') \\ h &\rightarrow \delta_{mo}^{C \downarrow K}(h) := h \quad \blacksquare \end{aligned}$$

3.2 Def.: $\text{Func}(C \downarrow K, D \downarrow TK)$

Let $K : \mathcal{B} \rightarrow \mathcal{C}$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ be functors and $C \in \text{Obj}_{\mathcal{C}}$, $D \in \text{Obj}_{\mathcal{D}}$. We define $\text{Func}(C \downarrow K, D \downarrow TK)$ as the class having for elements functors $V : C \downarrow K \rightarrow D \downarrow TK$ such that $\delta^{D \downarrow TK} \circ V = \delta^{C \downarrow K}$. ■

From this condition $\text{Func}(C \downarrow K, D \downarrow TK)$ may be characterized in terms of a map V^* as we see in the next result.

3.3 Res.: The condition $\delta^{D \downarrow TK} \circ V = \delta^{C \downarrow K}$ fixes the form of $V \in \text{Func}(C \downarrow K, D \downarrow TK)$ as follows

$$\begin{aligned} V_{ob} : \text{Obj}_{C \downarrow K} &\rightarrow \text{Obj}_{D \downarrow TK} \\ (f, B) &\rightarrow V_{ob}(f, B) := (V^*(f), B) \end{aligned}$$

$$\begin{aligned} V_{mo} : \text{Morf}_{C \downarrow K}((f, B), (f', B')) &\rightarrow \text{Morf}_{D \downarrow TK}((V^*(f), B), (V^*(f'), B')) \\ h &\rightarrow V_{mo}(h) = h \end{aligned}$$

where

$$\begin{aligned} V^* : \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{C}}(C, KB) &\rightarrow \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{D}}(D, TKB) \\ f : C \xrightarrow{\mathcal{C}} KB &\rightarrow V^*(f) : D \xrightarrow{\mathcal{D}} TKB \end{aligned}$$

satisfies

$$\forall f : C \xrightarrow{\mathcal{C}} KB, \forall f' : C \xrightarrow{\mathcal{C}} KB', \exists h : B \xrightarrow{\mathcal{B}} B' \text{ such that } f' = K(h)f \text{ and } V^*(f') = TK(h)V^*(f) \quad (4)$$

Proof: It follows straightforwardly from the condition $\delta^{D\downarrow TK} \circ V = \delta^{C\downarrow K}$. ■

3.4 Remark: Given f, f' the condition $f' = K(h)f$ restricts the form of $h : B \xrightarrow{\mathcal{B}} B'$. The other condition $V^*(f') = TK(h)V^*(f)$ restricts the form of V^* . Generally, they constitute independent conditions, but there are cases when $V^*(f') = TK(h)V^*(f)$ appears as a consequence of having $f' = K(h)f$.

3.5 Def.: $\delta_T : C \downarrow K \rightarrow TC \downarrow TK$

Let $K : \mathcal{B} \rightarrow \mathcal{C}$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ be functors. We define the functor $\delta_T : C \downarrow K \rightarrow TC \downarrow TK$ as

$$\begin{aligned} \delta_{Tob} : \text{Obj}_{C\downarrow K} &\rightarrow \text{Obj}_{TC\downarrow TK} \\ (f, B) &\rightarrow \delta_{Tob}(f, B) := (T(f), B) \end{aligned}$$

$$\begin{aligned} \delta_{Tmo} : \text{Morf}_{C\downarrow K}((f, B), (f', B')) &\rightarrow \text{Morf}_{TC\downarrow TK}((T(f), B), (T(f'), B')) \\ h &\rightarrow \delta_{Tmo}(h) := h \blacksquare \end{aligned}$$

Our next definition associates to every morphism $g : D \xrightarrow{\mathcal{D}} TC$ an induced functor between comma categories.

3.6 Def.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ be functors. Given a morphism $g : D \xrightarrow{\mathcal{D}} TC$, it induces a functor between comma categories $g^* : TC \downarrow TK \rightarrow D \downarrow TK$ defined as follows

$$\begin{aligned} g_{ob}^* : \text{Obj}_{TC\downarrow TK} &\rightarrow \text{Obj}_{D\downarrow TK} \\ (w, B) &\rightarrow g_{ob}^*(w, B) := (wg, B) \end{aligned}$$

$$\begin{aligned} g_{mo}^* : \text{Morf}_{TC\downarrow TK}((w_1, B_1), (w_2, B_2)) &\rightarrow \text{Morf}_{D\downarrow TK}((w_1g, B_1), (w_2g, B_2)) \\ u &\rightarrow g_{mo}^*(u) := u \blacksquare \end{aligned}$$

We are now equipped to define K -continuity of a functor according to Bacon-Cordier-Porter.

3.7 Def.: K -continuity

Let $K : \mathcal{B} \rightarrow \mathcal{C}$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ be functors. We say that T is K -continuous at $C \in \text{Obj}_{\mathcal{C}}$ iff $\forall D \in \text{Obj}_{\mathcal{D}}, \forall V^* : \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{C}}(C, KB) \rightarrow \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{D}}(D, TKB)$ satisfying (4)

$$\exists! g : D \xrightarrow{\mathcal{D}} TC, \forall f : C \xrightarrow{\mathcal{C}} KB : V^*(f) = T(f)g \quad (5)$$

where the last relation is summarized in the commutative diagram

$$\begin{array}{ccc} & D & \\ g \swarrow & & \searrow V^*(f) \\ TC & \xrightarrow{T(f)} & TKB \end{array}$$

We say that T is K -continuous if T is K continuous $\forall C \in \text{Obj}_{\mathcal{C}}$.

3.8 Remark: We notice that the form of $V^*(f)$ given in (5) satisfies condition (4).

We define continuity as a particular case of K -continuity as follows. Let us assume $\mathcal{B} = \mathcal{C}$ and $K = 1_{\mathcal{C}}$, then K -continuity becomes continuity as defined below:

3.9 Def.: Continuous Functor

Let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a covariant functor. T is continuous in $C \in \text{Obj}_{\mathcal{C}}$ iff $\forall D \in \text{Obj}_{\mathcal{D}}, \forall V \in \text{Func}(\mathcal{C} \downarrow 1_{\mathcal{C}}, D \downarrow T), \exists ! g : D \xrightarrow{\mathcal{D}} TC, V = g_T^* \delta_T$.

This condition is equivalent to

$$\begin{aligned} & \forall D \in \text{Obj}_{\mathcal{D}}, \forall V^* : \cup_{C' \in \text{Obj}_{\mathcal{C}}} \text{Morf}_{\mathcal{C}}(C, C') \rightarrow \cup_{C' \in \text{Obj}_{\mathcal{C}}} \text{Morf}_{\mathcal{D}}(D, TC') \text{ satisfying (4)} \\ & \exists ! g : D \xrightarrow{\mathcal{D}} TC, \forall f : C \xrightarrow{\mathcal{C}} C' : V^*(f) = T(f)g \quad \blacksquare \end{aligned} \quad (6)$$

We now examine how the concept of K -continuity implies continuity relative to inverse systems as given in §2.1.

3.10 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a projection. If $T : \mathcal{C} \rightarrow \mathcal{D}$ is K -continuous then T is continuous for inverse systems.

Proof: Let $\{X_{\alpha}, p_{\alpha\beta}\}_M$ be an inverse system in \mathcal{C} and let us assume there is defined the inverse limit $\{p_{\alpha} : X_{\infty} \xrightarrow{\mathcal{C}} X_{\alpha}\}_M = \varprojlim \{X_{\alpha}, p_{\alpha\beta}\}_M$. Given the covariant functor $T : \mathcal{C} \rightarrow \mathcal{D}$ we have that $\{TX_{\alpha}, T(p_{\alpha\beta})\}_M$ is an inverse system in \mathcal{D} and $\{T(p_{\alpha}) : TX_{\infty} \xrightarrow{\mathcal{D}} TX_{\alpha}\}_M$ satisfies

$$T(p_{\alpha}) = T(p_{\alpha\beta})T(p_{\beta}) . \quad (7)$$

Let $\{u_{\alpha} : W \xrightarrow{\mathcal{D}} TX_{\alpha}\}_M$ be such that

$$u_{\alpha} = T(p_{\alpha\beta})u_{\beta} . \quad (8)$$

Since $K : \text{Morf}_{\mathcal{B}}(X_{\beta}, X_{\alpha}) \rightarrow \text{Morf}_{\mathcal{C}}(X_{\beta}, X_{\alpha})$ is a surjection, then for $p_{\alpha\beta} \in \text{Morf}_{\mathcal{C}}(X_{\beta}, X_{\alpha}), \exists q_{\alpha\beta} \in \text{Morf}_{\mathcal{B}}(X_{\beta}, X_{\alpha})$ such that $K(q_{\alpha\beta}) = p_{\alpha\beta}$. But $p_{\alpha} = p_{\alpha\beta} p_{\beta}$ then

$$p_{\alpha} = K(q_{\alpha\beta})p_{\beta} . \quad (9)$$

Now, since $X_{\infty} \in \text{Obj}_{\mathcal{C}}$ and $W \in \text{Obj}_{\mathcal{D}}$ we have that $V \in \text{Func}(X_{\infty} \downarrow K, W \downarrow TK)$ is characterized by a map $V^* : \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{C}}(X_{\infty}, B) \rightarrow \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{D}}(W, TB)$ satisfying (4), which reads as

$$\forall f : X_{\infty} \xrightarrow{\mathcal{C}} B, \forall f' : X_{\infty} \xrightarrow{\mathcal{C}} B', \exists h : B \xrightarrow{\mathcal{B}} B' \text{ with } f' = K(h)f \text{ and } V^*(f') = TK(h)V^*(f) . \quad (10)$$

Consider now a particular choice for V^* such that for $f \equiv p_{\beta}, f' \equiv p_{\alpha} (\alpha \leq \beta)$ we have $V^*(p_{\alpha}) = u_{\alpha}, V^*(p_{\beta}) = u_{\beta}$. That this choice exists it is readily seen if we take $h \equiv q_{\alpha\beta}$ satisfying $K(q_{\alpha\beta}) = p_{\alpha\beta}$ for in this case we have that (8) and (9) guarantee (10). Since V is continuous, from (5) we have that

$$\exists ! g : W \xrightarrow{\mathcal{D}} TX_{\infty}, \forall p_{\alpha} : X_{\infty} \xrightarrow{\mathcal{C}} X_{\alpha}, V^*(p_{\alpha}) = T(p_{\alpha})g .$$

i.e.

$$u_\alpha = T(p_\alpha)g. \quad (11)$$

From (7), (8) and (11) the family $\{T(p_\alpha) : TX_\infty \xrightarrow{\mathcal{D}} TX_\alpha\}_M$ satisfies the analogue of conditions (2) and (3) relative to the inverse system $\{TX_\alpha, T(p_{\alpha\beta})\}_M$, therefore

$$\{T(p_\alpha) : TX_\infty \xrightarrow{\mathcal{D}} TX_\alpha\}_M = \varprojlim \{TX_\alpha, T(p_{\alpha\beta})\}_M$$

i.e. $T : \mathcal{C} \rightarrow \mathcal{D}$ is continuous for inverse systems. ■

3.11 Remark: As we have seen, the definition of K -continuity of $T : \mathcal{C} \rightarrow \mathcal{D}$ assumes the existence of a unique morphism g , but does not specify the conditions for this morphism to exist. However, in the result just proved, the existence and uniqueness of g stated in §3.7 follows as a consequence of existing the inverse limit of $\{TX_\alpha, T(p_{\alpha\beta})\}_M$.

3.12 Res.: Continuity implies continuity for inverse systems

Let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. If T is continuous then T is continuous for inverse systems.

Proof. It follows directly from §3.9 since continuity is a particular case of K -continuity taking $K = 1_{\mathcal{C}}$. ■

4 Hofmann's definition of continuity

Here we analyze the concept of continuity developed by Hofmann. The concept is established relative to a functor previously fixed and employs the standard definition of limit.

First we introduce the concept of constant functor induced by an object.

4.1 Def.: Let \mathcal{C} and \mathcal{D} be categories and $D \in \text{Obj}_{\mathcal{D}}$. We define a constant functor $D_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{D}$ as follows

$$D_{\mathcal{C}ob} : \text{Obj}_{\mathcal{C}} \rightarrow \text{Obj}_{\mathcal{D}}$$

$$C \rightarrow D_{\mathcal{C}ob} := D$$

$$D_{\mathcal{C}mo} : \text{Morf}_{\mathcal{C}} \rightarrow \text{Morf}_{\mathcal{D}}$$

$$h : C \xrightarrow{\mathcal{C}} C' \rightarrow D_{\mathcal{C}mo}(h) := 1_D \quad \blacksquare$$

Given a functor it induces a natural transformation as follows:

4.2 Def.: Given categories \mathcal{C} and \mathcal{D} and a morphism $F : D \xrightarrow{\mathcal{D}} D'$ we define a natural transformation

$$F_{\mathcal{C}} : D_{\mathcal{C}} \rightarrow D'_{\mathcal{C}}$$

$$F_{\mathcal{C}} : \text{Obj}_{\mathcal{C}} \rightarrow \text{Morf}_{\mathcal{D}}$$

$$C \rightarrow F_{\mathcal{C}}(C) := F \quad \blacksquare$$

We recall the standard definition of the limit of a functor:

4.3 Def.: Limit of a functor

Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a functor. The limit of K consists of a pair $(\lim K, \lambda^K)$ with $\lim K \in \text{Obj}_{\mathcal{C}}$ and $\lambda^K : (\lim K)_{\mathcal{B}} \rightarrow K$ such that $\forall C \in \text{Obj}_{\mathcal{C}}, \forall \alpha : C_{\mathcal{B}} \rightarrow K, \exists! \bar{\alpha} : C \xrightarrow{\mathcal{C}} \lim K$ such that $\lambda^K \bar{\alpha}_{\mathcal{B}} = \alpha$. ■

Condition $\lambda^K \bar{\alpha}_{\mathcal{B}} = \alpha$ may be expressed in terms of the commutative diagram below ($\forall B \in \text{Obj}_{\mathcal{B}}$):

$$\begin{array}{ccc} & C & \\ \bar{\alpha} \swarrow & & \searrow \alpha^{(B)} \\ \lim K & \xrightarrow{\lambda^{K(B)}} & KB \end{array}$$

We say that λ^K is the *limit morphism* and $\lim K$ is the *limit object*. As an abuse of notation we write $\lim K$ as a shorthand for the pair $(\lim K, \lambda^K)$.

We need a preliminary result:

4.4 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ be functors and let us assume that $\exists \lim K$. Then, $T \circ (\lim K)_{\mathcal{B}} = [T(\lim K)]_{\mathcal{B}}$.

Proof: It follows directly from definition 4.1. ■

The next result guarantees the existence of a unique functor T_K provided it exists the limits of K and TK .

4.5 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ be functors. If $\exists \lim K, \exists \lim TK$ then $\exists! T_K : T(\lim K) \xrightarrow{\mathcal{D}} \lim TK$ such that $\lambda^{TK} T_K = T \lambda^K$, i.e. the diagram below is commutative

$$\begin{array}{ccc} [T(\lim K)]_{\mathcal{B}} & \xrightarrow{T_{K\mathcal{B}}} & (\lim TK)_{\mathcal{B}} \\ & \searrow T \lambda^K & \swarrow \lambda^{TK} \\ & TK & \end{array}$$

Proof: Since it exists $\lim K$ it follows there is a natural transformation $\lambda^K : (\lim K)_{\mathcal{B}} \rightarrow K$. Using §4.4 we consider the natural transformation $T \lambda^K : [T(\lim K)]_{\mathcal{B}} \rightarrow TK$. Since it also exists $\lim TK$ there is a natural transformation $\lambda^{TK} : (\lim TK)_{\mathcal{B}} \rightarrow TK$ satisfying:

$$\forall \beta : D_{\mathcal{B}} \rightarrow TK, \exists! \bar{\beta} : D \xrightarrow{\mathcal{D}} \lim TK \text{ such that } \lambda^{TK} \bar{\beta}_{\mathcal{B}} = \beta. \quad (12)$$

Identifying in (12): $D \equiv T(\lim K)$ and $\beta \equiv T \lambda^K$ we have that $\exists! \bar{\beta} : T(\lim K) \xrightarrow{\mathcal{D}} \lim TK$ such that $\lambda^{TK} \bar{\beta}_{\mathcal{B}} = T \lambda^K$. We identify the morphism T_K with $\bar{\beta}$ and this ends our proof. ■

We are now equipped to define K -continuity.

4.6 Def.: K -continuity (according to Hofmann)

Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a functor. The functor $T : \mathcal{C} \rightarrow \mathcal{D}$ is K -continuous iff

- i. $\exists \lim K \Rightarrow \exists \lim TK$
- ii. $T_K : T(\lim K) \rightarrow \lim TK$ is an isomorphism ■

We need a preliminary result:

4.7 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a projection such that $\exists \lim K$. For every inverse system on \mathcal{C} , $\{X_\alpha, p_{\alpha\beta}\}_M$, it exists $\{p_\alpha : \lim K \xrightarrow{\mathcal{C}} X_\alpha\}_M$ such that $\{p_\alpha : \lim K \xrightarrow{\mathcal{C}} X_\alpha\}_M = \varprojlim \{X_\alpha, p_{\alpha\beta}\}_M$.

Proof: Let $\{X_\alpha, p_{\alpha\beta}\}_M$ be an inverse system in \mathcal{C} . If K is a projection then it exists $q_{\alpha\beta} : X_\beta \xrightarrow{\mathcal{B}} X_\alpha$ with $K(q_{\alpha\beta}) = p_{\alpha\beta}$. If it exists $\lim K$ we have the following conditions i, ii, iii:

i. There is a natural transformation $\lambda^K : (\lim K)_\mathcal{B} \rightarrow K$ such that for the $q_{\alpha\beta} : X_\beta \xrightarrow{\mathcal{B}} X_\alpha$ we have $\lambda^K(X_\alpha) = K(q_{\alpha\beta})\lambda^K(X_\beta)$, i.e.

$$\lambda^K(X_\alpha) = p_{\alpha\beta}\lambda^K(X_\beta). \quad (13)$$

ii. For any $\beta : W_\mathcal{B} \rightarrow K$ we also have $\beta(X_\alpha) : W \xrightarrow{\mathcal{C}} X_\alpha$ satisfies

$$\beta(X_\alpha) = p_{\alpha\beta}\beta(X_\beta). \quad (14)$$

iii. There is a unique $\bar{\beta} : W \xrightarrow{\mathcal{C}} \lim K$ such that

$$\lambda^K(X_\alpha)\bar{\beta} = \beta(X_\alpha). \quad (15)$$

Then, if we identify $p_\alpha = \lambda^K(X_\alpha)$ and $u_\alpha \equiv \beta(X_\alpha)$ we notice that relations (13), (14), (15) correspond to the conditions (2) and (3). Then $\{p_\alpha : \lim K \xrightarrow{\mathcal{C}} X_\alpha\}_M = \varprojlim \{X_\alpha, p_{\alpha\beta}\}_M$. ■

We will now examine how K -continuity implies continuity relative to inverse systems.

4.8 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a projection. If $\exists \lim K$ and $T : \mathcal{C} \rightarrow \mathcal{D}$ is K -continuous then T is continuous for inverse systems.

Proof: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a projection and assume it exists $\lim K$. Let $T : \mathcal{C} \rightarrow \mathcal{D}$ be K -continuous. By definition $\exists \lim TK$, $\exists T_K : T(\lim K) \xrightarrow{\mathcal{D}} \lim TK$, which is an isomorphism. Let $\{X_\alpha, p_{\alpha\beta}\}_M$ be an inverse system on \mathcal{C} and let us assume there is defined the inverse limit.

Since $K : \mathcal{B} \rightarrow \mathcal{C}$ is a projection then for $p_{\alpha\beta} : X_\beta \xrightarrow{\mathcal{C}} X_\alpha$ we have $q_{\alpha\beta} : X_\beta \xrightarrow{\mathcal{B}} X_\alpha$ such that $K(q_{\alpha\beta}) = p_{\alpha\beta}$.

If it exists $\lim K$ we have defined a natural transformation $\lambda^K : (\lim K)_\mathcal{B} \rightarrow K$ such that $\lambda^K(X_\alpha) : \lim K \xrightarrow{\mathcal{C}} X_\alpha$ satisfies $\lambda^K(X_\alpha) = p_{\alpha\beta}\lambda^K(X_\beta)$. From 4.7 we identify $\{\lambda^K(X_\alpha) : \lim K \xrightarrow{\mathcal{C}} X_\alpha\}_M = \varprojlim \{X_\alpha, p_{\alpha\beta}\}_M$.

The functor T determines an inverse system $\{TX_\alpha, T(p_{\alpha\beta})\}_M$ in \mathcal{D} and $T(\lambda^K(X_\alpha)) : T \lim K \xrightarrow{\mathcal{D}} TX_\alpha$ satisfies

$$T(\lambda^K(X_\alpha)) = T(p_{\alpha\beta})T(\lambda^K(X_\beta)). \quad (16)$$

If it exists $\lim TK$ we have defined a natural transformation $\lambda^{TK} : (\lim TK)_\mathcal{B} \rightarrow TK$ such that $\lambda^{TK}(X_\alpha) : \lim TK \xrightarrow{\mathcal{D}} TX_\alpha$. Given $\beta : W_\mathcal{B} \rightarrow TK$ we consider $\beta(X_\alpha) : W \xrightarrow{\mathcal{D}} TX_\alpha$, which satisfies

$$\beta(X_\alpha) = T(p_{\alpha\beta})\beta(X_\beta). \quad (17)$$

Then $\exists! \bar{\beta} : W \xrightarrow{\mathcal{D}} \lim TK$ with $\lambda^{TK}(X_\alpha)\bar{\beta} = \beta(X_\alpha)$.

From 4.5 T_K satisfies $\lambda^{TK}(X_\alpha)T_K = T(\lambda^K(X_\alpha))$ and since T_K is isomorphism we obtain $\lambda^{TK}(X_\alpha) = T(\lambda^K(X_\alpha))T_K^{-1} \therefore \lambda^{TK}(X_\alpha)\bar{\beta} = T(\lambda^K(X_\alpha))T_K^{-1}\bar{\beta}$, which becomes

$$\beta(X_\alpha) = T(\lambda^K(X_\alpha))T_K^{-1}\bar{\beta}. \quad (18)$$

Then considering (16), (17) and (18) we have shown that $\{T(\lambda^K(X_\alpha)) : T \lim K \xrightarrow{\mathcal{D}} TX_\alpha\}$ satisfies the properties (2) and (3) relative to the inverse system $\{TX_\alpha, T(p_{\alpha\beta})\}$, therefore

$$\{T(\lambda^K(X_\alpha)) : T \lim K \xrightarrow{\mathcal{D}} TX_\alpha\} = \varprojlim \{TX_\alpha, T(p_{\alpha\beta})\}$$

i.e. T is continuous for inverse systems. ■

We now propose a modification of Hofmann's definition in order to relate it to the Bacon-Cordier-Porter's definition.

4.9 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a functor. The functor $T : \mathcal{C} \rightarrow \mathcal{D}$ is K -continuous iff

i. $\exists \lim K \Rightarrow \exists \lim TK$

ii. $T_K : T(\lim K) \rightarrow \lim TK$ is an isomorphism

iii. $\forall \alpha : C_{\mathcal{B}} \rightarrow K, \forall \beta : D_{\mathcal{B}} \rightarrow TK, \exists ! \chi : D \xrightarrow{\mathcal{D}} TC$ such that $T_K^{-1}\bar{\beta} = T(\bar{\alpha})\chi$

i.e. the diagram below is commutative

$$\begin{array}{ccc} D & \xrightarrow{\bar{\beta}} & \lim TK \\ \chi \downarrow & & \downarrow T_K^{-1} \\ TC & \xrightarrow{T(\bar{\alpha})} & T(\lim K) \end{array}$$

4.10 Res.: Let $K : \mathcal{B} \rightarrow \mathcal{C}$ be a functor such that $\exists \lim K$. If $T : \mathcal{C} \rightarrow \mathcal{D}$ is K -continuous according to definition 4.9 then T is K -continuous according to the Bacon-Porter-Cordier definition.

Proof: Let $C \in \text{Obj}_{\mathcal{C}}, D \in \text{Obj}_{\mathcal{D}}$ and consider a map

$$V^* : \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{C}}(C, KB) \rightarrow \cup_{B \in \text{Obj}_{\mathcal{B}}} \text{Morf}_{\mathcal{D}}(D, TKB)$$

satisfying (4). In order to show that T is K -continuous we must verify (5). Then let us assume $f : C \xrightarrow{\mathcal{C}} KB, f' : C \xrightarrow{\mathcal{C}} KB'$ satisfying $f' = K(h)f$ and $V^*(f') = TK(h)V^*(f)$ for a certain $h : B \xrightarrow{\mathcal{B}} B'$.

Since $\exists \lim K$ let us consider a natural transformation $\alpha : C_{\mathcal{B}} \rightarrow K$ satisfying $\alpha(B) = f, \alpha(B') = f'$. Then for the given $h : B \xrightarrow{\mathcal{B}} B'$ we have $\alpha(B') = K(h)\alpha(B)$. We also have $\lambda^K(B)\bar{\alpha} = \alpha(B)$ for a unique $\bar{\alpha} : C \xrightarrow{\mathcal{C}} \lim K$.

Since $\exists \lim TK$ let us consider a natural transformation $\beta : D_{\mathcal{B}} \rightarrow TK$ satisfying $\beta(B) = V^*(f), \beta(B') = V^*(f')$. Then, for the same $h : B \xrightarrow{\mathcal{B}} B'$ previously considered we have $\beta(B') = TK(h)\beta(B)$. In addition we also have the condition $\lambda^{TK}(B)\bar{\beta} = \beta(B)$ for a unique $\bar{\beta} : D \xrightarrow{\mathcal{D}} \lim TK$.

The K -continuity of T provide us with an isomorphism $T_K : T(\lim K) \xrightarrow{\mathcal{D}} \lim TK$ satisfying

$\lambda^{TK}(B) = T(\lambda^K(B))T_K^{-1}$, then $\lambda^{TK}(B)\bar{\beta} = T(\lambda^K(B))T_K^{-1}\bar{\beta}$ and from §4.9.iii $\exists! \chi : D \xrightarrow{\mathcal{D}} TC$ such that $\lambda^{TK}(B)\bar{\beta} = T(\lambda^K(B))T(\bar{\alpha})\chi = T(\alpha(B))\chi$ i.e. $\beta(B) = T(\alpha(B))\chi$, or in a more suitable notation: $V^*(f) = T(f)\chi$. Then we have proven (5) and this shows that T is K -continuous according to the Bacon-Cordier-Porter definition. ■

5 Conclusion

The concept of continuity of a functor is not consensually established and in our work we have focused our attention on three definitions. The first one, due to Holsztynski, deals with inverse systems and arises in the context of Shape theory. The second one, due to Bacon, Cordier and Porter, was also developed in the context of Shape theory and employs a framework that depends neither on inverse systems nor on the limit concept. The third definition, due to Hofmann, defines continuity using the concept of limit and seeks application in topological algebras. In its original form none of them includes simultaneously the other two as particular cases.

It is only with the addition of condition §4.9iii to Hofmann's original definition that this modified form becomes a general definition including the two previous ones.

It is not clear what elements could be added to or even what modifications could be made on the Bacon-Cordier-Porter definition in order to make it the most general one. From the analysis of sections 3 and 4 we observe that for $f : C \xrightarrow{\mathcal{C}} KX_\alpha$ the key point is to harmonize equations $V^*(f) = T(f)g$ (see (5)) with $V^*(f) = T(\lambda^K(X_\alpha))T_K^{-1}\bar{\beta}$, which is the farther we can go in §4.10 without imposing condition §4.9iii. Such attempt, if possible, may demand some modifications on the definitions given in section §3, which consequently would affect the content of the Shape theory formulated by Bacon-Cordier-Porter leading to a modified form. The analysis of the properties of this modified Shape theory deserve some investigation.

Acknowledgement: This work is done in honour of $\overline{\text{IHHP}}$, $\overline{\text{TC}}$, $\overline{\text{INA}}$.

References

- [1] S. Eilenberg and N. Steenrod, *Foundations of Algebraic Topology*, Princeton University Press, 1952.
- [2] W. Holsztynski, *An extension and axiomatic characterization of Borsuk's theory of shape*, *Fundamenta Mathematicae* **70** (1971), no. 2, 157-168.
- [3] P. Bacon, *Axiomatic Shape theory*, *Proc. Am. Math. Soc.* **53** (1975), no. 2, 489-496.
- [4] J.-M. Cordier and T. Porter, *Shape Theory*, Dover, 2008.
- [5] K. H. Hofmann, *Category theoretical methods in topological algebras*, *Lecture Notes in Mathematics*, vol 540 (1976) 345-403.